

King's Research Portal

DOI:

[10.1017/S0033291720000574](https://doi.org/10.1017/S0033291720000574)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Lukito, S. D. A. W., Norman, L. J., Carlisi, C. O., Radua, J., Hart, H. C., Simonoff, E., & Rubia, K. (2020). Comparative Meta-analyses of Brain Structural and Functional Abnormalities during Cognitive Control in Attention-Deficit/Hyperactivity Disorder and Autism Spectrum Disorder. *Psychological Medicine*, 50(6), 894-919. <https://doi.org/10.1017/S0033291720000574>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

TITLE PAGE

Comparative Meta-analyses of Brain Structural and Functional Abnormalities during Cognitive Control in Attention-Deficit/Hyperactivity Disorder and Autism Spectrum Disorder

Steve Lukito^{1*}, PhD, Luke Norman^{1,2,3}, PhD, Christina Carlisi^{1,4}, PhD, Joaquim Radua^{5,6,7}, MD, PhD, Heledd Hart¹, PhD, Emily Simonoff¹, MD & Katya Rubia¹, PhD*

¹ Department of Child and Adolescent Psychiatry; Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK

² Department of Psychiatry, University of Michigan, Ann Arbor, Michigan, USA

³ The Social and Behavioral Research Branch, National Human Genome Research Institute, National Institute of Health, Bethesda, Maryland, USA

⁴ Division of Psychology and Language Sciences, University College London, UK

⁵ Department of Psychosis Studies; Institute of Psychiatry, Psychology and Neuroscience; King's College London; UK

⁶ Imaging of mood- and anxiety-related disorders (IMARD) group, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), CIBERSAM, Barcelona, Spain

⁷ Centre for Psychiatric Research and Education, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

* Correspondence and reprints request:

Dr Steve Lukito

Department of Child and Adolescent Psychiatry; Institute of Psychiatry, Psychology and Neuroscience, King's College London, 16 De Crespigny Park, London, SE5 8AF, UK

Word count: 4718

Abstract: 250

ABSTRACT

Background: People with attention-deficit/hyperactivity disorder (ADHD) and autism spectrum disorder (ASD) have abnormalities in frontal, temporal, parietal and striato-thalamic networks. It is unclear to what extent these abnormalities are distinctive or shared. This comparative meta-analysis aimed to identify the most consistent disorder-differentiating and shared structural and functional abnormalities.

Methods: Systematic literature search was conducted for whole-brain voxel-based morphometry (VBM) and fMRI studies of cognitive control comparing people with ASD or ADHD with typically developing controls. Regional gray matter volume (GMV) and fMRI abnormalities during cognitive control were compared in the overall sample and in age-, sex- and IQ-matched subgroups with seed-based d mapping meta-analytic methods.

Results: Eighty-six independent VBM (1533 ADHD and 1295 controls; 1445 ASD and 1477 controls) and 60 fMRI datasets (1001 ADHD and 1004 controls; 335 ASD and 353 controls) were identified. The VBM meta-analyses revealed ADHD-differentiating *decreased* ventromedial orbitofrontal ($z = 2.22, p < .0001$) but ASD-differentiating *increased* bilateral temporal and right dorsolateral prefrontal GMV ($zs \geq 1.64, ps \leq .002$). The fMRI meta-analyses of cognitive control revealed ASD-differentiating medial prefrontal underactivation but overactivation in bilateral ventrolateral prefrontal cortices and precuneus ($zs \geq 1.04, ps \leq .003$). During motor response inhibition specifically, ADHD relative to ASD showed right inferior fronto-striatal underactivation ($zs \geq 1.14, ps \leq .003$) but shared right anterior insula underactivation.

Conclusions: People with ADHD and ASD have mostly distinct structural abnormalities, with enlarged fronto-temporal GMV in ASD and reduced orbitofrontal GMV in ADHD; and mostly distinct functional abnormalities, which were more pronounced in ASD.

TEXT

1. Introduction

Attention-deficit/hyperactivity disorder (ADHD) and autism spectrum disorder (ASD) are distinct neurodevelopmental conditions. ADHD is characterized by age-inappropriate symptoms of inattention, hyperactivity and impulsivity, whereas ASD is characterized by difficulties in social interaction/communication and stereotypical repetitive behavior (American Psychiatric Association (APA), 2013). The estimated prevalence of ADHD (5-7%) is higher than ASD (1-2%) worldwide (Elsabbagh et al., 2012; Polanczyk, Willcutt, Salum, Kieling, & Rohde, 2014). In both disorders, brain structural abnormalities have been observed in frontal, temporal, parietal and striato-thalamic networks (Ecker, 2017; Norman et al., 2016), although the extent of their overlap is poorly understood. Furthermore, both disorders are associated with cognitive control deficits (F. Craig et al., 2016), although ASD relative to ADHD seem less severely impaired (see meta-analyses Lipszyc & Schachar, 2010; Willcutt, Sonuga-Barke, Nigg, & Sergeant, 2008) and display more heterogeneous performance (Geurts, van den Bergh, & Ruzzano, 2014; Kuiper, Verhoeven, & Geurts, 2016). Exploring abnormalities in brain structure and cognitive control function could help elucidate the disorder-differentiating and shared difficulties in ASD and ADHD.

In ADHD, brain abnormalities are thought to represent delayed maturation in brain structure and function mediating late-developing cognitive functions such as cognitive control (Hoogman et al., 2017; Shaw et al., 2007; Sripada, Kessler, & Angstadt, 2014).

Correspondingly, previous voxel-based morphometry (VBM) meta-analyses have revealed reduced gray matter volume (GMV) in ventromedial orbitofrontal cortex (vmOFC) and basal ganglia (BG) (Frodl & Skokauskas, 2012; Nakao, Radua, Rubia, & Mataix-Cols, 2011; Norman et al., 2016). In addition, mega-analytic studies by the ENIGMA consortium of subcortical and cortical structure involving over 1700 and 2200 individuals with ADHD,

respectively, found reduced GMV in BG, amygdala and hippocampus (Hoogman et al., 2017), as well as reduced cortical surface areas and thickness in several frontal, temporal and parietal regions (Hoogman et al., 2019).

Converging with these findings are meta-analytic reports of underactivation in lateral and medial frontostriatal networks in ADHD relative to controls during cognitive control such as in right inferior frontal gyrus (IFG)/anterior insula (AI), supplementary motor area (SMA)/anterior cingulate cortex (ACC), and the striatum (Hart, Radua, Nakao, Mataix-Cols, & Rubia, 2013; McCarthy, Skokauskas, & Frodl, 2014; Norman et al., 2016). The role of right IFG as potential biomarker of ADHD in particular has been discussed in several studies and a meta-analysis showing disorder-specificity of underactivation in right and/or left IFG during cognitive control in ADHD children relative to pediatric bipolar disorder (Passarotti, Sweeney, & Pavuluri, 2010), obsessive-compulsive disorder (Norman et al., 2016) and conduct disorder (Rubia, 2018; Rubia, Halari, et al., 2010).

In ASD, findings from neuroimaging, post-mortem and head circumference development studies have led to the “early brain overgrowth theory” (Courchesne, Campbell, & Solso, 2011; Lange et al., 2015; Redcay & Courchesne, 2005; Schumann et al., 2010), which, however, has been contested in recent studies (Ecker, Schmeisser, Loth, & Murphy, 2017; Raznahan et al., 2013). Meta-analytic and mega-analytic findings have varied but GMV reduction in the striatum and amygdala/hippocampus were among the more frequently reported (Cauda et al., 2011; DeRamus & Kana, 2015; Nickl-Jockschat et al., 2012; van Rooij et al., 2018; Via, Radua, Cardoner, Happe, & Mataix-Cols, 2011). Another recent meta-analysis of VBM studies in over 900 ASD patients, however, primarily found cortical abnormalities such as GMV decrease in medial prefrontal cortex (mPFC) and posterior insula; and increase in left anterior temporal, right inferior temporo-parietal, left dorsolateral prefrontal (dlPFC) and precentral cortices (Carlisi et al., 2017). Furthermore, the ENIGMA

mega-analysis has found cortical thickness increase in frontal and decrease in temporal areas (van Rooij et al., 2018).

The whole-brain meta-analysis of fMRI studies of cognitive control in ASD (Carlisi et al., 2017) has found underactivation in salience and executive network areas such as mPFC, left dorsolateral and right ventrolateral prefrontal cortices (vlPFC)/AI, left cerebellum, inferior parietal lobe (IPL) and right premotor area; and overactivation in right temporo-parietal regions including default mode network (DMN) areas.

While these meta-analytic findings suggest partially overlapping brain abnormalities in ASD and ADHD, only direct comparisons can elucidate their differences and commonalities. Among such studies, a VBM study showed reduced GMV in ADHD relative to ASD in right posterior cerebellum and ASD-differentiating reduced left middle/superior temporal gyri (M/STG) (Lim et al., 2015). Furthermore, fMRI studies have shown ASD-differentiating mPFC underactivation during reward reversal (Chantiluke, Barrett, Giampietro, Brammer, Simmons, Murphy, et al., 2015), shared dlPFC underactivation during sustained attention and working memory (Chantiluke, Barrett, Giampietro, Brammer, Simmons, & Rubia, 2015; Christakou et al., 2013); and shared precuneus abnormalities during sustained attention, reward reversal and the resting state (Chantiluke, Barrett, Giampietro, Brammer, Simmons, Murphy, et al., 2015; Christakou et al., 2013; Di Martino et al., 2013). Importantly, during successful motor inhibition, ADHD-differentiating underactivation was observed in left vlPFC and BG and ASD-differentiating overactivation in bilateral IFG (Chantiluke, Barrett, Giampietro, Santosh, et al., 2015). Few imaging studies, however, have directly compared the two disorders and their relatively small sample sizes have limited statistical power.

We therefore conducted meta-analyses of structural and functional abnormalities related to functions that are commonly impaired in ASD and ADHD, i.e., cognitive control. The aim of the study was to establish the most consistently disorder-differentiating and shared structural

and functional deficits, which is important for developing disorder-specific or transdiagnostic treatment. Guided by previous comparative studies and meta-analyses in ASD and/or ADHD, we hypothesized that these disorders are characterized by both shared abnormalities in salience, cognitive control, and default mode networks as well as disorder-differentiating impairments. Structurally, we hypothesized ADHD-differentiating reduced GMV in BG/insula relative to ASD (Lim et al., 2015; Norman et al., 2016), and ASD-differentiating increased GMV in frontal and temporo-parietal regions (Carlisi et al., 2017; Lim et al., 2015). During cognitive control, we expected ADHD-differentiating reduced activation in right IFG and BG relative to ASD (Chantiluke, Barrett, Giampietro, Santosh, et al., 2015; Norman et al., 2016; Passarotti et al., 2010; Rubia et al., 2009), ASD-differentiating underactivation relative to ADHD in the medial frontal part of the cognitive control network (Carlisi et al., 2017; Chantiluke, Barrett, Giampietro, Brammer, Simmons, Murphy, et al., 2015), and shared abnormal overactivation in precuneus in both disorders (Chantiluke, Barrett, Giampietro, Brammer, Simmons, Murphy, et al., 2015; Christakou et al., 2013; Di Martino et al., 2013).

2. Methods

2.1. Publication Search and Study Inclusion

Systematic literature search was conducted for peer-reviewed English language publications in PubMed, Scopus, Web of Science and ScienceDirect until 30th November 2018, for whole-brain fMRI or VBM studies in ASD and ADHD. Manual search was conducted in reference lists of previous meta-analyses. Included studies compared the ASD or ADHD groups against typically developing (TD) controls on GMV or on cognitive-control fMRI BOLD signal from stop-signal, go/no-go, switch, Stroop, Simon and flanker, using predefined inhibitory contrasts (Supplement 1). Only whole-brain neuroimaging data were included, to prevent bias towards specific neuroanatomy (Friston, Rotshtein, Geng, Sterzer, & Henson, 2006). Excluded studies used region of interest (ROI) approaches or involved <10 patients, which

were deemed lacking in statistical power (Desmond & Glover, 2002; Nakao et al., 2011; Thirion et al., 2007). Studies that potentially report duplicate data from other publications (including mega-analyses), have no TD controls, or have provided no peak coordinates, were excluded. When samples overlapped, the largest sample was included. Mean age, proportion of males, mean IQ, cognitive control tasks, current and lifetime psychostimulant use (i.e., proportion of participants being prescribed the medication) and effect size and coordinate location of peaks for regional GMV and activation differences were extracted from each study and authors were contacted for missing information. Reports of exclusion or inclusion of people with co-occurring ASD or ADHD in the counterpart disorder groups were also extracted from the literature. Data were extracted from all studies by SL, 64% of which were also extracted by LN and CC who achieved 100% agreement. This meta-analysis was not pre-registered in a time-stamped, institutional registry prior to the research being conducted.

2.2. Meta-Analyses

The anisotropic seed-based *d* Mapping (AES-SDM) meta-analytic software (www.sdmproject.com; Supplement 1) employed in previous meta-analyses of ASD and ADHD (Carlisi et al., 2017; Frodl & Skokauskas, 2012; McCarthy et al., 2014; Norman et al., 2016) was used for the present voxel-wise meta-analyses. The software can accommodate statistical parametric maps or peak coordinates and effect sizes (*t*-scores) data. For the latter, AES-SDM computes signed (positive/negative) effect sizes and variance maps of brain regional differences between clinical and control groups by convolving an anisotropic non-normalized Gaussian kernel with Hedges effect size of each peak. Voxels correlated with the peak values are assigned higher effect sizes. Maps are then combined across studies based on random-effect model, accounting for sample size, within-study variability and between-study heterogeneity. Correlated datasets (e.g., when the same group of participants completed several cognitive tasks) were included in the meta-analysis as a

single set (Norman et al., 2016), adjusted for shared variance of brain activation or structure across datasets. The Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines were followed (Stroup et al., 2000).

Age, sex and IQ were compared across patient and control groups prior to meta-analyses in STATA14 (StataCorp, 2015), using sample-size weighted *F*-statistics. First, VBM meta-analyses were conducted in each disorder relative to TD controls and then between the two disorders. Second, equivalent fMRI meta-analyses were conducted to examine the neural impairments observed during cognitive control within and between disorders using all available data. To explore specific cognitive-control subconstructs (e.g., Bari & Robbins, 2013), we conducted fMRI sub-meta-analyses of prepotent motor response inhibition, including studies employing the go/no-go and stop-signal tasks only. No other cognitive control subconstructs were explored due to insufficient data. To find the most consistent disorder-differentiating neural abnormalities regardless of demographic characteristics, we conducted sensitivity meta-analyses in an age-, sex- and IQ-matched subgroup, instead of covarying group differences in these variables that could result in false positives (Dennis et al., 2009; Miller & Chapman, 2001). An iterative algorithm computed group differences in age, sex and IQ for all possible combinations of studies, selecting the largest subgroups with statistically nonsignificant group differences in the three variables (Carlisi et al., 2017). Findings surviving the sensitivity analyses in the matched subgroups are the only ones reported and discussed.

Furthermore, conjunction analyses were used to assess overlapping fMRI and GMV abnormalities within and across disorders (using threshold $p < .005$). To assess consistency of findings across age groups, supplementary age-stratified sub-meta-analyses were conducted in overall pediatric or adult age groups and in their respective age-, sex- and IQ-matched subgroups, where possible (Supplements 3 and 4). These were complemented with a series of sample-size weighted correlational analyses between the mean sample age

and extracted effect sizes of brain abnormalities averaged over 10-mm radius spheres centered at the peaks of disorder-differentiating or shared abnormalities, applying Bonferroni multiple comparison correction (Supplement 5). Non-parametric correlations were chosen due to the bimodal distribution of age among studies. Following previous meta-analyses (Hart et al., 2013; Norman et al., 2016), psychostimulant effects were explored in separate meta-regressions between brain structure/function and lifetime psychostimulant exposure, i.e., proportions of samples who have ever been exposed to psychostimulants, or current psychostimulant exposure, in brain regions that are found impaired in ADHD (Supplement 6). Supplementary meta-analyses covarying for task types (motor response inhibition, cognitive interference, motor switching, combination of tasks) and behavioral task performance (impaired, unimpaired) were also conducted to assess their impacts on disorder-differentiating findings (Supplement 7).

A statistical threshold $p < .005$ was used for all meta-analyses (Radua & Mataix-Cols, 2009; Radua, Mataix-Cols, et al., 2012), and a reduced threshold $p < .0005$ and a cluster extent 20 voxels was used in the meta-regressions to control for false positives (Radua, Borgwardt, et al., 2012). Egger's tests assessed potential publication bias in the disorder-differentiating and shared findings. Jack-knife sensitivity analyses examined replicability of findings through iterative whole-brain meta-analyses, leaving one dataset out at a time (Supplement 8; Radua & Mataix-Cols, 2009).

3. Results

3.1. Search Results and Sample Characteristics

The systematic literature search identified 140 VBM and fMRI articles of cognitive control in ASD and ADHD. Accounting for datasets of same clinical participants and, where possible, separating data from pediatric and adult participants resulted in 86 independent VBM datasets (38 datasets including 1533 ADHD participants and 1295 controls; 48 datasets

involving 1445 ASD participants and 1477 controls) and 60 independent cognitive-control fMRI datasets (42 datasets involving 1001 ADHD participants and 1004 controls; 18 datasets involving 335 ASD participants and 353 controls) (Tables 1-2). Groups did not differ in age in VBM, $F(3, 83) = 1.24$, $p = .30$, and fMRI datasets, $F(3, 56) = .68$, $p = .57$, but did in sex in VBM, $F(3, 83) = 6.03$, $p = .0009$, and fMRI datasets, $F(3, 56) = 5.93$, $p = .0014$; and in IQ in VBM, $F(3, 67) = 16.2$, $p < .0001$ and fMRI datasets $F(3, 45) = 15.5$, $p < .0001$. See Table 3 for demographic characteristics.

3.2. Disorder-Differentiating and Shared Brain Structure Abnormalities

3.2.1. ADHD VBM

Relative to TD controls, ADHD had reduced GMV in vmOFC/vmPFC/rostral ACC (rACC), extending into right caudate, and in right putamen/posterior insula/STG, left precingulate gyrus (pre-CG), right rostrolateral PFC and left vlPFC/STG/temporal pole (Table 4A-i; Figure 1A-i). Age-stratified analyses showed reduced GMV in right STG/putamen/posterior insula, right caudate and rACC/ventromedial PFC (vmPFC) among pediatric participants and vmOFC/subcallosal gyrus in both age groups (Supplements 3,4). Current and lifetime psychostimulant exposures were associated with increased vmOFC GMV (Supplement 6).

3.2.2. ASD VBM

Relative to TD controls, ASD had GMV reduction in dACC/dorsomedial PFC (dmPFC), left cerebellum, right hippocampus/PHG/FFG and dorsomedial thalamus; and enhancement in left anterior inferior/middle/superior temporal gyri (I/M/STG)/posterior insula, bilateral precuneus/posterior cingulate cortex (PCC) and right middle/superior frontal gyrus (M/SFG) (Table 4A-ii; Figure 1A-ii). Most abnormalities in the overall sample, i.e., in left cerebellum, dorsomedial thalamus, precuneus/PCC and right MFG/SFG/dlPFC GMV were also found

among pediatric participants, and enhanced left anterior STG/MTG GMV was found in both age subgroups (Supplements 3,4).

3.2.3. ADHD vs. ASD VBM

People with ADHD, relative to ASD, had consistently disorder-differentiating reduced vmOFC/rACC/left caudate GMV; while people with ASD, relative to ADHD, showed increased GMV in left anterior STG/MTG, right MFG/SFG/dIPFC, and right posterior MTG/STG (Table 4A-iii,iv; Figure 1A-iii,iv). Reduced vmOFC GMV was most consistently ADHD-differentiating in adults, while the medial prefrontal GMV reduction in rACC/dmPFC was more apparent among pediatric participants (Supplements 3,4).

3.3. Disorder-Differentiating and Shared Brain Abnormalities during Cognitive Control

3.3.1. ADHD fMRI

During cognitive control, ADHD relative to TD controls, showed underactivation in right thalamus/caudate, left MTG/STG/superior temporal pole, bilateral SMA/dmPFC, left IFG/AI/temporal pole, right AI/putamen, left postcingulate gyrus (post-CG), left MFG/dIPFC and right MTG (Table 4B-i, Figure 1B-i). Age-stratified analyses showed underactivated dmPFC clusters and left post-CG among pediatric participants, in all regions but the dmPFC and left post-CG in adults; and in left M/STG/temporal pole in both groups (Supplements 3,4). Lifetime psychostimulant exposure was positively associated with activation in left IFG (Supplement 6).

During motor response inhibition specifically, underactivation was found in ADHD relative to TD controls in left MFG/dIPFC, left anterior MTG/STG, left post-CG, right vIPFC/OFC/AI, right IFG, and right caudate (Table 4C-i, Figure 1C-i). Age-stratified analyses showed that underactivation in left MTG/STG was apparent among pediatric participants, underactivation

in left MFG/dIPFC was apparent in adults, while underactivation in right vIPFC/AI was apparent in both age groups (Supplements 3,4).

3.3.2. ASD fMRI

During cognitive control, ASD relative to TD controls showed underactivation in ACC/midcingulate/dmPFC, left MFG/dIPFC, right MFG/dIPFC, left IPL and left lingual gyrus/cerebellum; and overactivation in left precuneus/midcingulate, right inferior occipital gyrus (IOG), left vIPFC/OFC, left MFG/rostrolateral PFC and right IFG (Table 4B-ii, Figure 1B-ii). Underactivation in rdACC/dmPFC and left IPL; and overactivation in precuneus, left vIPFC/OFC and left MFG/dIPFC were found in adults (Supplements 3,4). No pediatric sub-meta-analyses were conducted due to insufficient fMRI data.

During motor response inhibition, underactivation was found in ASD relative to TD controls in right AI/vIPFC, left cerebellum, right MFG/dIPFC and right PCC/precuneus; while overactivation was found in left vIPFC/OFC and right IOG/fusiform gyrus (FFG)/ITG (Table 4C-ii; Figure 1C-ii). No age-stratified sub-meta-analyses were conducted due to insufficient data.

3.3.3. ADHD vs. ASD fMRI

During cognitive control, ASD-differentiating underactivation was found in rACC/midcingulate/dmPFC and left MFG/dIPFC; and overactivation was found in precuneus, right IOG/FFG, right IFG and left vIPFC/OFC (Table 4B-iii,iv; Figure 1B-iii,iv). The age-stratified analysis, which was conducted only in adults due to available data, found ASD-differentiating underactivation relative to ADHD in rdACC/dmPFC and overactivation in precuneus and left vIPFC/OFC (Supplement 4). No ADHD-differentiating abnormalities were found.

During motor response inhibition, ADHD-differentiating underactivation was found in right caudate and IFG. Furthermore, ASD-differentiating underactivation was found in left lingual gyrus/FFG/cerebellum, right precuneus and right MFG/dIPFC; while overactivation was found in right IOG/FFG, left vIPFC/OFC and left MFG/SFG/dIPFC (Table 4C-iii,iv; Figure 1C-iii,iv). Underactivation in right AI (MNI coordinates: 40, 20, -6; 51 voxels) was shared between disorders.

3.5. Multimodal VBM and fMRI Analyses

In ADHD relative to TD controls, overlapping reduced GMV and fMRI underactivation was found during cognitive control in right caudate nucleus (MNI coordinates: 10, 10, 8; 194 voxels). In ASD relative to TD controls, reduced GMV and fMRI underactivation was found in dACC/mPFC (MNI coordinates: 0, 40, 16; 575 voxels).

3.6. Controlling for Task Type and Performance in fMRI Meta-Analysis

The disorder-differentiating impairment during cognitive control was unchanged after covarying for task types and performance. During motor response inhibition, ADHD-differentiating underactivation in right IFG and caudate did not survive covarying for task performance (Supplement 7).

3.7. Publication Bias and Jack-Knife Reliability Findings

Egger tests suggest no significant publication bias for the structural, $ts(84) = .32-2.29$, $ps \geq .1$, and functional abnormalities during cognitive control, $ts(58) = .04-1.90$, $ps \geq .38$, and during motor response inhibition specifically, $ts(40) = .41-1.37$, $ps \geq .99$ (Bonferroni adjusted p -values). Jack-knife reliability analyses suggest robust disorder-differentiating findings (Supplement 8).

4. Discussion

We aimed to elucidate the most consistent disorder-differentiating and shared brain abnormalities in ASD and ADHD. The findings revealed predominantly disorder-differentiating abnormalities, particularly striking in the VBM meta-analysis, where we found ADHD-differentiating *reduced* GMV in vmOFC; and ASD-differentiating *increased* GMV in bilateral temporal and right lateral prefrontal cortices. In fMRI, the findings overall showed predominantly ASD-differentiating abnormalities, including underactivation in medial frontal and overactivation in bilateral ventrolateral regions and precuneus during cognitive control. During motor response inhibition specifically, ADHD-differentiating underactivation was in right IFG and caudate, while ASD had differentiating underactivation mostly in posterior regions, and overactivation in left dorsal and ventral frontal regions (Figure 2). Both disorders shared underactivation in right AI. The findings overall suggest that people with ADHD and ASD have mostly distinct brain abnormalities.

The ADHD-differentiating decreased GMV in vmOFC relative to ASD may be related to common reports of reward-based decision-making neural network dysfunctions in ADHD (e.g., Chantiluke et al., 2014; Cubillo, Halari, Smith, Taylor, & Rubia, 2012; Rubia, 2018; Tegelbeckers et al., 2018), and supports the hypothesis of distinctive reward processing in ASD and ADHD (Kohls et al., 2014; van Dongen et al., 2015). Reduced vmOFC GMV in ADHD compared to TD controls extends relatively recent meta-analytic findings (McGrath & Stoodley, 2019; Norman et al., 2016), and there have been correlational reports between OFC GMV reduction and increased ADHD symptoms in large-scale general population studies (Albaugh et al., 2017; Fuentes et al., 2012; Korponay et al., 2017). Interestingly, our age-stratified results suggest that the ADHD-differentiating deficit features more consistently in adulthood, when co-occurring addiction behavioral problems increase (e.g., Breyer et al., 2009; Ortal et al., 2015), more so in ADHD than in ASD (e.g., Sizoo et al., 2010; Solberg et al., 2019).

The increased left anterior and right posterior temporal GMV in ASD is a consistent VBM meta-analytic finding in ASD relative to TD controls (Carlisi et al., 2017; Cauda et al., 2011; DeRamus & Kana, 2015; Duerden, Mak-Fan, Taylor, & Roberts, 2012). Enhanced left anterior GMV was the only mega-analytic impairment finding in over 400 people with ASD but it did not survive in a smaller age- and sex-matched subgroups (Riddle, Cascio, & Woodward, 2017). Our meta-analysis may have increased statistical power to detect the differences. The ASD-differentiating increased left temporal GMV extends findings from a small VBM ASD/ADHD comparative study (Lim et al., 2015). The left temporal lobe plays roles in semantic and language processing (Binder et al., 2011), while the right posterior temporoparietal cortices plays a role in social interaction and mentalizing, i.e., the ability to attribute mental states in others (Krall et al., 2015). These structures are important for social cognition, which has been thought to be an ASD-specific impairment (Kana et al., 2015; Lombardo, Chakrabarti, Bullmore, & Baron-Cohen, 2011; White, Frith, Rellecke, Al-Noor, & Gilbert, 2014).

The ASD-differentiating increased right dlPFC GMV has not converged in smaller meta-analyses (Carlisi et al., 2017; DeRamus & Kana, 2015), possibly reflecting heterogeneity in findings. Right dlPFC is important for cognitive control, working memory and cognitive flexibility (Szczepanski & Knight, 2014). Neuronal and grey matter overgrowth have been reported in pediatric ASD in small ROI-based neuroimaging and post-mortem studies (Carper & Courchesne, 2005; Courchesne, Mouton, et al., 2011; Mitchell et al., 2009), but they have not been replicated in wider age range of ASD cases with cognitive flexibility deficits (Griebeling et al., 2010).

The disorder-differentiating GMV *increase* in ASD and *decrease* in ADHD may possibly be related to the disorders' contrasting developmental trajectories (Courchesne, Campbell, et al., 2011; Shaw et al., 2007). However, while most of our ADHD-differentiating findings showed reduction in structure and function that may possibly reflect delayed maturation in

ADHD, the ASD-differentiating enhanced clusters were not found consistently in the pediatric sub-meta-analysis. Recent findings suggest that early brain overgrowth is not a defining characteristic of toddlers with high-risk for developing ASD (Hazlett et al., 2017), and may be specific to boys with regressive autism (Nordahl et al., 2011), who are underrepresented across studies. Thus, our findings are unlikely to reflect early brain overgrowth.

In the fMRI meta-analyses, ASD-differentiating impairments were predominant medial and left middle prefrontal underactivation and in bilateral ventrolateral prefrontal overactivation during cognitive control, while ADHD-differentiating underactivation emerged during motor inhibition in right inferior frontal and striatal regions. The ADHD-differentiating underactivation in right IFG and caudate, key regions implicated in inhibitory control (Rae, Hughes, Weaver, Anderson, & Rowe, 2014; Zhang, Geng, & Lee, 2017), are consistent with previous meta-analytic findings in ADHD during cognitive control (Hart et al., 2013; McCarthy et al., 2014; Norman et al., 2016) and extends the role of these regions, in particular right IFG, as disorder-specific neurofunctional biomarker of ADHD, as it has previously been observed relative to bipolar, obsessive compulsive and conduct disorders (Norman et al., 2016; Passarotti et al., 2010; Rubia, 2018; Rubia, Cubillo, et al., 2010; Rubia, Halari, et al., 2010; Rubia et al., 2009).

The mPFC underactivation in ASD is a consistent meta-analytic finding during cognitive control and related functions (Carlisi et al., 2017; Di Martino et al., 2009; Philip et al., 2012). The region is also implicated in functions such as sensory, emotion and social processing domains typically impaired in ASD (Kana, Patriquin, Black, Channell, & Wicker, 2016; Martinez-Sanchis, 2014; South & Rodgers, 2017). Intriguingly, the mPFC underactivation was ASD-differentiating relative to ADHD. While both medial and lateral PFC are activated during motor inhibition (Rae et al., 2014; Zhang et al., 2017), the activation of mPFC are more reliably tapped by cognitive interference tasks thus it may have closer associations to change or conflict detection (Aron, Herz, Brown, Forstmann, & Zaghoul, 2016). Whether our

findings represent dissociated frontal brain functional impairment in ASD from ADHD in sub-constructs of cognitive control should thus be tested further meta-analytically when more fMRI findings are available.

While both disorders show functional abnormalities in prefrontal cortex in both hemispheres, the ASD-differentiating atypical activation was more predominantly on the left, in contrast to the right-lateralized ADHD-differentiating fronto-striatal underactivation, although atypical brain lateralization is not uncommon finding in ASD (Dichter, 2012; Floris & Howells, 2018; Kleinmans, Muller, Cohen, & Courchesne, 2008). Most importantly, comparisons between each disorder relative their respective TD controls show that neural impairment in ADHD is overwhelmingly characterized by fronto-striato-temporal underactivation, which is in line with previous findings of a developmental lag of cognitive control networks (Sripada et al., 2014), whereas the impairments in ASD implicate both under- and overactivation clusters in executive, attentional and, even, fronto-parieto-occipital visuo-perceptual brain regions. Of note, abnormally enhanced recruitment of posterior brain regions in ASD has been observed during working memory (Koshino et al., 2005; Vogan et al., 2014), psychomotor vigilance (Christakou et al., 2013; Murphy et al., 2014) and temporal discounting (Chantiluke et al., 2014; Norman et al., 2017). This thus suggests a complex pattern of neurofunctional abnormality in ASD, with compromised cortical specialization, possibly reflecting both neural impairments inherent to ASD and, possibly, ensuing neuronal compensation (see review Floris & Howells, 2018).

The ASD-differentiating overactivation in PCC/precuneus, which was also unimpaired in ADHD relative to their respective controls, was unexpected, however. The overactivation in precuneus/PCC in ASD, which has been shown in a variety of tasks and resting-state condition (Christakou et al., 2013; Kennedy, Redcay, & Courchesne, 2006; Murdaugh et al., 2012; Spencer et al., 2012), presumably indicates DMN dysregulation. The lack of overactivation in PCC/precuneus in ADHD during cognitive control were apparent in

previous meta-analyses (McCarthy et al., 2014; Norman et al., 2016), and could potentially be related to psychostimulant exposure which has shown to normalize DMN functioning (Liddle et al., 2011; Peterson et al., 2009; Rubia et al., 2014).

Finally, shared functional abnormality between disorders was found during motor response inhibition only in right AI, which is part of the salience network (Menon & Uddin, 2010). The limited shared abnormality may seem at odds with typical observation of psychiatric comorbidities in clinical settings, but individuals with ASD and ADHD are typically selectively recruited into imaging studies, excluding specific psychiatric comorbidities, which could have emphasized the differences between disorders, at the cost of their representativeness.

On the other hand, the findings of disorder differences are more likely to be underestimated. First, typically stringent statistical corrections applied by individual studies could have concealed true group differences in coordinate-based meta-analyses. The ADHD-differentiating findings, particularly, are likely underestimated by co-occurring ADHD in the ASD groups due to past preclusion of ADHD diagnosis in ASD (APA, 2000). Studies attempting to include more representative samples of the disorders may have included comorbid ASD and ADHD cases, perhaps at higher rates in ASD than in ADHD, as shown in prevalence studies in community representative samples (Gjevik, Eldevik, Fjæran-Granum, & Sponheim, 2011; Hollingdale, Woodhouse, Young, Fridman, & Mandy, 2019; Salazar et al., 2015; Simonoff et al., 2008). Long-term medication exposure is also likely to have masked the ADHD-differentiating findings.

Other limitations of the meta-analyses include the fewer number of participants with ASD relative to ADHD, which could have reduced power for detecting small ASD-differentiating impairments and increased the probability of false positive findings, particularly in the fMRI meta- and sub-meta-analyses. The representativeness of the meta-analytic findings may also be limited by the fact that many studies, particularly in ADHD, investigated males

specifically. In ASD, the majority of studies have focused on adults and predominantly high-functioning individuals only (with few exceptions, i.e., Cai et al., 2018; Contarino et al., 2016; Retico et al., 2016; Riva et al., 2013; Wang et al., 2017; Q. Yang et al., 2018), even though individuals with learning disability made up a significant proportion of the ASD population (e.g., Charman et al., 2011). Furthermore, brain-behavior correlations could not be examined due to variations of disorder trait measures across studies. Finally, the influence of task discrepancies across studies could not be fully assessed due to limited data for the non-motor cognitive control tasks.

5. Conclusions

These comparative meta-analyses of brain structure and function show predominantly disorder-differentiating abnormalities in the form of *decreased* GMV in vmOFC reward processing regions in ADHD and *increased* GMV in ASD in frontal and temporal cortices, part of the central executive and social cognition networks. Disorder-differentiating fMRI deficits were predominantly observed in ASD in medial frontal executive, ventrolateral prefrontal and DMN regions when including all cognitive control tasks; and in ADHD in inferior fronto-striatal regions during motor response inhibition only. Shared functional abnormality was limited to right AI during motor response inhibition. Therefore, people with ADHD and ASD appear to have mostly distinct brain structure and cognitive-control functional abnormalities. The findings contribute to the elucidation of the differential brain abnormalities in the two disorders, which is important for the understanding of their underlying pathophysiology and could ultimately aid in the development of future, disorder-differentiated behavioral, pharmacological or neurotherapeutic treatments.

STATEMENTS

Acknowledgement: We thank all authors of the included studies who provided additional data for this meta-analysis. We also thank Toby Wise for insightful discussions on the statistical methods used in this meta-analysis.

Financial support: SL and LN were supported by Medical Research Council and Institute of Psychiatry, Psychology, and Neuroscience (IoPPN) PhD Excellence awards. CC is supported by a Wellcome Trust Sir Henry Wellcome Postdoctoral Fellowship (206459/Z/17/Z). JR receives support from the Instituto de Salud Carlos III and co-funded by European Union (ERDF/ESF, 'Investing in your future', Miguel Servet Research Contract CPII19/00009 and Research Project Grant PI19/00394). HH was supported by a grant from the Lila Weston Foundation for Medical Research and the Kids Company to KR. ES and KR receive research support from the National Institute of Health Research (NIHR) Biomedical Research Centre at South London and Maudsley Foundation Trust (IS-BRC-1215-20018). ES furthermore is supported by the NIHR through a program grant (RP-PG-1211-20016) and Senior Investigator Award (NF-SI-0514-10073), the European Union Innovative Medicines Initiative (EU-IMI 115300), Autistica (7237) Medical Research Council (MR/R000832/1, MR/P019293/1), the Economic and Social Research Council (ESRC 003041/1) and Guy's and St Thomas' Charitable Foundation (GSTT EF1150502) and the Maudsley Charity. KR and SL are currently supported by an MRC Grant MR/P012647/1 to KR. This study is supported by funding from the UK Department of Health via the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) for Mental Health at South London and the Maudsley National Health Service (NHS) Foundation Trust and the IoPPN, King's College London, which also support both senior authors, ES and KR. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Conflicts of interest: KR has received grants from Lilly and Shire pharmaceuticals and speaker's bureau from Shire, Lilly and Medice. Other authors report no biomedical financial interests or potential conflicts of interest.

Ethical standards: The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

FIGURE LEGENDS

Fig. 1. Brain abnormalities in the ADHD and ASD groups. Abnormalities in the (A) gray matter volume and brain activations during (B) cognitive control and (C) motor response inhibition. Rows (i) and (ii) show abnormalities in ADHD and ASD relative to typically developing (TD) controls. Abnormalities in ADHD versus ASD, each relative to TD controls are shown in (iii) overall samples and (iv) age-, sex-, and IQ-matched subgroups. In the overall sample, shared reduced rACC/mPFC GMV (MNI coordinates: 0,46,12; 119 voxels), and subthreshold overactivation in ADHD and underactivation in ASD in dmPFC (MNI coordinates: -2, 34, 36; 82 voxels) were observed but did not survive sensitivity analysis. A statistical threshold of $p < .005$ with a cluster extent of 20 voxels were used in all analyses.

Fig. 2. Summary of consistent disorder-differentiating and shared brain abnormalities in ADHD and ASD. Findings of (A) gray matter volume abnormalities and brain activation abnormalities during (B) cognitive control and (C) motor response inhibition. A statistical threshold of $p < .005$ with a cluster extent of 20 voxels were used in all analyses.

TABLES

Table 1. Sample Characteristics and Summary Findings of VBM and fMRI Cognitive Control Studies in ADHD

Source	Age group	Task	Patients			Controls			Summary findings
			N (% male)	Age, y	IQ	N (% male)	Age, y	IQ	
(A) VBM studies in ADHD									
Ahrendts et al. (2011)	Adult	--	31 (65)	31.2	n/a	31 (65)	31.5	n/a	ADHD<TD: L/R occipital lobe
Montes et al. (2010)	Adult	--	20 (50)	29.0	102.9	20 (50)	27.6	100.2	ADHD<TD: R caudate
Amico, Stauber, Koutsouleris, and Frodl (2011)	Adult	--	20 (75)	33.6	n/a	20 (75)	34.7	n/a	--
Bonath, Tegelbeckers, Wilke, Flechtner, and Krauel (2016)	Pediat.	--	18 (100)	13.6	106.8	18 (100)	14.1	108.1	ADHD<TD: L/R cerebellum, occipital cortex, L/R hippocampus/amygdala complex, L ACC
Bralten et al. (2016)	Mixed	--	307 (68)	17.1	97.1	196 (49)	16.7	106.6	ADHD<TD: L pre-CG, L OFC, R frontal pole, L paracingulate/cingulate cortex/frontal pole, bilateral medial FG/paracingulate/cingulate/ subcallosal cortices
Brieber et al. (2007)	Pediat.	--	15 (100)	13.1	104.1	15 (100)	13.3	107.7	ADHD<TD: L/R caudate nucleus, L/R hippocampus, R MFG, L insula, L STG, L MOG; ADHD>TD: L/R SPL, post-CG, L middle cingulate, precuneus, L IPL
Carmona et al. (2005)	Pediat.	--	25 (84)	10.8	n/a	25 (84)	11.2	n/a	ADHD<TD: L pre-/post-CG, paracentral lobule, R rectal gyrus, R inferior and superior orbital frontal, L/R cerebellum
Depue, Burgess, Bidwell, Willcutt, and Banich (2010)	Adult	--	31 (61)	20.0	114.2	21 (65)	19.3	112.6	--
Gehricke et al. (2017)	Adult	--	32 (81)	25.3	n/a	40 (83)	23.9	n/a	ADHD<TD: R M/IFG, R S/MTG, L caudate head, L PHG
He et al. (2015)	Pediat.	--	37 (100)	9.9	n/a	35 (100)	10.7	n/a	ADHD<TD: R OFC, R PMC, L PCC, L posterior midcingulate cortex
Iannaccone, Hauser, Ball, et al. (2015)	Pediat.	--	18 (50)	14.5	114.5	18 (61)	14.8	108.5	ADHD<TD: R medial SFG/ACC/SFG, L SMA/cingulate gyrus, L cerebellum; ADHD>TD: R pre- and post-CG
Jagger-Rickels, Kibby, and Constance (2018)	Pediat.	--	41 (44)	9.6	n/a	32 (56)	9.7	n/a	ADHD<TD: L MTG/STS, R STS L/R caudate/ putamen, L/R SFG, L/R insula, L/R thalamus, R medial OFC, L MFG, L/R occipital lobe, L medial SFG, L pre-CG, R post cingulate/anterior lingual, R calcarine/cuneus, L supramarginal gyrus
Johnston et al. (2014)	Pediat.	--	34 (100)	12.5	99.8	34 (100)	13.2	103.7	ADHD<TD: Brainstem, putamen
Kappel et al. (2015)	Pediat.	--	14 (71)	9.8	104.6	10 (80)	11.0	111.9	ADHD<TD: R STG, R Heschl's gyrus, R rolandic operculum; ADHD>TD: L paracentral lobule, L/R middle orbital gyrus, R FFG, L rectal gyrus

	Adult	--	16 (94)	23.5	97.8	20 (100)	23.7	108.4	ADHD<TD: R supramarginal gyrus, R precuneus, L OFG, R hippocampus, L rectal gyrus
Kaya et al. (2018)	Pediat.	--	19 (74)	10.3	113.5	18 (67)	10.2	119.7	ADHD>TD: L/R SFG, R MFG, L/R SMA, L/R pre-CG, L post-CG, R MOG, L cuneus
Kobel et al. (2010)	Pediat.	--	14 (100)	10.4	n/a	12 (100)	10.9	n/a	ADHD<TD: R S/MTG
Kumar, Arya, and Agarwal (2017)	Pediat.	--	18 (100)	9.6	92.1	18 (100)	9.7	109.7	ADHD<TD: L OFC, L MFG/ dIPFC, L MTG, L cerebellum
Li et al. (2015)	Pediat.	--	30 (100)	10.3	121.7	30 (100)	10.3	107.1	ADHD<TD: R insula, R OFC
Lim et al. (2015)	Pediat.	--	44 (100)	13.6	92.2	33 (100)	14.3	110	ADHD<TD: L/R cerebellum, L IPL, pre-/post-CG
Maier et al. (2015)	Adult	--	131 (48)	34.5	113.1	95 (47)	37.7	121	--
McAlonan et al. (2007)	Pediat.	--	28 (100)	9.9	109.9	31 (100)	9.6	116.5	ADHD<TD: R MFG, L/R SFG/IFG, R globus pallidus/putamen/thalamus, precuneus, L IPL, L SOG, R cerebellum vermis
Moreno-Alcázar et al. (2016)	Adult	--	44 (66)	31.6	105.0	44 (66)	32.6	106.0	ADHD<TD: R SMA/SFG, subgenual ACC, R DLPFC
Onnink et al. (2014)	Adult	--	119 (39)	36.3	107.5	107 (42)	36.9	110.2	--
Overmeyer et al. (2001)	Pediat.	--	18 (83)	10.4	99	16 (94)	10.3	n/a	ADHD<TD: R PCC, R SFG, L/R putamen/globus pallidus
Ramesh and Rai (2013)	Pediat.	--	15 (26)	16.8	n/a	15 (26)	16.7	n/a	ADHD<TD: L cingulate gyrus, R mid-cingulate gyrus, L SFG, L/R medial SFG, L temporal lobe, L MOG, L cuneus, L STG, L supramarginal gyrus
Roman-Urrestarazu et al. (2016)	Adult	--	49 (76)	22.2	96.6	34 (50)	22.9	112.2	ADHD<TD: L/R caudate
Saad et al. (2017)	Pediat.	--	16 (75)	12.8	n/a	28 (68)	13.1	n/a	--
	Pediat.	--	18 (72)	13.7	n/a	28 (68)	13.1	n/a	--
Sasayama et al. (2010)	Pediat.	--	18 (72)	10.6	90	17 (71)	10.0	n/a	ADHD<TD: L/R amygdala/temporal pole/OFC, L/R occipital cortex, R STS, L parietal cortex, L MFG, L temporal pole, R rectal gyrus, L PHG
Seidman et al. (2011)	Adult	--	74 (51)	37.3	116	54 (46)	34.3	115.8	--
Sethi et al. (2017)	Adult	--	30 (63)	33.7	109.0	30 (63)	32.6	110.1	ADHD<TD: R IPC
Shimada et al. (2015)	Pediat.	--	17 (88)	10.3	95.3	15 (73)	12.8	104.1	ADHD<TD: L putamen
Stevens and Haney-Caron (2012)	Pediat.	--	24 (67)	15.7	n/a	24 (70)	16.0	n/a	--
van Wingen et al. (2013)	Adult	--	14 (100)	32.0	104	15 (100)	37.0	99	ADHD<TD: R putamen, R cerebellum; ADHD>TD: L/R midbrain, R pre-CG
Vilgis, Sun, Chen, Silk, and Vance (2016)	Pediat.	--	48 (100)	12.6	92.2	31 (100)	12.8	109.6	ADHD<TD: R SPL, L IPL, L/R S/MFG, medial FG, R MTG
Villemonteix et al. (2015)	Pediat.	--	33 (55)	10.3	105.6	24 (50)	10.0	109.7	ADHD<TD: R insula, R MTG,
	Pediat.	--	20 (80)	10.4	107.4	24 (50)	10.0	109.7	ADHD<TD: R MFG, R pre-CG
Wang, Jiang, Cao, and Wang (2007)	Pediat.	--	12 (100)	13.4	n/a	12 (100)	13.5	n/a	ADHD<TD: L SPL, R MFG, R MTG, R BG; ADHD>TD: R occipital lobe

P. Yang et al. (2008)	Pediat.	--	57 (61)	11.1	97.9	57 (60)	11.7	n/a	ADHD<TD: L/R caudate, L/R cerebellum, R calcarine sulcus, L cuneus
(B) fMRI cognitive control studies in ADHD									
Banich et al. (2009)	Adult	Stroop	23 (61)	20.0	116	23 (57)	19.0	113	ADHD>TD: R MFG
Bhaijiwala, Chevrier, and Schachar (2014)	Pediat.	Stop	12 (58)	13.8	n/a	12 (50)	15.4	n/a	ADHD<TD: R MFG, R MTG, L cerebellum; ADHD>TD: R MFG/SFG
Booth et al. (2005)	Pediat.	GNG	12 (67)	11.0	n/a	12 (58)	11.7	n/a	ADHD<TD: L/R pre-CG, L/R caudate, R caudate head, R IFG, L/R thalamus.
Carmona et al. (2012)	Adult	GNG	19 (100)	33.6	110.9	19 (100)	29.4	111.7	ADHD<TD: R STG
Chantiluke, Barrett, Giampietro, Santosh, et al. (2015)	Pediat.	Stop	18 (100)	14.3	95	25 (100)	13.4	109	ADHD<TD: L OFC/STL/putamen/GP, L IPL
Chen et al. (2015)	Adult	GNG	29 (100)	24.9	n/a	25 (100)	25.6	n/a	--
Chou, Chia, Shang, and Gau (2015)	Pediat.	Stroop	42 (81)	10.5	108.5	20 (80)	12.0	106.5	ADHD<TD: R MFG/IFG (atomoxetine group pre-treatment), L IFG, L SPL, R medial FG (methylphenidate group pre-treatment); ADHD>TD: L/R SPL, pre-CG, L dIPFC, L dACC (atomoxetine group pre-treatment), post-CG (methylphenidate group pre-treatment)
Congdon et al. (2010)	Adult	Stop	35 (54)	30.9	n/a	62 (45)	30.8	n/a	--
Cubillo et al. (2010) ^a	Adult	Stop, Switch	11 (100)	29.0	92	14 (100)	28.0	106	ADHD<TD: L IFC/AI/PMC, R IFC/AI/caudate/thalamus/ putamen/PMC, L/R SMA/ACC (stop), R IFG/insula/caudate/ putamen L IFG/insula/ putamen/pre-/post-CG/inferior parietal (switch)
Cubillo et al. (2014)	Pediat.	Stop	19 (100)	13.1	92	29 (100)	13.8	110	ADHD<TD: L/R IFG, L M/ITG/parietal gyri, R cerebellum/FFG; ADHD>TD: L cerebellum, L/R PCC/occipital gyri, R STG/post-CG/posterior insula/ putamen
Cubillo, Halari, Giampietro, Taylor, and Rubia (2011) ^a	Adult	Simon	11 (100)	29.0	92	15 (100)	28.0	112	ADHD<TD: L OFC/IFC/mPFC/ACC/ caudate/PMC
Dibbets, Evers, Hurks, Marchetta, and Jolles (2009) ^b	Adult	GNG	16 (100)	28.9	n/a	13 (100)	28.1	n/a	--
Dibbets, Evers, Hurks, Bakker, and Jolles (2010) ^b	Adult	Switch	15 (100)	28.8	n/a	14 (100)	28.6	n/a	ADHD<TD: R putamen, R dorsal posterior CG, L mPFC, R thalamus, L OFC/ claustrum/post-CG; ADHD>TD: R MTG, R dACC, R precuneus, R lingual gyrus, L pre-CG/SMA, L insula
Durston, Mulder, Casey, Ziermans, and van Engeland (2006)	Pediat.	GNG	11 (100)	14.0	100	11 (100)	15.3	106	--
L. Y. Fan, Chou, and Gau (2017) ^c	Adult.	Stroop	12 (42)	28.9	115.8	12 (42)	30.3	118.3	ADHD>TD: R IFG, RACC
	Adult	Stroop	12 (42)	32.5	119.9	12 (42)	30.3	118.3	ADHD>TD: R IFG
L. Y. Fan, Shang, Tseng, Gau, and Chou (2018)	Pediat.	Stroop	27 (89)	12.1	105.2	27 (78)	11.8	110.4	ADHD < TD: R MFG, R post-CG, R SPL
Hwang et al. (2015)	Pediat.	Stroop	26 (65)	13.9	106.4	35 (51)	14.5	105.1	ADHD < TD: Medial FG, cerebellar tonsil

Iannaccone, Hauser, Staempfli, et al. (2015)	Pediat.	GNG/ Flanker	18 (50)	14.5	108.5	18 (61)	14.8	114.4	ADHD<TD: R MFG/dIPFC/SFG; ADHD>TD: L/R MTG/STG, L insula/putamen/pre-CG, L ITG/FFG/PHG/MTG, L pons, L amygdala/hippocampus, L FFG/lingual gyrus, L post-CG/IPG, R brainstem/pons/cerebellum
Janssen, Heslenfeld, Mourik, Logan, and Oosterlaan (2015)	Pediat.	Stop	21 (90)	10.6	98.6	17 (76)	10.3	108.7	ADHD<TD: L/R mPFC, R MFG, R IFG, R pre-CG, R ACC, L SFG; ADHD>TD: L/R ACC, L/R MFG, R mPFC, R PCC, R cuneus, L IFG, L post-CG, L pre-CG.
Konrad, Neufang, Hanisch, Fink, and Herpertz-Dahlmann (2006)	Pediat.	Flanker	16 (100)	10.2	103	16 (100)	10.3	105	ADHD<TD: L pre-CG, R putamen; ADHD>TD: L medial SPL
Kooistra et al. (2010)	Adult	GNG	11 (100)	21.5	110	11 (100)	10.1	125	ADHD>TD: L/R ACC, R supramarginal gyrus, L angular gyrus, L/R supramarginal gyrus, R occipital gyrus, L frontal pole, L/R MFG, L/R IFG, L SFG, paracingulate gyrus, medial thalamus, L caudate nucleus, L posterior putamen, L lateral occipital gyrus, L/R pre-CG, R post-CG, R mPFC, L PCC, L STG
J. Ma et al. (2012)	Pediat.	GNG	15 (53)	9.8	100.2	15 (53)	22.3	102.6	ADHD>TD: R ITF, R midbrain, R pre-CG, R calcarine, R IOG, R MOG, L/R cerebellum, L post-CG, L IFG, R hippocampus
I. Ma et al. (2016)	Pediat.	Stroop	25 (76)	15.4	98.3	33 (67)	15.3	108.9	--
Massat et al. (2018)	Pediat.	Stop	18 (44)	10.0	106.8	19 (47)	10.6	114.4	ADHD>TD: L caudate body, L ventral putamen R dorsal putamen, R insula, L caudate tail, R middle CC, R ACC
Passarotti et al. (2010)	Pediat.	Stop	11 (55)	13.1	107.6	15 (48)	9.9	101.2	ADHD<TD: R MFG, R mPFC, L/R I/SFG, L STG; ADHD>TD: R caudate tail, L caudate, L anterior cerebellar vermis
Peterson et al. (2009)	Pediat.	Stroop	16 (81)	14.1	101.2	20 (60)	14.1	118.5	ADHD<TD: L ACC, insula R, precuneus, thalamus, caudate; ADHD>TD: R SFG, hippocampus, L ACC
Rasmussen et al. (2016)	Adult	GNG	50 (82)	24.8	102.1	23 (70)	24.1	109.2	ADHD<TD: R supramarginal gyrus, superior parietal, angular gyrus, L/R MFG. R frontal pole, L/R SPL, L/R IFG, L/R pre-/post-CG, R caudate/accumbens, L/R precuneus, R OFC, R posterior cingulate gyrus, R insula, R thalamus
Rubia, Smith, Brammer, Toone, and Taylor (2005)	Pediat.	Stop	16 (100)	13.2	100	21 (100)	13.4	95	ADHD<TD: R IFC, OFC, pre-CG, STL
Rubia, Halari, Cubillo, et al. (2011) ^d	Pediat.	Simon	12 (100)	13.0	90	13 (100)	14.0	102	ADHD<TD: R IFC/inferior parietal, L vmPFC/BG/thalamus, R SMA/ACC/PCC/SPL, L STL/MTL/occipital
Rubia, Halari, Mohammad, Taylor, and Brammer (2011) ^d	Pediat.	Stop	12 (100)	13.0	90	13 (100)	13.0	102	ADHD<TD: B IFC, insula, ACC, pre-SMA; thalamus; R MTL, occipital, IPL, precuneus, PCC, cerebellum
Schulz et al. (2004)	Pediat.	GNG	10 (100)	17.9	88.4	9 (100)	17.5	91.9	ADHD<TD: R pre-CG, R ITG, L hippocampus, R lingual gyrus, L/R cerebellum; ADHD>TD: L/R MFG, L/R IFG, L medial FG, L ACC, L/R IPL, R precuneus
Schulz et al. (2014)	Adult	GNG	14 (100)	23.3	n/a	14 (100)	22.8	n/a	--

Schulz et al. (2017)	Adult	Stroop	27 (89)	24.2	n/a	28 (86)	24.6	n/a	ADHD<TD: L/R cerebellum, L MTG, R lingual gyrus, R IFG, R MTG, R OFG, R SOG, R precuneus, R IPL, R MOG
Sebastian et al. (2012) ^e	Adult	GNG Stroop Stop	20 (55)	33.3	115.3	24 (46)	17.5	115.7	ADHD<TD: R caudate (GNG), R GP (Stop), R post-CG, L para-central lobule/mid-CC, L STG/MTG, R temporal pole, R GP (Stroop)
Shang, Sheng, Yang, Chou, and Gau (2018) ^f	Adult	Stroop	25 (48)	29.1	112.8	30 (50)	28.2	115.4	ADHD<TD: L IFG, L/R caudate nucleus
	Adult	Stroop	25 (56)	28.5	113.1	30 (50)	28.2	115.4	ADHD<TD: L IFG, L caudate nucleus, L SPL
Siniatchkin et al. (2012)	Pediat.	GNG	12 (83)	9.3	n/a	12 (75)	30.3	n/a	ADHD<TD: ACC, R dIPFC, R caudate nucleus
Smith, Taylor, Brammer, Toone, and Rubia (2006) ^g	Pediat.	GNG	17(100)	12.8	n/a	18 (100)	9.3	n/a	ADHD<TD: L MFG
		Stroop	19 (100)	12.9	n/a	24 (100)	12.8	n/a	--
		Switch	14 (100)	13.3	n/a	27 (100)	12.9	n/a	ADHD<TD: L STG/IFG/pre-CG/insula, R MTG/IFG, IPL/STG
Spinelli et al. (2011)	Pediat.	GNG	13 (69)	10.6	109.2	17 (47)	13.3	108.8	ADHD>TD: R pre-CG, MFG
Szekely et al. (2018)	Adult	Stop	64 (56)	24.0	n/a	84 (57)	24.5	n/a	--
Tamm, Menon, Ringel, and Reiss (2004)	Pediat.	GNG	10 (100)	16.0	109.2	12 (100)	10.6	111.6	ADHD<TD: R ACC/SMA/ MFG; ADHD>TD: L M/I/STG
Thornton, Bray, Langevin, and Dewey (2018)	Pediat.	GNG	20 (90)	12.4	109.4	20 (40)	10.6	112.6	--
van Hulst, de Zeeuw, Rijks, Neggers, and Durston (2017)	Pediat.	GNG	24 (100)	11.2	105.6	26 (100)	10.5	117.3	--
van Rooij et al. (2015)	Pediat.	Stop	108 (64)	15.1	92.7	77 (49)	16.0	109.2	ADHD<TD: L STG, pre-CG; ADHD>TD: L SMA
	Adult	Stop	77 (78)	20.3	99.1	45 (33)	14.6	106.4	ADHD<TD: L SFG, R hippocampal gyrus
Zamorano et al. (2017)	Pediat.	Stroop	17 (100)	11.6	104.2	17 (100)	11.7	109.8	ADHD>TD: R medial/IFG

Abbreviations. N=sample size, y=year, pediat=pediatric (child/adolescent) sample, ADHD=attentional-deficit/hyperactivity disorder, TD=typically developing controls, L/R=Left/Right, GNG=Go/No-Go. Brain regions (in alphabetical order): ACC=anterior cingulate cortex, AI=anterior insula, BG=basal ganglia, CC=cingulate cortex, dIPFC=dorsolateral prefrontal cortex, dACC=dorsal ACC, FFG=fusiform gyrus, GP=globus pallidus, I/M/SFG=inferior/middle/superior frontal gyrus, I/SPL=inferior/ superior parietal lobe, medial FG=medial frontal gyrus, mPFC=medial prefrontal cortex, M/STG=middle/superior temporal gyrus, OFC=orbitofrontal cortex, OFG=orbitofrontal gyrus, PCC=posterior cingulate cortex, PHG=parahippocampal gyrus, PMC=premotor cortex, pre-/post-CG=pre-/post-central gyrus, SMA=supplementary motor cortex, STL=superior temporal lobe, STS=superior temporal sulcus, vmPFC=ventromedial prefrontal cortex, a-g datasets were combined, adjusting their variance according to the method outlined in Norman et al. (2016). See Supplement Table S2a for further information.

Table 2. Sample Characteristics and Summary Findings of VBM and fMRI Cognitive Control Studies in ASD

Source	Age group	Task	Patients			Controls			Summary findings
			N (% male)	Age, y	IQ	N (% male)	Age, y	IQ	
(A) VBM studies in ASD									
Abell et al. (1999)	Adult	--	15 (80)	28.8	n/a	15 (80)	25.3	n/a	ASD<TD: R paracingulate, L IFG, L occipito-temporal junction; ASD>TD: L amygdala, L/R anterior cerebellum, L MTG, R ITG
Boddaert et al. (2004)	Pediat.	--	21 (76)	9.3	n/a	12 (58)	10.8	n/a	ASD<TD: L/R STS
Bonilha et al. (2008)	Pediat.	--	12 (100)	12.4	n/a	16 (100)	13.2	n/a	ASD>TD: L/R IFG, L/R cuneus, L/R ACC, R PCC, claustrum, L/R precuneus, L/R STG, L/R MTG, L ITG, thalamus (pulvinar), L/R SFG, L/R SPL, L insula, L putamen, R caudate, L/R FFG, L/R occipital gyrus, L lingual gyrus, L pre-CG, L post-CG, L/R thalamus, L/R PHG, L/R mPFC, R IPL
Brieber et al. (2007)	Pediat.	--	15 (100)	14.2	106.8	15 (100)	13.3	107.7	ASD<TD: L/R ITG/hippocampus/ amygdala, L MOG, L PMC, L hippocampus; ASD>TD: R supramarginal gyrus, L post-CG
Cai et al. (2018)	Pediat.	--	38 (84)	9.6	75.8	27 (96)	8.3	98.6	ASD<TD: R cerebellum anterior lobe, R precuneus, R angular gyrus; ASD>TD: L ITG, L/R MTG
Cheng, Chou, Fan, and Lin (2011)	Pediat.	--	25 (100)	13.7	101.6	25 (100)	13.5	109.0	ASD<TD: R IFG, R pre-CG, L post-CG, cuneus, thalamus, lingual gyrus, STG; ASD>TD: ACC, paracentral lobule, SPL, precuneus, MFG, FFG, subcallosal gyrus
Contarino, Bulgheroni, Annunziata, Erbetta, and Riva (2016)	Pediat.	--	25 (88)	6.1	56	25 (65)	6.1	103	--
M. Craig et al. (2007)	Adult	--	14 (0)	37.9	103.4	19 (0)	35.0	111.2	ASD<TD: Cuneus, L ITG/STG, R MTG, R ACC
D'Mello, Crocetti, Mostofsky, and Stoodley (2015)	Pediat.	--	35 (86)	10.4	n/a	35 (60)	10.4	n/a	ASD<TD: R lingual gyrus, R cerebellum, R angular gyrus; ASD>TD: L PCC/precuneus, R SFG, L MOG
Ecker et al. (2012)	Adult	--	89 (100)	27.0	110	89 (100)	28.0	113.0	ASD<TD: R occipital/ITG/MTG, R cerebellum, R posterior FFG, R lingual gyrus, L/R inferior and R SOG, R cuneus, precuneus, R PCC; ASD>TD: L /R ITG/MTG/STG/ L/R FFG/PHG/insula, L IFG, L putamen/caudate, L thalamus, L/R dIPFC/ MFG/pre-/post-CG, R IPL
Foster et al. (2015)	Pediat.	--	38 (100)	12.4	102.5	46 (100)	12.6	113.1	ASD<TD: R STG, L/R supramarginal gyrus, L cerebellum; ASD>TD: R central sulcus, L medial FG, L/R IFG, L/R pre-CG, L MFG, L pre-SMA, R SFG, L ACC, L OFC, L ITG/STG. L/R MTG, L Heschl's gyrus, R lingual gyrus, L FFG, L post-CG, L PCC, L precuneus, R supramarginal/angular gyrus, L IOG, L/R cuneus, L putamen, L caudate
Freitag et al. (2008)	Mixed	--	15 (87)	17.5	101.2	15 (87)	18.6	112.1	ASD<TD: R intraparietal sulcus
Greimel et al. (2013)	Mixed	--	47 (100)	18.3	107.5	51 (100)	21.4	112.5	ASD<TD: ACC, L/R posterior STS, R MTG

Groen, Buitelaar, van der Gaag, and Zwiers (2011)	Pediat.	--	17 (82)	14.4	98.0	25 (88)	15.5	105.0	--
Hyde, Samson, Evans, and Motttron (2010)	Mixed	--	15 (100)	22.7	100.4	15 (100)	19.2	106.6	ASD<TD: R post-CG, L/R pre-CG; ASD>TD: Brainstem/midbrain, reticular, medial FG/OFC, L/R MFG
Itahashi et al. (2015)	Adult	--	46 (100)	30.2	106.0	46 (100)	30.5	109.2	--
Katz et al. (2016)	Adult	--	23 (100)	26.6	n/a	32 (100)	29.8	n/a	ASD>TD: L/R OFC, ACC
Kaufmann et al. (2013)	Pediat.	--	10 (80)	14.7	102.3	10 (80)	13.8	109.5	ASD<TD: Lateral portion of R precuneus; ASD>TD: L medial FG, R precuneus
Ke et al. (2008)	Pediat.	--	17 (82)	8.9	108.8	15 (80)	9.7	109.8	ASD<TD: R PHG; ASD>TD: L/R supramarginal gyrus, post-CG, R MFG, R cerebellum
Kosaka et al. (2010)	Adult	--	32 (100)	23.8	101.6	40 (100)	22.5	109.7	ASD<TD: R insula, R IFG, R IPL
Kurth et al. (2011)	Pediat.	--	52 (73)	11.2	102.2	52 (73)	11.1	106.0	ASD<TD: Hypothalamus
Kwon, Ow, Pedatella, Lotspeich, and Reiss (2004)	Pediat.	--	20 (100)	13.5	n/a	13 (100)	13.6	n/a	ASD<TD: R ITG, R entorhinal cortex, R FFG
Langen et al. (2009)	Pediat.	--	99 (92)	12.9	107.6	89 (92)	12.4	110.0	--
Lim et al. (2015)	Pediat.	--	19 (100)	14.9	113.0	33 (100)	14.9	110.0	ASD>TD: L MTG/STG, L medial FG
Lin, Ni, Lai, Tseng, and Gau (2015)	Pediat.	--	28 (100)	10.7	106.9	43 (100)	10.6	115.2	ASD<TD: R post-CG, precuneus, L MOG; ASD>TD: L subcallosal gyrus, L/R sublobar areas
	Pediat.	--	40 (100)	14.7	101.5	18 (100)	15.5	108.7	--
	Adult	--	18 (100)	22.2	99.6	29 (100)	23.4	116.8	ASD>TD: L/R SFG, L MFG
Lin, Tseng, Lai, Chang, and Gau (2017)	Pediat.	--	20 (100)	13.5	103.8	54 (100)	12.8	112.5	ASD>TD: L cerebellum crus I
McAlonan et al. (2002)	Adult	--	17 (90)	32.0	96.0	24 (92)	33.0	114.0	ASD<TD: R cerebellum, L/R lenticular nucleus, R CG, L/R MFG, R SFG, precuneus
McAlonan et al. (2008)	Pediat.	--	33 (82)	11.6	113.2	55 (86)	10.7	117.1	ASD<TD: L dlPFC, L/R BG, L/R inferior cerebellar vermis, L STS, posterior parietal cortices
Mengotti et al. (2011)	Pediat.	--	20 (90)	7.0	n/a	22 (91)	7.7	n/a	ASD<TD: L SMA, R IFG; ASD>TD: R IPL, L/R ITG, L SPL, R SOG, precuneus
Mueller et al. (2013)	Adult	--	12 (75)	35.5	111.3	12 (67)	33.3	110.8	ASD<TD: L/R TPJ, L/R temporal lobe, L/R IFG, L MFG, L frontal pole, L/R mPFC, paracingulate gyrus
Ni et al. (2018)	Pediat.	--	81 (100)	12.6	107.4	61 (100)	12.4	112.0	ASD<TD: L lateral occipital/superior parietal cortex, L frontal pole
Pereira et al. (2018)	Mixed		19 (82)	17.4	99.8	25 (66)	18.5	105.8	ASD<TD: L/R cerebellum, posterior lobe, R cerebellum anterior lobe, L FFG, L/R CG, L paracentral lobule, R MFG, L claustrum, R medial FG, L PHG L amygdala, L post-CG, R SFG; ASD>TD: L/R brainstem, R cerebellum, posterior lobe
Poulin-Lord et al. (2014)	Mixed	--	23 (87)	19.8	100.3	22 (86)	22.6	107.3	--
Poustka et al. (2012)	Pediat.	--	18 (89)	9.7	111.0	18 (89)	9.7	112.8	--

Radeloff et al. (2014)	Mixed	--	34 (91)	19.1	105.7	26 (85)	19.5	107.8	--
Retico et al. (2016)	Pediat.	--	76 (50)	4.6	71	76 (50)	4.6	73	ASD<TD: L posterior MTG, L cerebellum, L/R temporal pole; ASD>TD: L precuneus, L PCC, R posterior STG, R MTG/ITG, R mPFC, L SPL, mid-CC
Riedel et al. (2014)	Adult	--	30 (63)	35.4	124.5	30 (63)	35.5	123.6	--
Riva et al. (2013)	Pediat.	--	26 (89)	5.8	51.6	21 (62)	6.8	n/a	ASD<TD: L/R cerebellum crus II, vermis 8/9, L/R hippocampus, L IFG, L MOG/SOG, L ITG, L SFG. R IOG, R post-CG
Rojas et al. (2006)	Mixed	--	24 (100)	20.8	94.8	23 (100)	21.4	118.7	--
Sato et al. (2017)	Adult	--	36 (69)	27.0	110.4	36 (69)	24.9	n/a	ASD<TD: R IOG/ITG/MTG, L FFG, L/R amygdala, R PHG, R L/MFG, R SMA, L/R CG, ACC, L/R medial SFG, R OFG
Schmitz et al. (2006)	Adult	--	10 (100)	38.0	105.0	12 (100)	39.0	106	ASD>TD: L IFG, ACC, R SFG, L/R MFG
Toal et al. (2010)	Adult	--	65 (88)	31.0	98.0	33 (91)	32.0	105	ASD<TD: R cerebellum/PHG/ FFG, R ITG/STG L PHG/STG/TG/FFG/ cerebellum
Waiter et al. (2004)	Pediat.	--	16 (100)	15.4	100.4	16 (100)	15.5	99.7	ASD<TD: R thalamus; ASD>TD: L SFG, R FFG, R mPFC, L MTG, R PCC, L/R STG, L lingual gyrus, L IFG, L MFG, L IOG, L PHG
Wang et al. (2017)	Pediat.	--	31 (100)	4.8	62.5	31 (100)	4.8	97.1	ASD>TD: L STG, L post-CG
Wilson, Tregellas, Hagerman, Rogers, and Rojas (2009)	Adult	--	10 (80)	30.1	91.5	10 (70)	29.4	127.2	--
Q. Yang et al. (2018)	Pediat.	--	16 (63)	10.4	45.3	16 (63)	10.5	97.5	ASD<TD: L cerebellum, L caudate
(B) fMRI studies in ASD									
Ambrosino et al. (2014)	Pediat.	GNG	19 (100)	11.5	112	19 (100)	11.1	120	--
Chantiluke, Barrett, Giampietro, Santosh, et al. (2015)	Pediat.	Stop	19 (100)	14.7	112	25 (100)	13.4	109	ASD<TD: L IPL; ASD>TD: R IFC, L IFG/MFG
Daly et al. (2014)	Adult	GNG	14 (100)	31	115	14 (100)	31	123	ASD<TD: R IFG, L thalamus; ASD>TD: R caudate, R cerebellum
Denisova et al. (2017)	Mixed	Simon	20 (95)	22.7	n/a	20 (95)	23.2	n/a	ASD>TD: R lingual gyrus, R cerebellum, R ITG
Duerden et al. (2013)	Adult	GNG	16 (69)	27.2	112	17 (71)	30.7	114	ASD<TD: R MFG; ASD>TD: R IFG, R FFG
J. Fan et al. (2012)	Adult	Flanker	12 (75)	30	115	12 (83)	28	120	ASD<TD: ACC
Gooskens et al. (2018)	Pediat.	Stop	26 (35)	11.3	108.9	53 (45)	10.8	111.9	--
Kana, Keller, Minshew, and Just (2007)	Adult	GNG	12 (92)	26.8	110	12 (92)	22.5	117	ASD<TD: L ITG, R PHG, R calcarine sulcus, R PMC, R mid-CC, L/R post-CG, R insula/IFG, L lingual gyrus
Kennedy et al. (2006)	Adult	Stroop	15 (100)	25.5	96	14 (100)	26.1	n/a	ASD>TD: mPFC, precuneus
Prat, Stocco, Neuhaus, and Kleinhans (2016)	Adult	GNG	16 (63)	25.3	107.4	17 (65)	25.6	111.0	--
Schmitz et al. (2006)	Adult	GNG, Switch, Stroop	10 (100)	38	105	12 (100)	39	106	ASD>TD: L M/IFG, OFC (GNG), insula (Stroop), and IPL, mesial parietal lobe (Switch)

Shafritz, Dichter, Baranek, and Belger (2008)	Adult	Switch	18 (89)	22.3	103	15 (87)	24.3	111	ASD<TD: L dIPFC, ACC, L intraparietal sulcus, L BG, L insula
Shafritz, Bregman, Ikuta, and Szeszko (2015)	Mixed	GNG	15 (80)	18.1	102	15 (80)	18.4	115	ASD<TD: R IFG/insula
Solomon et al. (2014)	Pediat.	Switch	27 (19)	15.4	108	27 (19)	16.1	113	ASD<TD: L PCC, L lingual gyrus, L MOG
Vaidya et al. (2011)	Pediat.	Stroop	11 (100)	10.8	114	14 (100)	11.0	119	ASD<TD: ACC, L MFG, R caudate
van Hulst et al. (2017)	Pediat.	GNG	26 (100)	10.8	109.6	26 (100)	10.5	117.3	--
Velasquez et al. (2017)	Adult	GNG	19 (68)	25.8	111	22 (73)	29.0	112	ASD<TD: R ACC, post-CG, R lateral occipital cortex
Yerys et al. (2015)	Pediat.	Switch	20 (80)	11.3	115	19 (68)	11.4	120	ASD>TD: L MFG/ pre-CG, L SFG/ dACC, R IFG

Abbreviations. N=sample size, y=year, pediat.=pediatric (child/adolescent) sample, ASD=autism spectrum disorder, TD=typically developing controls, L/R=Left/Right, GNG = Go/No-Go. Brain region (in alphabetical order): ACC=anterior cingulate cortex, BG=basal ganglia, dACC=dorsal ACC, dIPFC=dorsolateral prefrontal cortex, FFG=fusiform gyrus, I/M/medial/SFG=inferior/middle/medial/superior frontal gyrus, I/M/STG=inferior/middle/superior temporal gyrus, I/SPL=inferior/superior parietal lobe, mPFC=medial prefrontal cortex, OFC=orbital frontal cortex, OFG=orbital frontal gyrus, PCC=posterior cingulate cortex, pre-/post-CG=pre-/post-central gyrus, PHG=parahippocampal gyrus, PMC=premotor cortex, SMA=supplementary motor area, STS=superior temporal sulcus. See Supplement Table S2b for further information.

Table 3. Characteristics of Overall Sample and Subgroups Matched on Age, Sex and IQ

	ADHD	ASD	TD _{ADHD}	TD _{ASD}
(A) Overall Sample				
VBM				
N	1533	1445	1295	1477
% males	69	88	67	87
Mean age (range), y	20.9 (6-65)	17.2 (2-59)	21.1 (6-65)	16.4 (2-58)
Mean FSIQ (SD)	103 (7.2)	99 (17.3)	110 (10.8)	108 (5.1)
fMRI during Cognitive Control				
N	1001	335	1004	353
% males	77	83	69	80
Mean age (range), y	18.3 (7-50)	20.7 (7-52)	18.6 (7-50)	19.2 (7-52)
Mean FSIQ (SD)	102 (7.1)	109 (5.2)	109 (13.2)	114 (4.4)
fMRI during Motor Response Inhibition				
N	745	212	743	232
% males	77	82	69	78
Mean age (range), y	18.9 (7-50)	21.6 (8-52)	19.0 (7-50)	19.1 (8-52)
Mean FSIQ (SD)	101 (7.2)	109 (3.7)	109 (14.8)	114 (4.4)
(B) Matched Subgroups				
VBM				
N	825	758	778	781
% males	78	80	78	80
Mean age (range), y	16.7 (6-65)	17.4 (3-59)	17.4 (6-65)	16.1 (2-58)
Mean FSIQ (SD)	102 (8.3)	97 (18.0)	109 (4.7)	107 (12.6)
fMRI during Cognitive Control				
N	586	335	628	353
% males	83	83	79	80
Mean age (range), y	17.6 (7-50)	20.7 (7-52)	17.6 (7-50)	19.2 (7-52)
Mean FSIQ (SD)	105 (7.7)	109 (4.8)	109 (16.0)	114 (4.1)
fMRI during Motor Response Inhibition				
N	354	212	421	232
% males	78	82	73	78
Mean age (range), y	21.3 (7-50)	21.6 (8-52)	21.6 (7-50)	19.1 (8-52)
Mean FSIQ (SD)	106 (9.6)	109 (3.6)	114 (7.8)	114 (4.3)

Abbreviations: N = overall number of subjects, TD_{ADHD} = typically developing controls in the ADHD studies, TD_{ASD} = typically developing controls in the ASD studies, VBM = voxel-based morphometry, fMRI = functional Magnetic Resonance Imaging, % males = proportion of males among the samples, y = year, FSIQ = full scale IQ, and SD = standard deviation. The demographic characteristics presented here were calculated from the independent datasets with non-overlapping participants.

Table 4. Brain Structural and Functional Abnormalities in ADHD relative to ASD

Contrasts	MNI coord. x, y, z	SDM Z	p-value	Voxels No	BA
(A) Structural Abnormalities					
(i) ADHD (vs. TD)					
ADHD < TD					
L/R vmOFC/vmPFC/rACC and R caudate	2,48,-18	2.48	.00009	2902	11/10
R putamen/posterior insula/STG	30,-4,4	1.98	.0009	902	48/22
L pre-CG	-40,-6,56	1.88	.001	110	6
R rostralateral PFC	28,66,-4	1.98	.0009	79	11
L vIPFC/STG/temporal pole	-26,16,-24	1.93	.001	57	38
ADHD > TD					
Nil	--	--	--	--	--
(ii) ASD (vs. TD)					
ASD < TD					
L/R dACC/dmPFC	4,42,18	1.61	.0004	575	24/32
L cerebellum	-10,-66,-48	1.35	.002	113	
R hippocampus/PHG/FFG	24,-6,-22	1.38	.002	104	34/20/36
Dorsomedial thalamus	0,0,20	1.39	.001	46	
ASD > TD					
L anterior I/M/STG/posterior insula	-60,-20,-14	2.55	.000002	1828	20/21/22/48
L/R PCC/precuneus,	-8,-50,26	1.99	.0003	437	23/26/7
R MFG/SFG/dIPFC	24,48,24	2.13	.0001	218	46
ADHD (vs. TD) vs. ASD (vs. TD)					
(iii) Overall Sample					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
vmOFC/rACC/L caudate	4,22,-10	2.22	.00009	1543	25/11
L post-CG	-42,-14,48	1.86	.0007	276	6
L MFG/SFG/dIPFC	-14,58,14	1.90	.0006	114	10
ASD (vs. TD) increased vs. ADHD (vs. TD)					
L anterior STG/MTG	-46,-16,-2	1.91	.0005	601	48/22/20/21
R MFG/SFG/dIPFC	24,46,24	1.98	.0004	150	46
R MFG/SFG	18,66,6	1.73	.001	91	10
L MTG/temporal pole	-46,6,-26	1.68	.002	56	20
R posterior MTG/STG	46,-36,4	1.64	.002	52	21
(iv) Matched Subgroups					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
vmOFC/rACC	2,48,-18	2.03	.001	442	11/10
L MOG	-22,-88,10	2.02	.001	85	18
ASD (vs. TD) reduced vs. ADHD (vs. TD)					
R PHG/amygdala/FFG/hippocampus	28,-6,-24	1.37	.0002	386	28/34/36/35
Dorsomedial thalamus	2,-6,16	1.18	.0006	92	
ASD (vs. TD) increased vs. ADHD (vs. TD)					
R MTG/STG/angular gyrus	50,-40,8	2.28	.0004	463	42/40/22/21/39
R MFG/SFG/dIPFC	24,44,24	2.00	.001	65	46
L anterior MTG/STG	-50,-20,-8	1.84	.003	36	48/22/21/20
PCC	-6,-50,28	1.84	.003	33	23/30
(B) Functional Abnormalities during Cognitive Control					

(i) ADHD (vs. TD)						
ADHD < TD						
R thalamus/caudate	10,-4,12	1.74	.0001	449		
L M/STG/superior temporal pole	-52,-16,-12	1.31	.001	285	21/22/38	
L/R SMA/dmPFC	4,6,52	1.33	.001	130	6/24	
L IFG/AI/temporal pole	-50,14,4	1.25	.002	82	45/48	
R AI/putamen	32,20,0	1.27	.002	65	48	
L post-CG	-48,-12,54	1.26	.002	53	6	
L MFG/dIPFC	-30,30,44	1.24	.002	35	9	
R MTG	46,-2,-24	1.18	.003	33	21	
ADHD > TD						
Nil	--	--	--	--	--	--
(ii) ASD (vs. TD)						
ASD < TD						
L/R ACC/midcingulate/dmPFC	0,34,20	2.12	.000001	2708	24/10/32	
L MFG/dIPFC	-42,34,24	1.42	.0003	370	45/46/48	
R MFG/dIPFC	42,10,48	1.28	.0007	113	9	
L IPL	-34,-46,52	1.11	.002	91	40	
L lingual gyrus/cerebellum, lobule IV/V	-14,-48,-10	1.08	.003	89	30/19	
ASD > TD						
Precuneus/midcingulate	-2,-44,56	1.48	.00007	1115	7/5/4	
R IOG	36,-70,-6	1.37	.0002	226	19	
L vIPFC/OFC	-30,40,-8	1.14	.0009	90	47/11	
L MFG/rostrolateral PFC	-30,48,10	1.14	.0008	81	10	
R IFG	42,22,18	1.08	.002	64	45	
ADHD (vs. TD) vs. ASD (vs. TD)						
(iii) Overall Sample						
ASD (vs. TD) reduced vs. ADHD (vs. TD)						
rACC/midcingulate/dmPFC	0,30,20	2.42	~0	2903	24/10/32	
L MFG/dIPFC	-44,34,24	1.19	.0004	319	45/46/48	
ASD (vs. TD) increased vs. ADHD (vs. TD)						
Precuneus	0,-42,50	1.50	.00009	1001	7/4	
R IOG/FFG	36,-70,-8	1.35	.0003	147	19	
R IFG	44,24,18	1.27	.0005	138	48/45	
L vIPFC/OFC	-30,44,-8	1.04	.002	32	47/11	
L MFG	-30,50,8	1.02	.003	25	10	
(iv) Matched Subgroups						
ADHD (vs. TD) reduced vs. ASD (vs. TD)						
SMA	-4, 12, 54	-1.33	.0003	353	6	
L STG/MTG/temporal pole	-40, -6, -12	-1.01	.003	40	48/21/38	
ASD (vs. TD) reduced vs. ADHD (vs. TD)						
rACC/dmPFC	2,28,20	2.30	.0000002	2815	24/32/9/10	
L MFG/dIPFC	-40,34,28	1.03	.001	211	45/45/48	
L IPL/SPL	-32,-46,54	1.06	.0009	192	40/7/2	
ASD (vs. TD) increased vs. ADHD (vs. TD)						
Precuneus/PCC	-2, -40, 50	1.25	.0006	433	5/4	
R FFG/IOG/ITG	36, -70, -8	1.49	.0001	247	19/37	
R IFG	44, 24, 18	1.24	.0007	88	48/45	

L vIPFC/OFC	-30, 40, -8	1.10	.002	57	11/47
(C) Functional Abnormalities during Motor Response Inhibition					
(i) ADHD (vs. TD)					
ADHD < TD					
L MFG/dIPFC	-30,30,44	1.70	.0002	235	9/46/8
L anterior MTG/STG	-52,-18,-10	1.57	.0005	210	22/21/20/48
L post-CG	-48,-10,54	1.52	.0006	198	6
R IFG	46,32,22	1.54	.0005	188	45/48/46
R vIPFC/OFC/AI	30,36,-16	1.30	.002	172	47/38
R caudate	10,20,0	1.38	.001	91	--
ADHD > TD					
Nil	--	--	--	--	--
(ii) ASD (vs. TD)					
ASD < TD					
R AI/vIPFC	44,20,0	1.42	.0004	491	47/45/48
L cerebellum, hemispheric lobule IV & V/ lingual gyrus/FFG	-14,-46,-10	1.33	.001	492	30/19/18/37
R MFG/dIPFC	40,12,50	1.43	.0004	251	9
R PCC/precuneus	16,-36,44	1.35	.0008	58	7
ASD > TD					
L vIPFC/OFC	-28,48,8	1.19	.0003	703	11/10/46/47
R IOG/FFG/ITG	36,-72,-8	1.50	.00001	489	19/37
ADHD (vs. TD) vs. ASD (vs. TD)					
(iii) Overall Sample					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
R caudate	10,22,0	1.20	.0008	98	--
R IFG	42,26,20	1.14	.001	97	48/45
ASD (vs. TD) reduced vs. ADHD (vs. TD)					
L lingual gyrus/FFG/cerebellum lobule IV	-18,-54,-12	1.17	.001	243	19/37/30
rACC/dmPFC	0,32,22	1.09	.002	203	24/32
L MFG/IFG	-44,34,16	1.11	.002	109	45/48
R precuneus	16,-36,44	1.27	.0008	37	7
R MFG/dIPFC	42,10,48	1.04	.003	20	9/6
ASD (vs. TD) increased vs. ADHD (vs. TD)					
R IOG/FFG	36,-70,-8	1.84	.00001	417	19/37/18
L vIPFC/OFC	-28,42,-14	1.21	.0007	372	11/47
L MFG/SFG/dIPFC	-30,46,6	1.12	.001	183	10/46/11
L posterior MTG	-48,-54,4	1.16	.001	63	37/21
(iv) Matched Subgroups					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
R caudate	8,20,2	1.17	.001	117	--
R IFG	42,26,20	1.18	.001	74	48
ASD (vs. TD) reduced vs. ADHD (vs. TD)					
L cerebellum lobule IV/FFG/lingual gyrus	-16,-48,-12	1.39	.0003	1082	19/37
R MFG/dIPFC	42,14,48	1.22	.0007	104	9/6
L SPL/precuneus	-18,-70,58	1.00	.002	50	7
R precuneus	16,-36,44	1.21	.0008	41	7
ASD (vs. TD) increased vs. ADHD (vs. TD)					

R IOG/FFG	36,-70,-10	-1.82 .00001	504	19/37
L vIPFC/OFC	-28,36,-8	-1.17 .001	265	11/47
L MFG/SFG/dIPFC	-28,54,8	-1.10 .002	171	10/46/11

Abbreviations: MNI = Montreal Neurological Institute, SDM = Seed-based d mapping, BA = Brodmann Area, TD = typically developing controls, brain regions (in alphabetical order): AI = anterior insula, dACC = dorsal anterior cingulate cortex, dIPFC = dorsolateral prefrontal cortex, dmPFC = dorsomedial prefrontal cortex, FFG = fusiform gyrus, IFG = inferior frontal gyrus, I/MOG = inferior/middle occipital gyrus, I/M/STG = inferior/middle/superior temporal gyrus, I/SPL = inferior/superior parietal lobe, M/SFG = middle/superior frontal gyrus, PCC = posterior cingulate cortex, PFC=prefrontal cortex, PHG=parahippocampal gyrus, pre-CG=precentral gyrus, rACC = rostral anterior cingulate cortex, SMA = supplementary motor area, vIPFC = ventrolateral prefrontal cortex, vmOFC = ventromedial orbitofrontal cortex, vmPFC = ventromedial prefrontal cortex. Bold prints indicate regions which survived the sensitivity analyses in subgroups matched on age, sex and IQ.

REFERENCES

- Abell, F., Krams, M., Ashburner, J., Passingham, R., Friston, K., Frackowiak, R., . . . Frith, U. (1999). The neuroanatomy of autism: A voxel-based whole brain analysis of structural scans. *Neuroreport*, 10(8), 1647-1651. doi:10.1097/00001756-199906030-00005
- Ahrendts, J., Rüscher, N., Wilke, M., Philipsen, A., Eickhoff, S. B., Glauche, V., . . . Tebartz Van Elst, L. (2011). Visual cortex abnormalities in adults with ADHD: A structural MRI study. *The World Journal of Biological Psychiatry*, 12(4), 260-270. doi:10.3109/15622975.2010.518624
- Albaugh, M. D., Orr, C., Chaarani, B., Althoff, R. R., Allgaier, N., D'Alberto, N., . . . Potter, A. S. (2017). Inattention and reaction time variability are linked to ventromedial prefrontal volume in adolescents. *Biological Psychiatry*, 82(9), 660-668. doi:10.1016/j.biopsych.2017.01.003
- Ambrosino, S., Bos, D. J., van Raalten, T. R., Kobussen, N. A., van Belle, J., Oranje, B., & Durston, S. (2014). Functional connectivity during cognitive control in children with autism spectrum disorder: An independent component analysis. *Journal of Neural Transmission*, 121(9), 1145-1155. doi:10.1007/s00702-014-1237-8
- American Psychiatric Association (APA). (2013). *Diagnostic and Statistical Manual of Mental Disorders: DSM-5* (5th ed.). Washington, DC: APA.
- American Psychiatric Association (APA) (Ed.) (2000). *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR)*. Washington, DC: APA.
- Amico, F., Stauber, J., Koutsouleris, N., & Frodl, T. (2011). Anterior cingulate cortex gray matter abnormalities in adults with attention deficit hyperactivity disorder: A voxel-based morphometry study. *Psychiatry Research: Neuroimaging*, 191(1), 31-35. doi:10.1016/j.psychres.2010.08.011
- Aron, A. R., Herz, D. M., Brown, P., Forstmann, B. U., & Zaghoul, K. (2016). Frontosubthalamic circuits for control of action and cognition. *Journal of Neuroscience*, 36(45), 11489-11495. doi:10.1523/jneurosci.2348-16.2016
- Banich, M. T., Burgess, G. C., Depue, B. E., Ruzic, L., Bidwell, L. C., Hitt-Laustsen, S., . . . Willcutt, E. G. (2009). The neural basis of sustained and transient attentional control in young adults with ADHD. *Neuropsychologia*, 47(14), 3095-3104. doi:10.1016/j.neuropsychologia.2009.07.005
- Bari, A., & Robbins, T. W. (2013). Inhibition and impulsivity: Behavioral and neural basis of response control. *Progress in Neurobiology*, 108, 44-79. doi:10.1016/j.pneurobio.2013.06.005
- Bhaijiwala, M., Chevrier, A., & Schachar, R. (2014). Withholding and canceling a response in ADHD adolescents. *Brain and Behavior*, 4(5), 602-614. doi:10.1002/brb3.244
- Binder, J. R., Gross, W. L., Allendorfer, J. B., Bonilha, L., Chapin, J., Edwards, J. C., . . . Weaver, K. E. (2011). Mapping anterior temporal lobe language areas with fMRI: A multicenter normative study. *NeuroImage*, 54(2), 1465-1475. doi:10.1016/j.neuroimage.2010.09.048
- Boddaert, N., Chabane, N., Gervais, H., Good, C. D., Bourgeois, M., Plumet, M. H., . . . Zilbovicius, M. (2004). Superior temporal sulcus anatomical abnormalities in childhood autism: A voxel-based morphometry MRI study. *NeuroImage*, 23(1), 364-369. doi:10.1016/j.neuroimage.2004.06.016
- Bonath, B., Tegelbeckers, J., Wilke, M., Flechtner, H. H., & Krauel, K. (2016). Regional gray matter volume differences between adolescents with ADHD and typically developing controls: Further evidence for anterior cingulate involvement. *Journal of Attention Disorders*. doi:10.1177/1087054715619682
- Bonilha, L., Cendes, F., Rorden, C., Eckert, M., Dalgalarondo, P., Li, L. M., & Steiner, C. E. (2008). Gray and white matter imbalance - Typical structural abnormality underlying

- classic autism? *Brain and Development*, 30(6), 396-401.
doi:10.1016/j.braindev.2007.11.006
- Booth, J. R., Burman, D. D., Meyer, J. R., Lei, Z., Trommer, B. L., Davenport, N. D., . . . Mesulam, M. M. (2005). Larger deficits in brain networks for response inhibition than for visual selective attention in attention deficit hyperactivity disorder (ADHD). *Journal of Child Psychology and Psychiatry*, 46(1), 94-111. doi:10.1111/j.1469-7610.2004.00337.x
- Bralten, J., Greven, C. U., Franke, B., Mennes, M., Zwiers, M. P., Rommelse, N. N. J., . . . Buitelaar, J. K. (2016). Voxel-based morphometry analysis reveals frontal brain differences in participants with ADHD and their unaffected siblings. *Journal of Psychiatry & Neuroscience*, 41(4), 272-279. doi:10.1503/jpn.140377
- Breyer, J. L., Botzet, A. M., Winters, K. C., Stinchfield, R. D., August, G., & Realmuto, G. (2009). Young adult gambling behaviors and their relationship with the persistence of ADHD. *Journal of Gambling Studies*, 25(2), 227-238. doi:10.1007/s10899-009-9126-z
- Brieber, S., Neufang, S., Bruning, N., Kamp-Becker, I., Remschmidt, H., Herpertz-Dahlmann, B., . . . Konrad, K. (2007). Structural brain abnormalities in adolescents with autism spectrum disorder and patients with attention deficit/hyperactivity disorder. *Journal of Child Psychology and Psychiatry*, 48(12), 1251-1258. doi:10.1111/j.1469-7610.2007.01799.x
- Cai, J., Hu, X., Guo, K., Yang, P., Situ, M., & Huang, Y. (2018). Increased left inferior temporal gyrus was found in both low function autism and high function autism. *Frontiers in Psychiatry*, 9, 542. doi:10.3389/fpsy.2018.00542
- Carlisi, C. O., Norman, L. J., Lukito, S. S., Radua, J., Mataix-Cols, D., & Rubia, K. (2017). Comparative multimodal meta-analysis of structural and functional brain abnormalities in autism spectrum disorder and obsessive-compulsive disorder. *Biological Psychiatry*, 82(2), 83-102. doi:10.1016/j.biopsych.2016.10.006
- Carmona, S., Hoekzema, E., Antoni Ramos-Quiroga, J., Richarte, V., Canals, C., Bosch, R., . . . Vilarroya, O. (2012). Response inhibition and reward anticipation in medication-naïve adults with attention-deficit/hyperactivity disorder: A within-subject case-control neuroimaging study. *Human Brain Mapping*, 33(10), 2350-2361. doi:10.1002/hbm.21368
- Carmona, S., Vilarroya, O., Bielsa, A., Tremols, V., Soliva, J. C., Rovira, M., . . . Bulbena, A. (2005). Global and regional gray matter reductions in ADHD: A voxel-based morphometric study. *Neuroscience Letters*, 389(2), 88-93. doi:10.1016/j.neulet.2005.07.020
- Carper, R. A., & Courchesne, E. (2005). Localized enlargement of the frontal cortex in early autism. *Biological Psychiatry*, 57(2), 126-133. doi:10.1016/j.biopsych.2004.11.005
- Cauda, F., Geda, E., Sacco, K., D'Agata, F., Duca, S., Geminiani, G., & Keller, R. (2011). Grey matter abnormality in autism spectrum disorder: An activation likelihood estimation meta-analysis study. *Journal of Neurology Neurosurgery and Psychiatry*, 82(12), 1304-1313. doi:10.1136/jnnp.2010.239111
- Chantiluke, K., Barrett, N., Giampietro, V., Brammer, M., Simmons, A., Murphy, D. G., & Rubia, K. (2015). Inverse effect of fluoxetine on medial prefrontal cortex activation during reward reversal in ADHD and autism. *Cerebral Cortex*, 25(7), 1757-1770. doi:10.1093/cercor/bht365
- Chantiluke, K., Barrett, N., Giampietro, V., Brammer, M., Simmons, A., & Rubia, K. (2015). Disorder-dissociated effects of fluoxetine on brain function of working memory in attention deficit hyperactivity disorder and autism spectrum disorder. *Psychological Medicine*, 45(6), 1195-1205. doi:10.1017/s0033291714002232
- Chantiluke, K., Barrett, N., Giampietro, V., Santosh, P., Brammer, M., Simmons, A., . . . Rubia, K. (2015). Inverse fluoxetine effects on inhibitory brain activation in non-comorbid boys with ADHD and with ASD. *Psychopharmacology*, 232(12), 2071-2082. doi:10.1007/s00213-014-3837-2

- Chantiluke, K., Christakou, A., Murphy, C. M., Giampietro, V., Daly, E. M., Ecker, C., . . . Rubia, K. (2014). Disorder-specific functional abnormalities during temporal discounting in youth with attention deficit hyperactivity disorder (ADHD), autism and comorbid ADHD and autism. *Psychiatry Research*, 223(2), 113-120. doi:10.1016/j.psychresns.2014.04.006
- Charman, T., Pickles, A., Simonoff, E., Chandler, S., Loucas, T., & Baird, G. (2011). IQ in children with autism spectrum disorders: Data from the Special Needs and Autism Project (SNAP). *Psychological Medicine*, 41(03), 619-627. doi:10.1017/s0033291710000991
- Chen, C.-Y., Yen, J.-Y., Yen, C.-F., Chen, C.-S., Liu, G.-C., Liang, C.-Y., & Ko, C.-H. (2015). Aberrant brain activation of error processing among adults with attention deficit and hyperactivity disorder. *Kaohsiung Journal of Medical Sciences*, 31(4), 179-187. doi:10.1016/j.kjms.2015.01.001
- Cheng, Y., Chou, K. H., Fan, Y. T., & Lin, C. P. (2011). ANS: Aberrant neurodevelopment of the social cognition network in adolescents with autism spectrum disorders. *PLoS One*, 6(4). doi:10.1371/journal.pone.0018905
- Chou, T.-L., Chia, S., Shang, C.-Y., & Gau, S. S.-F. (2015). Differential therapeutic effects of 12-week treatment of atomoxetine and methylphenidate on drug-naïve children with attention deficit/hyperactivity disorder: A counting Stroop functional MRI study. *European Neuropsychopharmacology*, 25(12), 2300-2310. doi:10.1016/j.euroneuro.2015.08.024
- Christakou, A., Murphy, C. M., Chantiluke, K., Cubillo, A. I., Smith, A. B., Giampietro, V., . . . Rubia, K. (2013). Disorder-specific functional abnormalities during sustained attention in youth with attention deficit hyperactivity disorder (ADHD) and with autism. *Molecular Psychiatry*, 18(2), 236-244. doi:10.1038/mp.2011.185
- Congdon, E., Mumford, J. A., Cohen, J. R., Galvan, A., Aron, A. R., Xue, G., . . . Poldrack, R. A. (2010). Engagement of large-scale networks is related to individual differences in inhibitory control. *NeuroImage*, 53(2), 653-663. doi:10.1016/j.neuroimage.2010.06.062
- Contarino, V. E., Bulgheroni, S., Annunziata, S., Erbetta, A., & Riva, D. (2016). Widespread focal cortical alterations in autism spectrum disorder with intellectual disability detected by threshold-free cluster enhancement. *American Journal of Neuroradiology*, 37(9), 1721-1726. doi:10.3174/ajnr.A4779
- Courchesne, E., Campbell, K., & Solso, S. (2011). Brain growth across the life span in autism: Age-specific changes in anatomical pathology. *Brain Research*, 1380, 138-145. doi:10.1016/j.brainres.2010.09.101
- Courchesne, E., Mouton, P. R., Calhoun, M. E., Semendeferi, K., Ahrens-Barbeau, C., Hallet, M. J., . . . Pierce, K. (2011). Neuron number and size in prefrontal cortex of children with autism. *JAMA*, 306(18), 2001-2010. doi:10.1001/jama.2011.1638
- Craig, F., Margari, F., Legrottaglie, A. R., Palumbi, R., de Giambattista, C., & Margari, L. (2016). A review of executive function deficits in autism spectrum disorder and attention-deficit/hyperactivity disorder. *Neuropsychiatric Disease and Treatment*, 12, 1191-1202. doi:10.2147/ndt.s104620
- Craig, M., Zaman, S. H., Daly, E. M., Cutter, W. J., Robertson, D. M. W., Hallahan, B., . . . Murphy, D. G. M. (2007). Women with autistic-spectrum disorder: magnetic resonance imaging study of brain anatomy. *British Journal of Psychiatry*, 191, 224-228. doi:10.1192/bjp.bp.106.034603
- Cubillo, A., Halari, R., Ecker, C., Giampietro, V., Taylor, E., & Rubia, K. (2010). Reduced activation and inter-regional functional connectivity of fronto-striatal networks in adults with childhood attention-deficit hyperactivity disorder (ADHD) and persisting symptoms during tasks of motor inhibition and cognitive switching. *Journal of Psychiatric Research*, 44(10), 629-639. doi:10.1016/j.jpsychires.2009.11.016
- Cubillo, A., Halari, R., Giampietro, V., Taylor, E., & Rubia, K. (2011). Fronto-striatal underactivation during interference inhibition and attention allocation in grown up

- children with attention deficit/hyperactivity disorder and persistent symptoms. *Psychiatry Research*, 193(1), 17-27. doi:10.1016/j.psychresns.2010.12.014
- Cubillo, A., Halari, R., Smith, A., Taylor, E., & Rubia, K. (2012). A review of fronto-striatal and fronto-cortical brain abnormalities in children and adults with attention deficit hyperactivity disorder (ADHD) and new evidence for dysfunction in adults with ADHD during motivation and attention. *Cortex*, 48(2), 194-215. doi:10.1016/j.cortex.2011.04.007
- Cubillo, A., Smith, A. B., Barrett, N., Giampietro, V., Brammer, M. J., Simmons, A., & Rubia, K. (2014). Shared and drug-specific effects of atomoxetine and methylphenidate on inhibitory brain dysfunction in medication-naïve ADHD boys. *Cerebral Cortex*, 24(1), 174-185. doi:10.1093/cercor/bhs296
- D'Mello, A. M., Crocetti, D., Mostofsky, S. H., & Stoodley, C. J. (2015). Cerebellar gray matter and lobular volumes correlate with core autism symptoms. *NeuroImage: Clinical*, 7, 631-639. doi:10.1016/j.nicl.2015.02.007
- Daly, E. M., Ecker, C., Hallahan, B., Deeley, Q., Craig, M., Murphy, C., . . . Murphy, D. G. M. (2014). Response inhibition and serotonin in autism: A functional MRI study using acute tryptophan depletion. *Brain*, 137, 2600-2610. doi:10.1093/brain/awu178
- Denisova, K., Zhao, G., Wang, Z., Goh, S., Huo, Y., & Peterson, B. S. (2017). Cortical interactions during the resolution of information processing demands in autism spectrum disorders. *Brain and Behavior*, 7(2), e00596. doi:10.1002/brb3.596
- Dennis, M., Francis, D. J., Cirino, P. T., Schachar, R., Barnes, M. A., & Fletcher, J. M. (2009). Why IQ is not a covariate in cognitive studies of neurodevelopmental disorders. *Journal of the International Neuropsychological Society*, 15(03), 331. doi:10.1017/s1355617709090481
- Depue, B. E., Burgess, G. C., Bidwell, L. C., Willcutt, E. G., & Banich, M. T. (2010). Behavioral performance predicts grey matter reductions in the right inferior frontal gyrus in young adults with combined type ADHD. *Psychiatry Research: Neuroimaging*, 182(3), 231-237. doi:10.1016/j.psychresns.2010.01.012
- DeRamus, T. P., & Kana, R. K. (2015). Anatomical likelihood estimation meta-analysis of grey and white matter anomalies in autism spectrum disorders. *NeuroImage: Clinical*, 7, 525-536. doi:10.1016/j.nicl.2014.11.004
- Desmond, J. E., & Glover, G. H. (2002). Estimating sample size in functional MRI (fMRI) neuroimaging studies: statistical power analyses. *Journal of Neuroscience Methods*, 118(2), 115-128. doi: 10.1016/s0165-0270(02)00121-8
- Di Martino, A., Ross, K., Uddin, L. Q., Sklar, A. B., Castellanos, F. X., & Milham, M. P. (2009). Functional brain correlates of social and nonsocial processes in autism spectrum disorders: An activation likelihood estimation meta-analysis. *Biological Psychiatry*, 65(1), 63-74. doi:10.1016/j.biopsych.2008.09.022
- Di Martino, A., Zuo, X.-N., Kelly, C., Grzadzinski, R., Mennes, M., Schvarcz, A., . . . Milham, M. P. (2013). Shared and distinct intrinsic functional network centrality in autism and attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 74(8), 623-632. doi:10.1016/j.biopsych.2013.02.011
- Dibbets, P., Evers, E. A. T., Hurks, P. P. M., Bakker, K., & Jolles, J. (2010). Differential brain activation patterns in adult attention-deficit hyperactivity disorder (ADHD) associated with task switching. *Neuropsychology*, 24(4), 413-423. doi:10.1037/a0018997
- Dibbets, P., Evers, L., Hurks, P., Marchetta, N., & Jolles, J. (2009). Differences in feedback- and inhibition-related neural activity in adult ADHD. *Brain and Cognition*, 70(1), 73-83. doi:10.1016/j.bandc.2009.01.001
- Dichter, G. S. (2012). Functional magnetic resonance imaging of autism spectrum disorders. *Dialogues in Clinical Neuroscience*, 14(3), 319-351. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3513685/>
- Duerden, E. G., Mak-Fan, K. M., Taylor, M. J., & Roberts, S. W. (2012). Regional differences in grey and white matter in children and adults with autism spectrum disorders: An activation likelihood estimate (ALE) meta-analysis. *Autism Research*, 5(1), 49-66. doi:10.1002/aur.235

- Duerden, E. G., Taylor, M. J., Soorya, L. V., Wang, T., Fan, J., & Anagnostou, E. (2013). Neural correlates of inhibition of socially relevant stimuli in adults with autism spectrum disorder. *Brain Research*, 1533, 80-90. doi:10.1016/j.brainres.2013.08.021
- Durston, S., Mulder, M., Casey, B. J., Ziermans, T., & van Engeland, H. (2006). Activation in ventral prefrontal cortex is sensitive to genetic vulnerability for attention-deficit hyperactivity disorder. *Biological Psychiatry*, 60(10), 1062-1070. doi:10.1016/j.biopsych.2005.12.020
- Ecker, C. (2017). The neuroanatomy of autism spectrum disorder: An overview of structural neuroimaging findings and their translatability to the clinical setting. *Autism*, 21(1), 18-28. doi:10.1177/1362361315627136
- Ecker, C., Schmeisser, M. J., Loth, E., & Murphy, D. G. (2017). Neuroanatomy and neuropathology of autism spectrum disorder in humans. *Advances in Anatomy, Embryology and Cell Biology*, 224, 27-48. doi:10.1007/978-3-319-52498-6_2
- Ecker, C., Suckling, J., Deoni, S. C., Lombardo, M. V., Bullmore, E. T., Baron-Cohen, S., . . . Murphy, D. G. M. (2012). Brain anatomy and its relationship to behavior in adults with autism spectrum disorder. *Archives of General Psychiatry*, 69(2), 195-209. doi:10.1001/archgenpsychiatry.2011.1251
- Elsabbagh, M., Divan, G., Koh, Y.-J., Kim, Y. S., Kauchali, S., Marcín, C., . . . Fombonne, E. (2012). Global prevalence of autism and other pervasive developmental disorders: global epidemiology of autism. *Autism Research*, 5(3), 160-179. doi:10.1002/aur.239
- Fan, J., Bernardi, S., Van Dam, N. T., Anagnostou, E., Gu, X., Martin, L., . . . Hof, P. R. (2012). Functional deficits of the attentional networks in autism. *Brain and Behavior*, 2(5), 647-660. doi:10.1002/brb3.90
- Fan, L. Y., Chou, T. L., & Gau, S. S. (2017). Neural correlates of atomoxetine improving inhibitory control and visual processing in drug-naïve adults with attention-deficit/hyperactivity disorder. *Human Brain Mapping*, 38(10), 4850-4864. doi:10.1002/hbm.23683
- Fan, L. Y., Shang, C. Y., Tseng, W. I., Gau, S. S., & Chou, T. L. (2018). Visual processing as a potential endophenotype in youths with attention-deficit/hyperactivity disorder: A sibling study design using the counting Stroop functional MRI. *Human Brain Mapping*, 39(10), 3827-3835. doi:10.1002/hbm.24214
- Floris, D. L., & Howells, H. (2018). Atypical structural and functional motor networks in autism. *Progress in Brain Research*, 238, 207-248. doi:10.1016/bs.pbr.2018.06.010
- Foster, N. E. V., Doyle-Thomas, K. A. R., Tryfon, A., Ouimet, T., Anagnostou, E., Evans, A. C., . . . Hyde, K. L. (2015). Structural gray matter differences during childhood development in autism spectrum disorder: A multimetric approach. *Pediatric Neurology*, 53(4), 350-359. doi:10.1016/j.pediatrneurol.2015.06.013
- Freitag, C. M., Konrad, C., Häberlen, M., Kleser, C., von Gontard, A., Reith, W., . . . Krick, C. (2008). Perception of biological motion in autism spectrum disorders. *Neuropsychologia*, 46(5), 1480-1494. doi:10.1016/j.neuropsychologia.2007.12.025
- Friston, K. J., Rotshtein, P., Geng, J. J., Sterzer, P., & Henson, R. N. (2006). A critique of functional localisers. *NeuroImage*, 30(4), 1077-1087. doi:10.1016/j.neuroimage.2005.08.012
- Frodl, T., & Skokauskas, N. (2012). Meta-analysis of structural MRI studies in children and adults with attention deficit hyperactivity disorder indicates treatment effects. *Acta Psychiatrica Scandinavica*, 125(2), 114-126. doi:10.1111/j.1600-0447.2011.01786.x
- Fuentes, P., Barros-Loscertales, A., Bustamante, J. C., Rosell, P., Costumero, V., & Avila, C. (2012). Individual differences in the behavioral inhibition system are associated with orbitofrontal cortex and precuneus gray matter volume. *Cognitive, Affective, & Behavioral Neuroscience*, 12(3), 491-498. doi:10.3758/s13415-012-0099-5
- Gehricke, J. G., Kruggel, F., Thampipop, T., Alejo, S. D., Tatos, E., Fallon, J., & Muftuler, L. T. (2017). The brain anatomy of attention-deficit/hyperactivity disorder in young adults - A magnetic resonance imaging study. *PLoS One*, 12(4). doi:10.1371/journal.pone.0175433

- Geurts, H. M., van den Bergh, S. F. W. M., & Ruzzano, L. (2014). Prepotent response inhibition and interference control in autism spectrum disorders: Two meta-analyses. *Autism Research*, 7(4), 407-420. doi:10.1002/aur.1369
- Gjevik, E., Eldevik, S., Fjæran-Granum, T., & Sponheim, E. (2011). Kiddie-SADS reveals high rates of DSM-IV disorders in children and adolescents with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 41(6), 761-769. doi:10.1007/s10803-010-1095-7
- Gooskens, B., Bos, D. J., Mensen, V. T., Shook, D. A., Bruchhage, M. M. K., Naaijen, J., . . . consortium, T. (2018). No evidence of differences in cognitive control in children with autism spectrum disorder or obsessive-compulsive disorder: An fMRI study. *Developmental Cognitive Neuroscience*, 100602. doi:10.1016/j.dcn.2018.11.004
- Greimel, E., Nehrkorn, B., Schulte-Ruether, M., Fink, G. R., Nickl-Jockschat, T., Herpertz-Dahlmann, B., . . . Eickhoff, S. B. (2013). Changes in grey matter development in autism spectrum disorder. *Brain Structure and Function*, 218(4), 929-942. doi:10.1007/s00429-012-0439-9
- Griebing, J., Minshew, N. J., Bodner, K., Libove, R., Bansal, R., Konasale, P., . . . Hardan, A. (2010). Dorsolateral prefrontal cortex magnetic resonance imaging measurements and cognitive performance in autism. *Journal of Child Neurology*, 25(7), 856-863. doi:10.1177/0883073809351313
- Groen, W. B., Buitelaar, J. K., van der Gaag, R. J., & Zwiers, M. P. (2011). Pervasive microstructural abnormalities in autism: A DTI study. *Journal of Psychiatry & Neuroscience*, 36(1), 32-40. doi:10.1503/jpn.090100
- Hart, H., Radua, J., Nakao, T., Mataix-Cols, D., & Rubia, K. (2013). Meta-analysis of functional magnetic resonance imaging studies of inhibition and attention in attention-deficit/hyperactivity disorder: exploring task-specific, stimulant medication, and age effects. *JAMA Psychiatry*, 70(2), 185-198. doi:10.1001/jamapsychiatry.2013.277
- Hazlett, H. C., Gu, H., Munsell, B. C., Kim, S. H., Styner, M., Wolff, J. J., . . . Statistical, A. (2017). Early brain development in infants at high risk for autism spectrum disorder. *Nature*, 542(7641), 348-351. doi:10.1038/nature21369
- He, N., Li, F., Li, Y., Guo, L., Chen, L., Huang, X., . . . Gong, Q. (2015). Neuroanatomical deficits correlate with executive dysfunction in boys with attention deficit hyperactivity disorder. *Neuroscience Letters*, 600, 45-49. doi:10.1016/j.neulet.2015.05.062
- Hollingdale, J., Woodhouse, E., Young, S., Fridman, A., & Mandy, W. (2019). Autistic spectrum disorder symptoms in children and adolescents with attention-deficit/hyperactivity disorder: A meta-analytical review. *Psychological Medicine*, 1-14. doi:10.1017/S0033291719002368
- Hoogman, M., Bralten, J., Hibar, D. P., Mennes, M., Zwiers, M. P., Schweren, L. S., . . . Franke, B. (2017). Subcortical brain volume differences in participants with attention deficit hyperactivity disorder in children and adults: A cross-sectional mega-analysis. *Lancet Psychiatry*, 4(4), 310-319. doi:10.1016/S2215-0366(17)30049-4
- Hoogman, M., Muetzel, R., Guimaraes, J. P., Shumskaya, E., Mennes, M., Zwiers, M. P., . . . Franke, B. (2019). Brain Imaging of the Cortex in ADHD: A coordinated analysis of large-scale clinical and population-based samples. *American Journal of Psychiatry*, 176(7), 531-542. doi:10.1176/appi.ajp.2019.18091033
- Hwang, S., White, S. F., Nolan, Z. T., Craig Williams, W., Sinclair, S., & Blair, R. J. R. (2015). Executive attention control and emotional responding in attention-deficit/hyperactivity disorder - A functional MRI study. *NeuroImage: Clinical*, 9, 545-554. doi:10.1016/j.nicl.2015.10.005
- Hyde, K. L., Samson, F., Evans, A. C., & Mottron, L. (2010). Neuroanatomical differences in brain areas implicated in perceptual and other core features of autism revealed by cortical thickness analysis and voxel-based morphometry. *Human Brain Mapping*, 31(4), 556-566. doi:10.1002/hbm.20887
- Iannaccone, R., Hauser, T. U., Ball, J., Brandeis, D., Walitza, S., & Brem, S. (2015). Classifying adolescent attention-deficit/hyperactivity disorder (ADHD) based on

- functional and structural imaging. *European Child & Adolescent Psychiatry*, 24(10), 1279-1289. doi:10.1007/s00787-015-0678-4
- Iannaccone, R., Hauser, T. U., Staempfli, P., Walitza, S., Brandeis, D., & Brem, S. (2015). Conflict monitoring and error processing: New insights from simultaneous EEG–fMRI. *NeuroImage*, 105, 395-407. doi:10.1016/j.neuroimage.2014.10.028
- Itahashi, T., Yamada, T., Nakamura, M., Watanabe, H., Yamagata, B., Jimbo, D., . . . Hashimoto, R. (2015). Linked alterations in gray and white matter morphology in adults with high-functioning autism spectrum disorder: A multimodal brain imaging study. *NeuroImage. Clinical*, 7, 155-169. doi:10.1016/j.nicl.2014.11.019
- Jagger-Rickels, A. C., Kibby, M. Y., & Constance, J. M. (2018). Global gray matter morphometry differences between children with reading disability, ADHD, and comorbid reading disability/ADHD. *Brain and Language*, 185, 54-66. doi:10.1016/j.bandl.2018.08.004
- Janssen, T. W. P., Heslenfeld, D. J., Mourik, R. V., Logan, G. D., & Oosterlaan, J. (2015). Neural correlates of response inhibition in children with attention-deficit/hyperactivity disorder: A controlled version of the stop-signal task. *Psychiatry Research: Neuroimaging*, 233(2), 278-284. doi:10.1016/j.psychresns.2015.07.007
- Johnston, B. A., Mwangi, B., Matthews, K., Coghill, D., Konrad, K., & Steele, J. D. (2014). Brainstem abnormalities in attention deficit hyperactivity disorder support high accuracy individual diagnostic classification. *Human Brain Mapping*, 35(10), 5179-5189. doi:10.1002/hbm.22542
- Kana, R. K., Keller, T. A., Minshew, N. J., & Just, M. A. (2007). Inhibitory control in high-functioning autism: decreased activation and underconnectivity in inhibition networks. *Biological Psychiatry*, 62(3), 198-206. doi:10.1016/j.biopsych.2006.08.004
- Kana, R. K., Maximo, J. O., Williams, D. L., Keller, T. A., Schipul, S. E., Cherkassky, V. L., . . . Just, M. A. (2015). Aberrant functioning of the theory-of-mind network in children and adolescents with autism. *Molecular Autism*, 6, 59. doi:10.1186/s13229-015-0052-x
- Kana, R. K., Patriquin, M. A., Black, B. S., Channell, M. M., & Wicker, B. (2016). Altered medial frontal and superior temporal response to implicit processing of emotions in autism. *Autism Research*, 9(1), 55-66. doi:10.1002/aur.1496
- Kappel, V., Lorenz, R. C., Streifling, M., Renneberg, B., Lehmkuhl, U., Stroehle, A., . . . Beck, A. (2015). Effect of brain structure and function on reward anticipation in children and adults with attention deficit hyperactivity disorder combined subtype. *Social Cognitive and Affective Neuroscience*, 10(7), 945-951. doi:10.1093/scan/nsu135
- Katz, J., d'Albis, M. A., Boisgontier, J., Poupon, C., Mangin, J. F., Guevara, P., . . . Houenou, J. (2016). Similar white matter but opposite grey matter changes in schizophrenia and high-functioning autism. *Acta Psychiatrica Scandinavica*, 134(1), 31-39. doi:10.1111/acps.12579
- Kaufmann, L., Zotter, S., Pixner, S., Starke, M., Haberlandt, E., Steinmayr-Gensluckner, M., . . . Marksteiner, J. (2013). Brief report: CANTAB performance and brain structure in pediatric patients with Asperger syndrome. *Journal of Autism and Developmental Disorders*, 43(6), 1483-1490. doi:10.1007/s10803-012-1686-6
- Kaya, B. S., Metin, B., Tas, Z. C., Buyukaslan, A., Soysal, A., Hatiloglu, D., & Tarhan, N. (2018). Gray matter increase in motor cortex in pediatric adhd: A voxel-based morphometry study. *Journal of Attention Disorders*, 22(7), 611-618. doi:10.1177/1087054716659139
- Ke, X., Hong, S., Tang, T., Zou, B., Li, H., Hang, Y., . . . Liu, Y. (2008). Voxel-based morphometry study on brain structure in children with high-functioning autism. *Neuroreport*, 19(9), 921-925. doi:10.1097/wnr.0b013e328300edf3
- Kennedy, D. P., Redcay, E., & Courchesne, E. (2006). Failing to deactivate: Resting functional abnormalities in autism. *Proceedings of the National Academy of Sciences of the United States of America*, 103(21), 8275-8280. doi:10.1073/pnas.0600674103

- Kleinhans, N. M., Muller, R. A., Cohen, D. N., & Courchesne, E. (2008). Atypical functional lateralization of language in autism spectrum disorders. *Brain Research*, 1221, 115-125. doi:10.1016/j.brainres.2008.04.080
- Kobel, M., Bechtel, N., Specht, K., Klarhoefer, M., Weber, P., Scheffler, K., . . . Penner, I.-K. (2010). Structural and functional imaging approaches in attention deficit/hyperactivity disorder: Does the temporal lobe play a key role? *Psychiatry Research: Neuroimaging*, 183(3), 230-236. doi:10.1016/j.psychresns.2010.03.010
- Kohls, G., Thönessen, H., Bartley, G. K., Grossheinrich, N., Fink, G. R., Herpertz-Dahlmann, B., & Konrad, K. (2014). Differentiating neural reward responsiveness in autism versus ADHD. *Developmental Cognitive Neuroscience*, 10, 104-116. doi:10.1016/j.dcn.2014.08.003
- Konrad, K., Neufang, S., Hanisch, C., Fink, G. R., & Herpertz-Dahlmann, B. (2006). Dysfunctional attentional networks in children with attention deficit/hyperactivity disorder: Evidence from an event-related functional magnetic resonance imaging study. *Biological Psychiatry*, 59(7), 643-651. doi:10.1016/j.biopsych.2005.08.013
- Kooistra, L., van der Meere, J. J., Edwards, J. D., Kaplan, B. J., Crawford, S., & Goodyear, B. G. (2010). Preliminary fMRI findings on the effects of event rate in adults with ADHD. *Journal of Neural Transmission*, 117(5), 655-662. doi:10.1007/s00702-010-0374-y
- Korponay, C., Dentico, D., Kral, T., Ly, M., Kruis, A., Goldman, R., . . . Davidson, R. J. (2017). Neurobiological correlates of impulsivity in healthy adults: Lower prefrontal gray matter volume and spontaneous eye-blink rate but greater resting-state functional connectivity in basal ganglia-thalamo-cortical circuitry. *NeuroImage*, 157, 288-296. doi:10.1016/j.neuroimage.2017.06.015
- Kosaka, H., Omori, M., Munesue, T., Ishitobi, M., Matsumura, Y., Takahashi, T., . . . Wada, Y. (2010). Smaller insula and inferior frontal volumes in young adults with pervasive developmental disorders. *NeuroImage*, 50(4), 1357-1363. doi:10.1016/j.neuroimage.2010.01.085
- Koshino, H., Carpenter, P. A., Minshew, N. J., Cherkassky, V. L., Keller, T. A., & Just, M. A. (2005). Functional connectivity in an fMRI working memory task in high-functioning autism. *NeuroImage*, 24(3), 810-821. doi:10.1016/j.neuroimage.2004.09.028
- Krall, S. C., Rottschy, C., Oberwelland, E., Bzdok, D., Fox, P. T., Eickhoff, S. B., . . . Konrad, K. (2015). The role of the right temporoparietal junction in attention and social interaction as revealed by ALE meta-analysis. *Brain Structure and Function*, 220(2), 587-604. doi:10.1007/s00429-014-0803-z
- Kuiper, M. W. M., Verhoeven, E. W. M., & Geurts, H. M. (2016). The role of interstimulus interval and "stimulus-type" in prepotent response inhibition abilities in people with ASD: A quantitative and qualitative review. *Autism Research*, 9(11), 1124-1141. doi:10.1002/aur.1631
- Kumar, U., Arya, A., & Agarwal, V. (2017). Neural alterations in ADHD children as indicated by voxel-based cortical thickness and morphometry analysis. *Brain and Development*, 39(5), 403-410. doi:10.1016/j.braindev.2016.12.002
- Kurth, F., Narr, K. L., Woods, R. P., O'Neill, J., Alger, J. R., Caplan, R., . . . Levitt, J. G. (2011). Diminished gray matter within the hypothalamus in autism disorder: A potential link to hormonal effects? *Biological Psychiatry*, 70(3), 278-282. doi:10.1016/j.biopsych.2011.03.026
- Kwon, H., Ow, A. W., Pedatella, K. E., Lotspeich, L. J., & Reiss, A. L. (2004). Voxel-based morphometry elucidates structural neuroanatomy of high-functioning autism and Asperger syndrome. *Developmental Medicine & Child Neurology*, 46(11), 760-764. doi:10.1017/s0012162204001306
- Lange, N., Travers, B. G., Bigler, E. D., Prigge, M. B. D., Froehlich, A. L., Nielsen, J. A., . . . Lainhart, J. E. (2015). Longitudinal volumetric brain changes in autism spectrum disorder ages 6-35 years. *Autism Research*, 8(1), 82-93. doi:10.1002/aur.1427

- Langen, M., Schnack, H. G., Nederveen, H., Bos, D., Lahuis, B. E., de Jonge, M. V., . . . Durston, S. (2009). Changes in the developmental trajectories of striatum in autism. *Biological Psychiatry*, 66(4), 327-333. doi:10.1016/j.biopsych.2009.03.017
- Li, X., Cao, Q., Pu, F., Li, D., Fan, Y., An, L., . . . Wang, Y. (2015). Abnormalities of structural covariance networks in drug-naïve boys with attention deficit hyperactivity disorder. *Psychiatry Research*, 231(3), 273-278. doi:10.1016/j.psychres.2015.01.006
- Liddle, E. B., Hollis, C., Batty, M. J., Groom, M. J., Totman, J. J., Liotti, M., . . . Liddle, P. F. (2011). Task-related default mode network modulation and inhibitory control in ADHD: effects of motivation and methylphenidate. *Journal of Child Psychology and Psychiatry*, 52(7), 761-771. doi:10.1111/j.1469-7610.2010.02333.x
- Lim, L., Chantiluke, K., Cubillo, A. I., Smith, A. B., Simmons, A., Mehta, M. A., & Rubia, K. (2015). Disorder-specific grey matter deficits in attention deficit hyperactivity disorder relative to autism spectrum disorder. *Psychological Medicine*, 45(5), 965-976. doi:10.1017/S0033291714001974
- Lin, H. Y., Ni, H.-C., Lai, M.-C., Tseng, W.-Y. I., & Gau, S. S.-F. (2015). Regional brain volume differences between males with and without autism spectrum disorder are highly age-dependent. *Molecular Autism*, 6, 29. doi:10.1186/s13229-015-0022-3
- Lin, H. Y., Tseng, W. I., Lai, M. C., Chang, Y. T., & Gau, S. S. (2017). Shared atypical brain anatomy and intrinsic functional architecture in male youth with autism spectrum disorder and their unaffected brothers. *Psychological Medicine*, 47(4), 639-654. doi:10.1017/S0033291716002695
- Lipszyc, J., & Schachar, R. (2010). Inhibitory control and psychopathology: A meta-analysis of studies using the stop signal task. *Journal of the International Neuropsychological Society*, 16(06), 1064-1076. doi:10.1017/S1355617710000895
- Lombardo, M. V., Chakrabarti, B., Bullmore, E. T., & Baron-Cohen, S. (2011). Specialization of right temporo-parietal junction for mentalizing and its relation to social impairments in autism. *NeuroImage*, 56(3), 1832-1838. doi:10.1016/j.neuroimage.2011.02.067
- Ma, I., van Holstein, M., Mies, G. W., Mennes, M., Buitelaar, J., Cools, R., . . . Scheres, A. (2016). Ventral striatal hyperconnectivity during rewarded interference control in adolescents with ADHD. *Cortex*, 82, 225-236. doi:10.1016/j.cortex.2016.05.021
- Ma, J., Lei, D., Jin, X., Du, X., Jiang, F., Li, F., . . . Shen, X. (2012). Compensatory brain activation in children with attention deficit/hyperactivity disorder during a simplified go/no-go task. *Journal of Neural Transmission*, 119(5), 613-619. doi:10.1007/s00702-011-0744-0
- Maier, S., Perlov, E., Graf, E., Dieter, E., Sobanski, E., Rump, M., . . . Elst, L. T. v. (2015). Discrete global but no focal gray matter volume reductions in unmedicated adult patients with attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 80(12), 905-915. doi:10.1016/j.biopsych.2015.05.012
- Martinez-Sanchis, S. (2014). Neurobiological foundations of multisensory integration in people with autism spectrum disorders: The role of the medial prefrontal cortex. *Frontiers in Human Neuroscience*, 8, 970. doi:10.3389/fnhum.2014.00970
- Massat, I., Slama, H., Villemonteix, T., Mary, A., Baijot, S., Albajara Saenz, A., . . . Peigneux, P. (2018). Hyperactivity in motor response inhibition networks in unmedicated children with attention deficit-hyperactivity disorder. *The World Journal of Biological Psychiatry*, 19(2), 101-111. doi:10.1080/15622975.2016.1237040
- McAlonan, G. M., Cheung, V., Cheung, C., Chua, S. E., Murphy, D. G. M., Suckling, J., . . . Ho, T. P. (2007). Mapping brain structure in attention deficit-hyperactivity disorder: A voxel-based MRI study of regional grey and white matter volume. *Psychiatry Research: Neuroimaging*, 154(2), 171-180. doi:10.1016/j.psychres.2006.09.006
- McAlonan, G. M., Daly, E., Kumari, V., Critchley, H. D., van Amelsvoort, T., Suckling, J., . . . Murphy, D. G. M. (2002). Brain anatomy and sensorimotor gating in Asperger's syndrome. *Brain*, 125, 1594-1606. doi:10.1093/brain/awf150
- McAlonan, G. M., Suckling, J., Wong, N., Cheung, V., Lienenkaemper, N., Cheung, C., & Chua, S. E. (2008). Distinct patterns of grey matter abnormality in high-functioning

- autism and Asperger's syndrome. *Journal of Child Psychology and Psychiatry*, 49(12), 1287-1295. doi:10.1111/j.1469-7610.2008.01933.x
- McCarthy, H., Skokauskas, N., & Frodl, T. (2014). Identifying a consistent pattern of neural function in attention deficit hyperactivity disorder: A meta-analysis. *Psychological Medicine*, 44(4), 869-880. doi:10.1017/s0033291713001037
- McGrath, L. M., & Stoodley, C. J. (2019). Are there shared neural correlates between dyslexia and ADHD? A meta-analysis of voxel-based morphometry studies. *Journal of Neurodevelopmental Disorders*, 11(1), 31. doi:10.1186/s11689-019-9287-8
- Mengotti, P., D'Agostini, S., Terlevic, R., De Colle, C., Biasizzo, E., Londero, D., . . . Brambilla, P. (2011). Altered white matter integrity and development in children with autism: A combined voxel-based morphometry and diffusion imaging study. *Brain Research Bulletin*, 84(2), 189-195. doi:10.1016/j.brainresbull.2010.12.002
- Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: A network model of insula function. *Brain Structure and Function*, 214(5-6), 655-667. doi:10.1007/s00429-010-0262-0
- Miller, G. M., & Chapman, J. P. (2001). Misunderstanding analysis of covariance. *Journal of Abnormal Psychology*, 110(1), 40-48. doi:10.1037//0021-843x.110.1.40
- Mitchell, S. R., Reiss, A. L., Tatusko, D. H., Ikuta, I., Kazmerski, D. B., Botti, J. A., . . . Kates, W. R. (2009). Neuroanatomic alterations and social and communication deficits in monozygotic twins discordant for autism disorder. *American Journal of Psychiatry*, 166(8), 917-925. doi:10.1176/appi.ajp.2009.08101538
- Montes, L. G. A., Ricardo-Garcell, J., de la Torre, L. B., Alcántara, H. P., García, R. B. M., Fernández-Bouzas, A., & Acosta, D. A. (2010). Clinical correlations of grey matter reductions in the caudate nucleus of adults with attention deficit hyperactivity disorder. *Journal of Psychiatry and Neuroscience*, 35(4), 238-246. doi:10.1503/jpn.090099
- Moreno-Alcázar, A., Ramos-Quiroga, J. A., Radua, J., Salavert, J., Palomar, G., Bosch, R., . . . Pomarol-Clotet, E. (2016). Brain abnormalities in adults with attention deficit hyperactivity disorder revealed by voxel-based morphometry. *Psychiatry Research: Neuroimaging*, 254, 41-47. doi:10.1016/j.psychresns.2016.06.002
- Mueller, S., Keeser, D., Samson, A. C., Kirsch, V., Blautzik, J., Grothe, M., . . . Meindl, T. (2013). Convergent findings of altered functional and structural brain connectivity in individuals with high functioning autism: A multimodal MRI study. *PLoS One*, 8(6), e67329. doi:10.1371/journal.pone.0067329
- Murdaugh, D. L., Shinkareva, S. V., Deshpande, H. R., Wang, J., Pennick, M. R., & Kana, R. K. (2012). Differential deactivation during mentalizing and classification of autism based on default mode network connectivity. *PLoS One*, 7(11), e50064. doi:10.1371/journal.pone.0050064
- Murphy, C. M., Christakou, A., Daly, E. M., Ecker, C., Giampietro, V., Brammer, M., . . . Rubia, K. (2014). Abnormal functional activation and maturation of fronto-striato-temporal and cerebellar regions during sustained attention in autism spectrum disorder. *American Journal of Psychiatry*, 171(10), 1107-1116. doi:10.1176/appi.ajp.2014.12030352
- Nakao, T., Radua, J., Rubia, K., & Mataix-Cols, D. (2011). Gray matter volume abnormalities in ADHD: Voxel-based meta-analysis exploring the effects of age and stimulant medication. *American Journal of Psychiatry*, 168(11), 1154-1163. doi:10.1176/appi.ajp.2011.11020281
- Ni, H. C., Lin, H. Y., Tseng, W. I., Chiu, Y. N., Wu, Y. Y., Tsai, W. C., & Gau, S. S. (2018). Neural correlates of impaired self-regulation in male youths with autism spectrum disorder: A voxel-based morphometry study. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 82, 233-241. doi:10.1016/j.pnpbp.2017.11.008
- Nickl-Jockschat, T., Habel, U., Michel, T. M., Manning, J., Laird, A. R., Fox, P. T., . . . Eickhoff, S. B. (2012). Brain structure anomalies in autism spectrum disorder--a

- meta-analysis of VBM studies using anatomic likelihood estimation. *Human Brain Mapping*, 33(6), 1470-1489. doi:10.1002/hbm.21299
- Nordahl, C. W., Lange, N., Li, D. D., Barnett, L. A., Lee, A., Buonocore, M. H., . . . Amaral, D. G. (2011). Brain enlargement is associated with regression in preschool-age boys with autism spectrum disorders. *Proceedings of the National Academy of Sciences of the United States of America*, 108(50), 20195-20200. doi:10.1073/pnas.1107560108
- Norman, L. J., Carlisi, C., Lukito, S., Hart, H., Mataix-Cols, D., Radua, J., & Rubia, K. (2016). Structural and functional brain abnormalities in attention-deficit/hyperactivity disorder and obsessive-compulsive disorder: A comparative meta-analysis. *JAMA Psychiatry*. doi:10.1001/jamapsychiatry.2016.0700
- Norman, L. J., Carlisi, C. O., Christakou, A., Chantiluke, K., Murphy, C., Simmons, A., . . . Rubia, K. (2017). Neural dysfunction during temporal discounting in paediatric attention-deficit/hyperactivity disorder and obsessive-compulsive disorder. *Psychiatry Research: Neuroimaging*, 269, 97-105. doi:10.1016/j.psychresns.2017.09.008
- Onnink, A. M. H., Zwiers, M. P., Hoogman, M., Mostert, J. C., Kan, C. C., Buitelaar, J., & Franke, B. (2014). Brain alterations in adult ADHD: Effects of gender, treatment and comorbid depression. *European Neuropsychopharmacology*, 24(3), 397-409. doi:10.1016/j.euroneuro.2013.11.011
- Ortal, S., van de Glind, G., Johan, F., Itai, B., Nir, Y., Iliyan, I., & van den Brink, W. (2015). The role of different aspects of impulsivity as independent risk factors for substance use disorders in patients with ADHD: A review. *Current Drug Abuse Review*, 8(2), 119-133. doi:10.2174/1874473708666150916112913
- Overmeyer, S., Bullmore, E. T., Suckling, J., Simmons, A., Williams, S. C., Santosh, P. J., & Taylor, E. (2001). Distributed grey and white matter deficits in hyperkinetic disorder: MRI evidence for anatomical abnormality in an attentional network. *Psychological Medicine*, 31(8), 1425-1435. doi:10.1017/s0033291701004706
- Passarotti, A. M., Sweeney, J. A., & Pavuluri, M. N. (2010). Neural correlates of response inhibition in pediatric bipolar disorder and attention deficit hyperactivity disorder. *Psychiatry Research: Neuroimaging*, 181(1), 36-43. doi:10.1016/j.psychresns.2009.07.002
- Pereira, A. M., Campos, B. M., Coan, A. C., Pegoraro, L. F., de Rezende, T. J. R., Obeso, I., . . . Cendes, F. (2018). Differences in cortical structure and functional mri connectivity in high functioning autism. *Frontiers in Neurology*, 9, 539. doi:10.3389/fneur.2018.00539
- Peterson, B. S., Potenza, M. N., Wang, Z., Zhu, H., Martin, A., Marsh, R., . . . Yu, S. (2009). An fMRI study of the effects of psychostimulants on default-mode processing during Stroop task performance in youths with ADHD. *American Journal of Psychiatry*, 166(11), 1286-1294. doi:10.1176/appi.ajp.2009.08050724
- Philip, R. C. M., Dauvermann, M. R., Whalley, H. C., Baynham, K., Lawrie, S. M., & Stanfield, A. C. (2012). A systematic review and meta-analysis of the fMRI investigation of autism spectrum disorders. *Neuroscience & Biobehavioral Reviews*, 36(2), 901-942. doi:10.1016/j.neubiorev.2011.10.008
- Polanczyk, G., Willcutt, E. G., Salum, G. A., Kieling, C., & Rohde, L. A. (2014). ADHD prevalence estimates across three decades: An updated systematic review and meta-regression analysis. *International Journal of Epidemiology*, 43(2), 434-442. doi:10.1093/ije/dyt261
- Poulin-Lord, M. P., Barbeau, E. B., Soulières, I., Monchi, O., Doyon, J., Benali, H., & Mottron, L. (2014). Increased topographical variability of task-related activation in perceptive and motor associative regions in adult autistics. *NeuroImage: Clinical*, 4, 444-453. doi:10.1016/j.nicl.2014.02.008
- Poustka, L., Jennen-Steinmetz, C., Henze, R., Vomstein, K., Haffner, J., & Sieltjes, B. (2012). Fronto-temporal disconnectivity and symptom severity in children with autism spectrum disorder. *The World Journal of Biological Psychiatry*, 13(4), 269-280. doi:10.3109/15622975.2011.591824

- Prat, C. S., Stocco, A., Neuhaus, E., & Kleinhaus, N. M. (2016). Basal ganglia impairments in autism spectrum disorder are related to abnormal signal gating to prefrontal cortex. *Neuropsychologia*, 91, 268-281. doi:10.1016/j.neuropsychologia.2016.08.007
- Radeloff, D., Ciaramidaro, A., Siniatchkin, M., Hainz, D., Schlitt, S., Weber, B., . . . Freitag, C. M. (2014). Structural alterations of the social brain: A comparison between schizophrenia and autism. *PLoS One*, 9(9), e106539. doi:10.1371/journal.pone.0106539
- Radua, J., Borgwardt, S., Crescini, A., Mataix-Cols, D., Meyer-Lindenberg, A., McGuire, P. K., & Fusar-Poli, P. (2012). Multimodal meta-analysis of structural and functional brain changes in first episode psychosis and the effects of antipsychotic medication. *Neuroscience & Biobehavioral Review*, 36(10), 2325-2333. doi:10.1016/j.neubiorev.2012.07.012
- Radua, J., & Mataix-Cols, D. (2009). Voxel-wise meta-analysis of grey matter changes in obsessive-compulsive disorder. *British Journal of Psychiatry*, 195(5), 393-402. doi:10.1192/bjp.bp.108.055046
- Radua, J., Mataix-Cols, D., Phillips, M. L., El-Hage, W., Kronhaus, D. M., Cardoner, N., & Surguladze, S. (2012). A new meta-analytic method for neuroimaging studies that combines reported peak coordinates and statistical parametric maps. *European Psychiatry*, 27(8), 605-611. doi:10.1016/j.eurpsy.2011.04.001
- Rae, C. L., Hughes, L. E., Weaver, C., Anderson, M. C., & Rowe, J. B. (2014). Selection and stopping in voluntary action: A meta-analysis and combined fMRI study. *NeuroImage*, 86, 381-391. doi:10.1016/j.neuroimage.2013.10.012
- Ramesh, M. G., & Rai, K. S. (2013). Region-wise gray matter volume alterations in brain of adolescents with attention deficit hyperactive disorder: A voxel based morphometric analysis. *Indian Journal of Physiology and Pharmacology*, 57(3), 270-279. Retrieved from <https://www.researchgate.net/publication/277133055>
- Rasmussen, J., Casey, B. J., van Erp, T. G., Tamm, L., Epstein, J. N., Buss, C., . . . Group, M. T. A. N. (2016). ADHD and cannabis use in young adults examined using fMRI of a Go/NoGo task. *Brain Imaging and Behavior*, 10(3), 761-771. doi:10.1007/s11682-015-9438-9
- Raznahan, A., Wallace, G. L., Antezana, L., Greenstein, D., Lenroot, R., Thurm, A., . . . Giedd, J. N. (2013). Compared to what? Early brain overgrowth in autism and the perils of population norms. *Biological Psychiatry*, 74(8), 563-575. doi:10.1016/j.biopsych.2013.03.022
- Redcay, E., & Courchesne, E. (2005). When is the brain enlarged in autism? A meta-analysis of all brain size reports. *Biological Psychiatry*, 58(1), 1-9. doi:10.1016/j.biopsych.2005.03.026
- Retico, A., Giuliano, A., Tancredi, R., Cosenza, A., Apicella, F., Narzisi, A., . . . Calderoni, S. (2016). The effect of gender on the neuroanatomy of children with autism spectrum disorders: A support vector machine case-control study. *Molecular Autism*, 7, 5. doi:10.1186/s13229-015-0067-3
- Riddle, K., Cascio, C. J., & Woodward, N. D. (2017). Brain structure in autism: A voxel-based morphometry analysis of the Autism Brain Imaging Database Exchange (ABIDE). *Brain Imaging and Behavior*, 11(2), 541-551. doi:10.1007/s11682-016-9534-5
- Riedel, A., Maier, S., Ulbrich, M., Biscaldi, M., Ebert, D., Fangmeier, T., . . . Elst, L. T. v. (2014). No significant brain volume decreases or increases in adults with high-functioning autism spectrum disorder and above average intelligence: A voxel-based morphometric study. *Psychiatry Research: Neuroimaging*, 223(2), 67-74. doi:10.1016/j.pscychresns.2014.05.013
- Riva, D., Annunziata, S., Contarino, V., Erbetta, A., Aquino, D., & Bulgheroni, S. (2013). Gray matter reduction in the vermis and crus-II is associated with social and interaction deficits in low-functioning children with autistic spectrum disorders: A VBM-DARTEL study. *Cerebellum*, 12(5), 676-685. doi:10.1007/s12311-013-0469-8

- Rojas, D. C., Peterson, E., Winterrowd, E., Reite, M. L., Rogers, S. J., & Tregellas, J. R. (2006). Regional gray matter volumetric changes in autism associated with social and repetitive behavior symptoms. *BMC Psychiatry*, 6, 56. doi:10.1186/1471-244x-6-56
- Roman-Urrestarazu, A., Lindholm, P., Moilanen, I., Kiviniemi, V., Miettunen, J., Jaaskelainen, E., . . . Murray, G. K. (2016). Brain structural deficits and working memory fMRI dysfunction in young adults who were diagnosed with ADHD in adolescence. *European Child & Adolescent Psychiatry*, 25(5), 529-538. doi:10.1007/s00787-015-0755-8
- Rubia, K. (2018). Cognitive neuroscience of attention deficit hyperactivity disorder (adhd) and its clinical translation. *Frontiers in Human Neuroscience*, 12, 100. doi:10.3389/fnhum.2018.00100
- Rubia, K., Alegria, A. A., Cubillo, A. I., Smith, A. B., Brammer, M. J., & Radua, J. (2014). Effects of stimulants on brain function in attention-deficit/hyperactivity disorder: A systematic review and meta-analysis. *Biological Psychiatry*, 76(8), 616-628. doi:10.1016/j.biopsych.2013.10.016
- Rubia, K., Cubillo, A., Smith, A. B., Woolley, J., Heyman, I., & Brammer, M. J. (2010). Disorder-specific dysfunction in right inferior prefrontal cortex during two inhibition tasks in boys with attention-deficit hyperactivity disorder compared to boys with obsessive-compulsive disorder. *Human Brain Mapping*, 31(2), 287-299. doi:10.1002/hbm.20864
- Rubia, K., Halari, R., Cubillo, A., Mohammad, A. M., Scott, S., & Brammer, M. (2010). Disorder-specific inferior prefrontal hypofunction in boys with pure attention-deficit/hyperactivity disorder compared to boys with pure conduct disorder during cognitive flexibility. *Human Brain Mapping*, 31(12), 1823-1833. doi:10.1002/hbm.20975
- Rubia, K., Halari, R., Cubillo, A., Smith, A. B., Mohammad, A. M., Brammer, M., & Taylor, E. (2011). Methylphenidate normalizes fronto-striatal underactivation during interference inhibition in medication-naïve boys with attention-deficit hyperactivity disorder. *Neuropsychopharmacology*, 36(8), 1575-1586. doi:10.1038/npp.2011.30
- Rubia, K., Halari, R., Mohammad, A. M., Taylor, E., & Brammer, M. (2011). Methylphenidate normalizes frontocingulate underactivation during error processing in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 70(3), 255-262. doi:10.1016/j.biopsych.2011.04.018
- Rubia, K., Halari, R., Smith, A. B., Mohammad, M., Scott, S., & Brammer, M. J. (2009). Shared and disorder-specific prefrontal abnormalities in boys with pure attention-deficit/hyperactivity disorder compared to boys with pure CD during interference inhibition and attention allocation. *Journal of Child Psychology and Psychiatry*, 50(6), 669-678. doi:10.1111/j.1469-7610.2008.02022.x
- Rubia, K., Smith, A. B., Brammer, M. J., Toone, B., & Taylor, E. (2005). Abnormal brain activation during inhibition and error detection in medication-naïve adolescents with ADHD. *American Journal of Psychiatry*, 162(6), 1067-1075. doi:10.1176/appi.ajp.162.6.1067
- Saad, J. F., Griffiths, K. R., Kohn, M. R., Clarke, S., Williams, L. M., & Korgaonkar, M. S. (2017). Regional brain network organization distinguishes the combined and inattentive subtypes of attention deficit hyperactivity disorder. *NeuroImage: Clinical*, 15, 383-390. doi:10.1016/j.nicl.2017.05.016
- Salazar, F., Baird, G., Chandler, S., Tseng, E., O'sullivan, T., Howlin, P., . . . Simonoff, E. (2015). Co-occurring psychiatric disorders in preschool and elementary school-aged children with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 45(8), 2283-2294. doi:10.1007/s10803-015-2361-5
- Sasayama, D., Hayashida, A., Yamasue, H., Harada, Y., Kaneko, T., Kasai, K., . . . Amano, N. (2010). Neuroanatomical correlates of attention-deficit-hyperactivity disorder accounting for comorbid oppositional defiant disorder and conduct disorder.

- Psychiatry and Clinical Neuroscience*, 64(4), 394-402. doi:10.1111/j.1440-1819.2010.02102.x
- Sato, W., Uono, S., Kochiyama, T., Yoshimura, S., Sawada, R., Kubota, Y., . . . Toichi, M. (2017). structural correlates of reading the mind in the eyes in autism spectrum disorder. *Frontiers in Human Neuroscience*, 11, 361. doi:10.3389/fnhum.2017.00361
- Schmitz, N., Rubia, K., Daly, E., Smith, A., Williams, S., & Murphy, D. G. (2006). Neural correlates of executive function in autistic spectrum disorders. *Biological Psychiatry*, 59(1), 7-16. doi:10.1016/j.biopsych.2005.06.007
- Schulz, K. P., Bédard, A. C. V., Fan, J., Clerkin, S. M., Dima, D., Newcorn, J. H., & Halperin, J. M. (2014). Emotional bias of cognitive control in adults with childhood attention-deficit/hyperactivity disorder. *NeuroImage: Clinical*, 5, 1-9. doi:10.1016/j.nicl.2014.05.016
- Schulz, K. P., Fan, J., Tang, C. Y., Newcorn, J. H., Buchsbaum, M. S., Cheung, A. M., & Halperin, J. M. (2004). Response inhibition in adolescents diagnosed with attention deficit hyperactivity disorder during childhood: An event-related fMRI study. *American Journal of Psychiatry*, 161(9), 1650-1657. doi:10.1176/appi.ajp.161.9.1650
- Schulz, K. P., Li, X., Clerkin, S. M., Fan, J., Berwid, O. G., Newcorn, J. H., & Halperin, J. M. (2017). Prefrontal and parietal correlates of cognitive control related to the adult outcome of attention-deficit/hyperactivity disorder diagnosed in childhood. *Cortex*, 90, 1-11. doi:10.1016/j.cortex.2017.01.019
- Schumann, C. M., Bloss, C. S., Barnes, C. C., Wideman, G. M., Carper, R. A., Akshoomoff, N., . . . Courchesne, E. (2010). Longitudinal magnetic resonance imaging study of cortical development through early childhood in autism. *Journal of Neuroscience*, 30(12), 4419-4427. doi:10.1523/jneurosci.5714-09.2010
- Sebastian, A., Gerdes, B., Feige, B., Kloeppel, S., Lange, T., Philipsen, A., . . . Tiescher, O. (2012). Neural correlates of interference inhibition, action withholding and action cancellation in adult ADHD. *Psychiatry Research: Neuroimaging*, 202(2), 132-141. doi:10.1016/j.psychres.2012.02.010
- Seidman, L. J., Biederman, J., Liang, L., Valera, E. M., Monuteaux, M. C., Brown, A., . . . Makris, N. (2011). Gray matter alterations in adults with attention-deficit/hyperactivity disorder identified by voxel based morphometry. *Biological Psychiatry*, 69(9), 857-866. doi:10.1016/j.biopsych.2010.09.053
- Sethi, A., Evelyn-Rahr, E., Dowell, N., Jain, S., Voon, V., Critchley, H. D., . . . Cercignani, M. (2017). Magnetization transfer imaging identifies basal ganglia abnormalities in adult ADHD that are invisible to conventional T1 weighted voxel-based morphometry. *NeuroImage: Clinical*, 15, 8-14. doi:10.1016/j.nicl.2017.03.012
- Shafritz, K. M., Bregman, J. D., Ikuta, T., & Szeszko, P. R. (2015). Neural systems mediating decision-making and response inhibition for social and nonsocial stimuli in autism. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 60, 112-120. doi:10.1016/j.pnpbp.2015.03.001
- Shafritz, K. M., Dichter, G. S., Baranek, G. T., & Belger, A. (2008). The neural circuitry mediating shifts in behavioral response and cognitive set in autism. *Biological Psychiatry*, 63(10), 974-980. doi:10.1016/j.biopsych.2007.06.028
- Shang, C. Y., Sheng, C., Yang, L. K., Chou, T. L., & Gau, S. S. (2018). Differential brain activations in adult attention-deficit/ hyperactivity disorder subtypes: A counting Stroop functional MRI study. *Brain Imaging and Behavior*, 12(3), 882-890. doi:10.1007/s11682-017-9749-0
- Shaw, P., Eckstrand, K., Sharp, W., Blumenthal, J., Lerch, J. P., Greenstein, D., . . . Rapoport, J. L. (2007). Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *Proceedings of the National Academy of Sciences of the United States of America*, 104(49), 19649-19654. doi:10.1073/pnas.0707741104
- Shimada, K., Fujisawa, T. X., Takiguchi, S., Naruse, H., Kosaka, H., Okazawa, H., & Tomoda, A. (2015). Ethnic differences in COMT genetic effects on striatal grey matter alterations associated with childhood ADHD: A voxel-based morphometry

- study in a Japanese sample. *The World Journal of Biological Psychiatry*, 1-7. doi:10.3109/15622975.2015.1102325
- Simonoff, E., Pickles, A., Charman, T., Chandler, S., Loucas, T., & Baird, G. (2008). Psychiatric disorders in children with autism spectrum disorders: prevalence, comorbidity, and associated factors in a population-derived sample. *Journal of the American Academy of Child & Adolescent Psychiatry*, 47(8), 921-929. doi:10.1097/CHI.0b013e318179964f
- Siniatchkin, M., Glatthaar, N., von Mueller, G. G., Prehn-Kristensen, A., Wolff, S., Knoechel, S., . . . Gerber, W.-D. (2012). Behavioural treatment increases activity in the cognitive neuronal networks in children with attention deficit/hyperactivity disorder. *Brain Topography*, 25(3), 332-344. doi:10.1007/s10548-012-0221-6
- Sizoo, B., van den Brink, W., Koeter, M., Gorissen van Eenige, M., van Wijngaarden-Cremers, P., & van der Gaag, R. J. (2010). Treatment seeking adults with autism or ADHD and co-morbid substance use disorder: prevalence, risk factors and functional disability. *Drug and Alcohol Dependence*, 107(1), 44-50. doi:10.1016/j.drugalcdep.2009.09.003
- Smith, A. B., Taylor, E., Brammer, M., Toone, B., & Rubia, K. (2006). Task-specific hypoactivation in prefrontal and temporoparietal brain regions during motor inhibition and task switching in medication-naïve children and adolescents with attention deficit hyperactivity disorder. *American Journal of Psychiatry*, 163(6), 1044-1051. doi:10.1176/ajp.2006.163.6.1044
- Solberg, B. S., Zayats, T., Posserud, M. B., Halmoy, A., Engeland, A., Haavik, J., & Klungsoyr, K. (2019). Patterns of psychiatric comorbidity and genetic correlations provide new insights into differences between attention-deficit/hyperactivity disorder and autism spectrum disorder. *Biological Psychiatry*, 86(8), 587-598. doi:10.1016/j.biopsych.2019.04.021
- Solomon, M., Yoon, J. H., Ragland, J. D., Niendam, T. A., Lesh, T. A., Fairbrother, W., & Carter, C. S. (2014). The development of the neural substrates of cognitive control in adolescents with autism spectrum disorders. *Biological Psychiatry*, 76(5), 412-421. doi:10.1016/j.biopsych.2013.08.036
- South, M., & Rodgers, J. (2017). Sensory, emotional and cognitive contributions to anxiety in autism spectrum disorders. *Frontiers in Human Neuroscience*, 11, 20. doi:10.3389/fnhum.2017.00020
- Spencer, M. D., Chura, L. R., Holt, R. J., Suckling, J., Calder, A. J., Bullmore, E. T., & Baron-Cohen, S. (2012). Failure to deactivate the default mode network indicates a possible endophenotype of autism. *Molecular Autism*, 3(1), 15. doi:10.1186/2040-2392-3-15
- Spinelli, S., Joel, S., Nelson, T. E., Vasa, R. A., Pekar, J. J., & Mostofsky, S. H. (2011). Different neural patterns are associated with trials preceding inhibitory errors in children with and without attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 50(7), 705-715. doi:10.1016/j.jaac.2011.03.014
- Sripada, C. S., Kessler, D., & Angstadt, M. (2014). Lag in maturation of the brain's intrinsic functional architecture in attention-deficit/hyperactivity disorder. *Proceedings of the National Academy of Sciences of the United States of America*, 111(39), 14259-14264. doi:10.1073/pnas.1407787111
- StataCorp, L. P. (2015). *Stata: Release 14 Base Reference Manual*. College Station, TX: Stata Press.
- Stevens, M. C., & Haney-Caron, E. (2012). Comparison of brain volume abnormalities between ADHD and conduct disorder in adolescence. *Journal of Psychiatry and Neuroscience*, 37(6), 389-398. doi:10.1503/jpn.110148
- Stroup, D. F., Berlin, J. A., Morton, S. C., Olkin, I., Williamson, G. D., Rennie, D., . . . Thacker, S. B. (2000). Meta-analysis of observational studies in epidemiology: A proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA*, 283(15), 2008-2012. doi:10.1001/jama.283.15.2008

- Szczepanski, S. M., & Knight, R. T. (2014). Insights into human behavior from lesions to the prefrontal cortex. *Neuron*, 83(5), 1002-1018. doi:10.1016/j.neuron.2014.08.011
- Szekely, E., Schwantes-An, T. L., Justice, C. M., Sabourin, J. A., Jansen, P. R., Muetzel, R. L., . . . Shaw, P. (2018). Genetic associations with childhood brain growth, defined in two longitudinal cohorts. *Genetic Epidemiology*, 42(4), 405-414. doi:10.1002/gepi.22122
- Tamm, L., Menon, V., Ringel, J., & Reiss, A. L. (2004). Event-related fMRI evidence of frontotemporal involvement in aberrant response inhibition and task switching in attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43(11), 1430-1440. doi:10.1097/01.chi.0000140452.51205.8d
- Tegelbeckers, J., Kanowski, M., Krauel, K., Haynes, J. D., Breitling, C., Flechtner, H. H., & Kahnt, T. (2018). Orbitofrontal signaling of future reward is associated with hyperactivity in attention-deficit/hyperactivity disorder. *Journal of Neuroscience*, 38(30), 6779-6786. doi:10.1523/jneurosci.0411-18.2018
- Thirion, B., Pinel, P., Mériaux, S., Roche, A., Dehaene, S., & Poline, J.-B. (2007). Analysis of a large fMRI cohort: Statistical and methodological issues for group analyses. *NeuroImage*, 35(1), 105-120. doi:10.1016/j.neuroimage.2006.11.054
- Thornton, S., Bray, S., Langevin, L. M., & Dewey, D. (2018). Functional brain correlates of motor response inhibition in children with developmental coordination disorder and attention deficit/hyperactivity disorder. *Human Movement Science*, 59, 134-142. doi:10.1016/j.humov.2018.03.018
- Toal, F., Daly, E. M., Page, L., Deeley, Q., Hallahan, B., Bloemen, O., . . . Murphy, D. G. M. (2010). Clinical and anatomical heterogeneity in autistic spectrum disorder: A structural MRI study. *Psychological Medicine*, 40(7), 1171-1181. doi:10.1017/s0033291709991541
- Vaidya, C. J., Foss-Feig, J., Shook, D., Kaplan, L., Kenworthy, L., & Gaillard, W. D. (2011). Controlling attention to gaze and arrows in childhood: An fMRI study of typical development and autism spectrum disorders. *Developmental Science*, 14(4), 911-924. doi:10.1111/j.1467-7687.2011.01041.x
- van Dongen, E. V., von Rhein, D., O'Dwyer, L., Franke, B., Hartman, C. A., Heslenfeld, D. J., . . . Buitelaar, J. (2015). Distinct effects of ASD and ADHD symptoms on reward anticipation in participants with ADHD, their unaffected siblings and healthy controls: A cross-sectional study. *Molecular Autism*, 6, 48. doi:10.1186/s13229-015-0043-y
- van Hulst, B. M., de Zeeuw, P., Rijks, Y., Neggers, S. F. W., & Durston, S. (2017). What to expect and when to expect it: An fMRI study of expectancy in children with ADHD symptoms. *European Child & Adolescent Psychiatry*, 26(5), 583-590. doi:10.1007/s00787-016-0921-7
- van Rooij, D., Anagnostou, E., Arango, C., Auzias, G., Behrmann, M., Busatto, G. F., . . . Buitelaar, J. K. (2018). Cortical and subcortical brain morphometry differences between patients with autism spectrum disorder and healthy individuals across the lifespan: Results from the ENIGMA ASD working group. *American Journal of Psychiatry*, 175(4), 359-369. doi:10.1176/appi.ajp.2017.17010100
- van Rooij, D., Hoekstra, P. J., Mennes, M., von Rhein, D., Thissen, A. J. A. M., Hestfeldt, D., . . . Hartman, C. A. (2015). Distinguishing adolescents with ADHD from their unaffected siblings and healthy comparison subjects by neural activation patterns during response inhibition. *American Journal of Psychiatry*, 172(7), 674-683. doi:10.1176/appi.ajp.2014.13121635
- van Wingen, G. A., van den Brink, W., Veltman, D. J., Schmaal, L., Dom, G., Booij, J., & Crunelle, C. L. (2013). Reduced striatal brain volumes in non-medicated adult ADHD patients with comorbid cocaine dependence. *Drug and Alcohol Dependence*, 131(3), 198-203. doi:10.1016/j.drugalcdep.2013.05.007
- Velasquez, F., Qin, X. A., Reilly, M. A., Neuhaus, E., Estes, A., Aylward, E., & Kleinmans, N. M. (2017). Neural correlates of emotional inhibitory control in autism spectrum disorders. *Research in Developmental Disabilities*, 64, 64-77. doi:10.1016/j.ridd.2017.03.008

- Via, E., Radua, J., Cardoner, N., Happe, F., & Mataix-Cols, D. (2011). Meta-analysis of gray matter abnormalities in autism spectrum disorder: Should asperger disorder be subsumed under a broader umbrella of autistic spectrum disorder? *Archives of General Psychiatry*, 68(4), 409-418. doi:10.1001/archgenpsychiatry.2011.27
- Vilgis, V., Sun, L., Chen, J., Silk, T. J., & Vance, A. (2016). Global and local grey matter reductions in boys with ADHD combined type and ADHD inattentive type. *Psychiatry Research: Neuroimaging*, 254, 119-126. doi:10.1016/j.psychresns.2016.06.008
- Villemonteix, T., De Brito, S. A., Kavec, M., Baleriaux, D., Metens, T., Slama, H., . . . Massat, I. (2015). Grey matter volumes in treatment naive vs. chronically treated children with attention deficit/hyperactivity disorder: A combined approach. *European Neuropsychopharmacology*, 25(8), 1118-1127. doi:10.1016/j.euroneuro.2015.04.015
- Vogan, V. M., Morgan, B. R., Lee, W., Powell, T. L., Smith, M. L., & Taylor, M. J. (2014). The neural correlates of visuo-spatial working memory in children with autism spectrum disorder: effects of cognitive load. *Journal of Neurodevelopmental Disorders*, 6, 19. doi:10.1186/1866-1955-6-19
- Waiter, G. D., Williams, J. H. G., Murray, A. D., Gilchrist, A., Perrett, D. I., & Whiten, A. (2004). A voxel-based investigation of brain structure in male adolescents with autistic spectrum disorder. *NeuroImage*, 22(2), 619-625. doi:10.1016/j.neuroimage.2004.02.029
- Wang, J., Fu, K., Chen, L., Duan, X., Guo, X., Chen, H., . . . Chen, H. (2017). Increased gray matter volume and resting-state functional connectivity in somatosensory cortex and their relationship with autistic symptoms in young boys with autism spectrum disorder. *Frontiers in Physiology*, 8, 588. doi:10.3389/fphys.2017.00588
- Wang, J., Jiang, T., Cao, Q., & Wang, Y. (2007). Characterizing anatomic differences in boys with attention-deficit/hyperactivity disorder with the use of deformation-based morphometry. *American Journal of Neuroradiology*, 28(3), 543-547. Retrieved from <http://www.ajnr.org/content/28/3/543.long>
- White, S. J., Frith, U., Rellecke, J., Al-Noor, Z., & Gilbert, S. J. (2014). Autistic adolescents show atypical activation of the brain's mentalizing system even without a prior history of mentalizing problems. *Neuropsychologia*, 56, 17-25. doi:10.1016/j.neuropsychologia.2013.12.013
- Willcutt, E. G., Sonuga-Barke, E. J. S., Nigg, J. T., & Sergeant, J. A. (2008). Recent developments in neuropsychological models of childhood psychiatric disorders. In T. Banaschewski & L. A. Rohde (Eds.), *Advances in Biological Psychiatry* (pp. 195-226). Basel: Karger. Retrieved from <https://www.karger.com/Article/Abstract/118526>
- Wilson, L. B., Tregellas, J. R., Hagerman, R. J., Rogers, S. J., & Rojas, D. C. (2009). A voxel-based morphometry comparison of regional gray matter between fragile X syndrome and autism. *Psychiatry Research: Neuroimaging*, 174(2), 138-145. doi:10.1016/j.psychresns.2009.04.013
- Yang, P., Wang, P.-N., Chuang, K.-H., Jong, Y.-J., Chao, T.-C., & Wu, M.-T. (2008). Absence of gender effect on children with attention-deficit/hyperactivity disorder as assessed by optimized voxel-based morphometry. *Psychiatry Research: Neuroimaging*, 164(3), 245-253. doi:10.1016/j.psychresns.2007.12.013
- Yang, Q., Huang, P., Li, C., Fang, P., Zhao, N., Nan, J., . . . Cui, L. B. (2018). Mapping alterations of gray matter volume and white matter integrity in children with autism spectrum disorder: evidence from fMRI findings. *Neuroreport*, 29(14), 1188-1192. doi:10.1097/wnr.0000000000001094
- Yerys, B. E., Antezana, L., Weinblatt, R., Jankowski, K. F., Strang, J., Vaidya, C. J., . . . Kenworthy, L. (2015). Neural correlates of set-shifting in children with autism. *Autism Research*, 8(4), 386-397. doi:10.1002/aur.1454
- Zamorano, F., Billeke, P., Kausel, L., Larrain, J., Stecher, X., Hurtado, J. M., . . . Aboitiz, F. (2017). Lateral prefrontal activity as a compensatory strategy for deficits of cortical processing in attention deficit hyperactivity disorder. *Scientific Reports*, 7(1), 7181. doi:10.1038/s41598-017-07681-z

Zhang, R., Geng, X., & Lee, T. M. C. (2017). Large-scale functional neural network correlates of response inhibition: An fMRI meta-analysis. *Brain Structure and Function*, 222(9), 3973-3990. doi:10.1007/s00429-017-1443-x

Figure 1






Figure 2

(A) STRUCTURAL ABNORMALITIES


FUNCTIONAL ABNORMALITIES



(B) COGNITIVE CONTROL



(C) MOTOR RESPONSE INHIBITION



Reduced in ADHD Increased in ASD Reduced in ASD Reduced in ASD and ADHD

SUPPLEMENTARY MATERIALS

Comparative Meta-analyses of Brain Structural and Functional Abnormalities during Cognitive Control in Attention-Deficit/Hyperactivity Disorder and Autism Spectrum Disorder

Steve Lukito^{1*}, PhD, Luke Norman^{1,2,3}, PhD, Christina Carlisi^{1,4}, PhD, Joaquim Radua^{5,6,7}, MD, PhD, Heledd Hart¹, PhD, Emily Simonoff¹, MD & Katya Rubia¹, PhD

¹ Department of Child and Adolescent Psychiatry; Institute of Psychiatry, Psychology and Neuroscience; King's College London; UK

² Department of Psychiatry, University of Michigan, Ann Arbor, Michigan, USA

³ The Social and Behavioral Research Branch, National Human Genome Research Institute, National Institute of Health, Bethesda, Maryland, USA

⁴ Division of Psychology and Language Sciences, University College London, UK

⁵ Department of Psychosis Studies; Institute of Psychiatry, Psychology and Neuroscience; King's College London; UK

⁶ Imaging of mood- and anxiety-related disorders (IMARD) group, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), CIBERSAM, Barcelona, Spain

⁷ Centre for Psychiatric Research and Education, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

* Correspondence and reprints request:

Dr Steve Lukito

Department of Child and Adolescent Psychiatry; Institute of Psychiatry, Psychology and Neuroscience, King's College London, 16 De Crespigny Park, London, SE5 8AF, UK

Contents

Supplement 1: Literature Search and Methods	3
Literature Search.....	3
Seed-based <i>d</i> Mapping (SDM).....	4
Figure S1. Flow Diagram of Literature Search and Study Selection Process	5
Supplement 2: Full Characteristic Tables and Summary Findings of ADHD and ASD	
Neuroimaging Studies	6
Table S2a. Sample Characteristics and Summary Findings of VBM and fMRI Cognitive Control Studies in ADHD	6
Table S2b. Sample Characteristics and Summary Findings of VBM and fMRI Cognitive Control Studies in ASD	12
Supplement 3: Meta-analyses in Pediatric Sample with ADHD and ASD	16
Table S3a. Characteristics of Overall Pediatric Sample and Age-, Sex- and IQ- matched Subgroups	16
Table S3b. Structural and Functional Abnormalities in Pediatric Sample with ADHD and ASD	17
Figure S3. Structural and Functional Abnormalities in Pediatric Sample with ADHD and ASD	19
Supplement 4: Meta-analyses in Adults with ADHD and ASD	20
Table S4a. Characteristics of Adult Overall Sample and Subgroups Matched on Age, Sex and IQ 20	
Table S4b. Structural and Functional Abnormalities in Adults with ADHD and ASD.....	21
Figure S4. Structural and Functional Abnormalities in Adults with ADHD and ASD	24
Supplement 5: Associations between Brain Abnormalities and Age.....	25
Correlational Analyses between Brain Abnormalities and Age.....	25
Table S5. Spearman's Correlations between Disorder-Differentiating or Shared Abnormalities and Age	26
Additional Analyses Covarying for Age in the Sub-Meta-Analyses with Participants Matched on Age, Sex and IQ.....	27
Supplement 6: Associations between Brain Abnormalities and Psychostimulant Exposure	28
Figure S6a. Association between VBM Abnormalities and Current Psychostimulant Exposure in ADHD	28
Figure S6b. Association between fMRI Abnormalities and Lifetime Psychostimulant Exposure in ADHD	29
Supplement 7: Influences of Task Type or Performance in Disorder-differentiating Findings ..	30
Table S7. Task Types and Behavioural Performance Impairments in ADHD and ASD vs. TD	30
Comparing Task Type and Performance in ADHD and ASD vs. TD	31
Influences of Task Type and Performance in the Disorder-differentiating Findings during Cognitive Control.....	32
Influences of Performance in the Disorder-differentiating Findings during Motor Response Inhibition	33
Supplement 8: Jack-knife Analyses in Brain Structure and Function Abnormalities	34
Table S8a. Jack-knife Analyses of Reduced GMV Clusters among ADHD VBM Studies.....	34
Table S8b. Jack-knife Analyses of GMV Abnormalities among ASD VBM Studies	35
Table S8c. Jack-knife Analyses of Abnormal Brain Function during Cognitive Control among ADHD fMRI Studies.....	36
Table S8d. Jack-knife Analyses of Abnormal Brain Function during Cognitive Control among ASD fMRI Studies.....	37
Table S8e. Jack-knife Analyses of Abnormal Brain Function during Motor Response Inhibition among ADHD fMRI Studies	38
Table S8f. Jack-knife Analyses of Abnormal Brain Function during Motor Response Inhibition among ASD fMRI Studies	39
Figure S8. Pictorial Representation of the Jack-knife Analyses	40
References	41

Supplement 1: Literature Search and Methods

Literature Search

A comprehensive literature search was conducted by SL, LN, and CC in PubMed, Scopus, Web of Science and ScienceDirect to identify whole-brain functional magnetic resonance imaging (fMRI) or voxel-based morphometry (VBM) studies comparing individuals with ASD or ADHD against typical controls. The searches were conducted up to 30th November 2018 and used key words related to: (1) ASD, i.e., autism OR autistic OR Asperger OR ASD OR autism spectrum disorder OR pervasive developmental disorder; (2) ADHD, i.e., hyperkinetic OR ADHD or attention-deficit/hyperactivity disorder; (3) VBM; (4) inhibitory function, i.e., inhibition OR stop OR Stroop OR flanker OR go/no-go OR Simon OR interference OR executive function OR switch; and (5) neuroimaging, i.e., fMRI OR MRI.

The following syntaxes were used on PubMed to search for (1) ADHD VBM studies:

("ADHD"[Title/Abstract] OR "attention-deficit/hyperactivity disorder"[Title/Abstract] OR "hyperkinetic"[Title/Abstract]) AND ("VBM"[Title/Abstract] OR "voxel-based morphometry"[Title/Abstract]) and (2) ASD VBM studies: ("Autism"[Title/Abstract] OR "autistic"[Title/Abstract] OR "autism spectrum disorders"[Title/Abstract] OR "pervasive developmental disorders"[Title/Abstract] OR "ASD"[Title/Abstract] OR "PDD"[Title/Abstract] OR "Asperger"[Title/Abstract]) AND ("VBM"[Title/Abstract] OR "voxel-based morphometry"[Title/Abstract]).

The following syntaxes were used on PubMed to search for (1) ADHD fMRI studies:

((("ADHD"[Title/Abstract] OR "hyperkinetic"[Title/Abstract] OR "attention-deficit/hyperactivity disorders"[Title/Abstract])) AND ("inhibition"[Title/Abstract] OR "stop" [Title/Abstract] OR "stroop"[Title/Abstract] OR "flanker "[Title/Abstract] OR " go/no-go "[Title/Abstract] OR " Simon"[Title/Abstract] OR "interference"[Title/Abstract] OR "executive function"[Title/Abstract] OR " switch"[Title/Abstract])) AND ("MRI"[Title/Abstract] OR "fMRI"[Title/Abstract]), and (2) ASD fMRI studies: ((("Autism"[Title/Abstract] OR "autistic"[Title/Abstract] OR "autism spectrum disorders"[Title/Abstract] OR "pervasive developmental disorders"[Title/Abstract] OR "ASD"[Title/Abstract] OR "PDD"[Title/Abstract]

OR "Asperger"[Title/Abstract])) AND ("inhibition"[Title/Abstract] OR "stop" [Title/Abstract] OR "stroop"[Title/Abstract] OR "flanker "[Title/Abstract] OR " go/no-go "[Title/Abstract] OR " Simon"[Title/Abstract] OR "interference"[Title/Abstract] OR "executive function"[Title/Abstract] OR " switch"[Title/Abstract])) AND ("MRI"[Title/Abstract] OR "fMRI"[Title/Abstract]).


Only predefined contrasts comparing inhibitory control (i.e., stop, no-go, incongruent or switch trials) against a control condition (i.e., failed stop, go, oddball, congruent or repeated trials) were included in the fMRI meta-analysis. Data extracted included age, sex, IQ, current and historic exposure of psychostimulant medications and comorbidity of ASD and ADHD in the counterpart disorder groups. From the database searches, 2484 records were retrieved, and eight further records were identified from past meta-analyses. After duplicates were removed, 1290 records remained and were screened, yielding 243 full-text articles assessed for eligibility, and 140 included in the meta-analysis. Of the full-text articles excluded, 30 included samples already used in previous publications, eight included fewer than ten participants, 18 had no control participants, 26 were not whole-brain studies and 21 did not include eligible contrast.

Seed-based *d* Mapping (SDM)

The anisotropic seed-based *d* Mapping AES-SDM (www.sdmproject.com) is meta-analytic software for neuroimaging data. The software takes both statistical parametric maps of MRI/fMRI contrasts (Radua et al., 2012), and the conventional peak coordinates and effect sizes (*t*-scores) as input data. When provided with the latter type of data, AES-SDM estimates signed (positive/negative) effect size and variance maps that represent the difference of brain activations or gray matter structures between patient and control groups. These maps are computed by convolving an anisotropic non-normalized Gaussian kernel with Hedges effect size of each peak. Thus, voxels correlated with peaks are assigned higher effect sizes. Combinations of maps between studies are based on a random-effects model, accounting for sample size, within-study variability and between-study heterogeneity. An update of AES-SDM software also allows adjustment for correlated datasets, accounting

for shared variance of brain activations or structure across datasets (Norman et al., 2016), e.g., when pairs of datasets investigating two inhibitory tasks take places in the same or largely overlapping sample.

Figure S1. Flow Diagram of Literature Search and Study Selection Process



Flow diagram describing the literature search, which included the databases PubMed, ScienceDirect, Scopus and Web of Knowledge

Supplement 2: Full Characteristic Tables and Summary Findings of ADHD and ASD Neuroimaging Studies

Table S2a. Sample Characteristics and Summary Findings of VBM and fMRI Cognitive Control Studies in ADHD

Source	Age group	Task	Patients						Controls			Summary findings
			N (% male)	Age, Range, y	IQ	Life Stim %	Cur Stim %	Com ASD %	N (% male)	Age, y Range, y	IQ	
(A) VBM studies in ADHD												
Ahrendts et al. (2011)	Adult	--	31 (65)	31.2 (18-55)	n/a	3	0	n/a	31 (65)	31.5 (19-52)	n/a	ADHD<TD: L/R occipital lobe
Montes et al. (2010)	Adult	--	20 (50)	29.0 (25-35)	102.9	0	0	n/a	20 (50)	27.6 (25-35)	100.2	ADHD<TD: R caudate
Amico, Stauber, Koutsouleris, and Frodl (2011)	Adult	--	20 (75)	33.6 (n/a)	n/a	n/a	30	n/a	20 (75)	34.7 (n/a)	n/a	--
Bonath, Tegelbeckers, Wilke, Flechtner, and Krauel (2016)	Pediat.	--	18 (100)	13.6 (11-17)	106.8	78	56	n/a	18 (100)	14.1 (11-17)	108.1	ADHD<TD: L/R cerebellum, occipital cortex, L/R hippocampus/ amygdala complex, L ACC
Bralten et al. (2016)	Mixed	--	307 (68)	17.1 (8-30)	97.1	89	n/a	n/a	196 (49)	16.7 (8-30)	106.6	ADHD<TD: L pre-CG, L OFC, R frontal pole, L paracingulate/cingulate cortex/frontal pole, bilateral medial FG/ paracingulate/cingulate/ subcallosal cortices
Brieber et al. (2007)	Pediat.	--	15 (100)	13.1 (10-16)	104.1	n/a	67	0	15 (100)	13.3 (10-16)	107.7	ADHD<TD: L/R caudate nucleus, L/R hippocampus, R MFG, L insula, L STG, L MOG; ADHD>TD: L/R SPL, post-CG, L middle cingulate, precuneus, L IPL
Carmona et al. (2005)	Pediat.	--	25 (84)	10.8 (6-16)	n/a	100	100	n/a	25 (84)	11.2 (6-16)	n/a	ADHD<TD: L pre-/post-CG, paracentral lobule, R rectal gyrus, R inferior and superior orbital frontal, L/R cerebellum
Depue, Burgess, Bidwell, Willcutt, and Banich (2010)	Adult	--	31 (61)	20.0 (n/a)	114.2	77	77	n/a	21 (65)	19.3 (n/a)	112.6	--
Gehricke et al. (2017)	Adult	--	32 (81)	25.3 (n/a)	n/a	n/a	25	n/a	40 (83)	23.9 (n/a)	n/a	ADHD<TD: R M/IFG, R S/MTG, L caudate head, L PHG
He et al. (2015)	Pediat.	--	37 (100)	9.9 (7-16)	n/a	0	0	n/a	35 (100)	10.7 (8-15)	n/a	ADHD<TD: R OFC, R PMC, L PCC, L posterior midcingulate cortex
Iannaccone, Hauser, Ball, et al. (2015)	Pediat.	--	18 (50)	14.5 (12-16)	114.5	72	72	n/a	18 (61)	14.8 (12-16)	108.5	ADHD<TD: R medial SFG/ACC/SFG, L SMA/cingulate gyrus, L cerebellum; ADHD>TD: R pre- and post-CG
Jagger-Rickels, Kibby, and Constance (2018)	Pediat.	--	41 (44)	9.6 (8-12)	n/a	n/a	n/a	n/a	32 (56)	9.7 (8-12)	n/a	ADHD<TD: L MTG/STS, R STS L/R caudate/ putamen, L/R SFG, L/R insula, L/R thalamus, R medial OFC, L MFG, L/R occipital lobe, L medial SFG, L pre-CG, R post cingulate/anterior lingual, R calcarine/cuneus, L supramarginal gyrus

Johnston et al. (2014)	Pediat.	--	34 (100)	12.5 (n/a)	99.8	n/a	29	n/a	34 (100)	13.2 (n/a)	103.7	ADHD<TD: Brainstem, putamen
Kappel et al. (2015)	Pediat.	--	14 (71)	9.8 (8-12)	104.6	0	0	n/a	10 (80)	11.0 (8-12)	111.9	ADHD<TD: R STG, R Heschl's gyrus, R rolandic operculum; ADHD>TD: L paracentral lobule, L/R middle orbital gyrus, R FFG, L rectal gyrus
	Adult	--	16 (94)	23.5 (19-31)	97.8	38	0	n/a	20 (100)	23.7 (19-31)	108.4	ADHD<TD: R supramarginal gyrus, R precuneus, L OFG, R hippocampus, L rectal gyrus
Kaya et al. (2018)	Pediat.	--	19 (74)	10.3 (7-14)	113.5	0	0	n/a	18 (67)	10.2 (6-14)	119.7	ADHD>TD: L/R SFG, R MFG, L/R SMA, L/R pre-CG, L post-CG, R MOG, L cuneus
Kobel et al. (2010)	Pediat.	--	14 (100)	10.4 (9-13)	n/a	100	100	n/a	12 (100)	10.9 (9-13)	n/a	ADHD<TD: R S/MTG
Kumar, Arya, and Agarwal (2017)	Pediat.	--	18 (100)	9.6 (7-13)	92.1	n/a	0	0	18 (100)	9.7 (7-13)	109.7	ADHD<TD: L OFC, L MFG/ dIPFC, L MTG, L cerebellum
Li et al. (2015)	Pediat.	--	30 (100)	10.3 (8-14)	121.7	0	0	n/a	30 (100)	10.3 (8-14)	107.1	ADHD<TD: R insula, R OFC
Lim et al. (2015)	Pediat.	--	44 (100)	13.6 (10-18)	92.2	18	14	0	33 (100)	14.3 (10-18)	110	ADHD<TD: L/R cerebellum, L IPL, pre-/post-CG
Maier et al. (2015)	Adult	--	131 (48)	34.5 (18-58)	113.1	27	0	0	95 (47)	37.7 (n/a)	121	--
McAlonan et al. (2007)	Pediat.	--	28 (100)	9.9 (6-13)	109.9	100	100	n/a	31 (100)	9.6 (6-13)	116.5	ADHD<TD: R MFG, L/R SFG/IFG, R globus pallidus/putamen/ thalamus, precuneus, L IPL, L SOG, R cerebellum vermis
Moreno-Alcázar et al. (2016)	Adult	--	44 (66)	31.6 (18-54)	105.0	66	66	n/a	44 (66)	32.6 (18-54)	106.0	ADHD<TD: R SMA/SFG, subgenual ACC, R DLPFC
Onnink et al. (2014)	Adult	--	119 (39)	36.3, (n/a)	107.5	80	69	n/a	107 (42)	36.9 (n/a)	110.2	--
Overmeyer et al. (2001)	Pediat.	--	18 (83)	10.4 (8-13)	99	94	94	n/a	16 (94)	10.3 (7-14)	n/a	ADHD<TD: R PCC, R SFG, L/R putamen/globus pallidus
Ramesh and Rai (2013)	Pediat.	--	15 (26)	16.8 (11-20)	n/a	n/a	n/a	n/a	15 (26)	16.7 (11-20)	n/a	ADHD<TD: L cingulate gyrus, R mid-cingulate gyrus, L SFG, L/R medial SFG, L temporal lobe, L MOG, L cuneus, L STG, L supramarginal gyrus
Roman-Urrestarazu et al. (2016)	Adult	--	49 (76)	22.2 (20-24)	96.6	2	2	0	34 (50)	22.9 (20-24)	112.2	ADHD<TD: L/R caudate
Saad et al. (2017)	Pediat.	--	16 (75)	12.8 (8-17)	n/a	38	38	n/a	28 (68)	13.1 (8-17)	n/a	--
	Pediat.	--	18 (72)	13.7 (8-17)	n/a			n/a	28 (68)	13.1 (8-17)	n/a	--
Sasayama et al. (2010)	Pediat.	--	18 (72)	10.6 (6-16)	90	83	72	0	17 (71)	10.0 (6-14)	n/a	ADHD<TD: L/R amygdala/temporal pole/OFC, L/R occipital cortex, R STS, L parietal cortex, L MFG, L temporal pole, R rectal gyrus, L PHG

Seidman et al. (2011)	Adult	--	74 (51)	37.3 (18-59)	116	69	28	n/a	54 (46)	34.3 (18-59)	115.8	--
Sethi et al. (2017)	Adult	--	30 (63)	33.7 (18-65)	109.0	100	100	n/a	30 (63)	32.6 (18-65)	110.1	ADHD<TD: R IPC
Shimada et al. (2015)	Pediat.	--	17 (88)	10.3 (n/a)	95.3	n/a	n/a	n/a	15 (73)	12.8 (n/a)	104.1	ADHD<TD: L putamen
Stevens and Haney-Caron (2012)	Pediat.	--	24 (67)	15.7 (12-18)	n/a	n/a	n/a	n/a	24 (70)	16.0 (12-18)	n/a	--
van Wingen et al. (2013)	Adult	--	14 (100)	32.0 (22-50)	104	0	0	n/a	15 (100)	37.0 (22-50)	99	ADHD<TD: R putamen, R cerebellum; ADHD>TD: L/R midbrain, R pre-CG
Vilgis, Sun, Chen, Silk, and Vance (2016)	Pediat.	--	48 (100)	12.6 (8-17)	92.2	25	25	n/a	31 (100)	12.8 (8-17)	109.6	ADHD<TD: R SPL, L IPL, L/R S/MFG, medial FG, R MTG
Villemontheix et al. (2015)	Pediat.	--	33 (55)	10.3 (7-13)	105.6	0	0	n/a	24 (50)	10 (7-13)	109.7	ADHD<TD: R insula, R MTG,
	Pediat.	--	20 (80)	10.4 (7-13)	107.4	100	100	n/a	24 (50)	10 (7-13)	109.7	ADHD<TD: R MFG, R pre-CG
Wang, Jiang, Cao, and Wang (2007)	Pediat.	--	12 (100)	13.4 (n/a)	n/a	n/a	0	n/a	12 (100)	13.5 (n/a)	n/a	ADHD<TD: L SPL, R MFG, R MTG, R BG; ADHD>TD: R occipital lobe
P. Yang et al. (2008)	Pediat.	--	57 (61)	11.1 (7-17)	97.9	n/a	86	n/a	57 (60)	11.7 (7-17)	n/a	ADHD<TD: L/R caudate, L/R cerebellum, R calcarine sulcus, L cuneus
(B) fMRI cognitive control studies in ADHD												
Banich et al. (2009)	Adult	Stroop	23 (61)	20.0 (n/a)	116	87	61	n/a	23 (57)	19.0 (n/a)	113	ADHD>TD: R MFG
Bhajiwal, Chevrier, and Schachar (2014)	Pediat.	Stop	12 (58)	13.8 (9-18)	n/a	100	100	0	12 (50)	15.4 (9-18)	n/a	ADHD<TD: R MFG, R MTG, L cerebellum; ADHD>TD: R MFG/SFG
Booth et al. (2005)	Pediat.	GNG	12 (67)	11.0 (9-12)	n/a	100	100	n/a	12 (58)	11.7 (9-12)	n/a	ADHD<TD: L/R pre-CG, L/R caudate, R caudate head, R IFG, L/R thalamus.
Carmona et al. (2012)	Adult	GNG	19 (100)	33.6 (n/a)	110.9	0	0	n/a	19 (100)	29.4 (n/a)	111.7	ADHD<TD: R STG
Chantiluke, Barrett, Giampietro, Santosh, et al. (2015)	Pediat.	Stop	18 (100)	14.3 (10-17)	95	78	61	0	25 (100)	13.4 (10-17)	109	ADHD<TD: L OFC/STL/putamen/GP, L IPL
Chen et al. (2015)	Adult	GNG	29 (100)	24.9 (n/a)	n/a	0	0	n/a	25 (100)	25.6 (n/a)	n/a	--
Chou, Chia, Shang, and Gau (2015)	Pediat.	Stroop	42 (81)	10.5 (7-17)	108.5	0	0	n/a	20 (80)	12.0 (8-17)	106.5	ADHD<TD: R MFG/IFG (atomoxetine group pre-treatment), L IFG, L SPL, R medial FG (methylphenidate group pre-treatment); ADHD>TD: L/R SPL, pre-CG, L dlPFC, L dACC (atomoxetine group pre-treatment), post-CG (methylphenidate group pre-treatment)
Congdon et al. (2010)	Adult	Stop	35 (54)	30.9 (21-50)	n/a	n/a	29	n/a	62 (45)	30.8 (21-50)	n/a	--

Cubillo et al. (2010) ^a	Adult	Stop, Switch	11 (100)	29.0 (26-30)	92	0	0	n/a	14 (100)	28.0 (26-30)	106	ADHD<TD: L IFC/AI/PMC, R IFC/AI/caudate/thalamus/putamen/PMC, L/R SMA/ACC (stop), R IFG/insula/caudate/ putamen L IFG/insula/ putamen/pre-/post-CG/inferior parietal (switch)
Cubillo et al. (2014)	Pediat.	Stop	19 (100)	13.1 (10-17)	92	0	0	0	29 (100)	13.8 (10-17)	110	ADHD<TD: L/R IFG, L M/ITG/parietal gyri, R cerebellum/FFG; ADHD>TD: L cerebellum, L/R PCC/occipital gyri, R STG/post-CG/posterior insula/putamen
Cubillo, Halari, Giampietro, Taylor, and Rubia (2011) ^a	Adult	Simon	11 (100)	29.0 (26-30)	92	0	0	n/a	15 (100)	28.0 (26-30)	112	ADHD<TD: L OFC/IFC/mPFC/ACC/ caudate/PMC
Dibbets, Evers, Hurks, Marchetta, and Jolles (2009) ^b	Adult	GNG	16 (100)	28.9 (21-42)	n/a	88	88	n/a	13 (100)	28.1 (21-41)	n/a	--
Dibbets, Evers, Hurks, Bakker, and Jolles (2010) ^b	Adult	Switch	15 (100)	28.8 (21-42)	n/a	80	80	n/a	14 (100)	28.6 (21-41)	n/a	ADHD<TD: R putamen, R dorsal posterior CG, L mPFC, R thalamus, L OFC/ claustrum/post-CG; ADHD>TD: R MTG, R dACC, R precuneus, R lingual gyrus, L pre-CG/SMA, L insula
Durston, Mulder, Casey, Ziermans, and van Engeland (2006)	Pediat.	GNG	11 (100)	14.0 (8-20)	100	82	55	n/a	11 (100)	15.3 (8-20)	106	--
L. Y. Fan, Chou, and Gau (2017) ^c	Adult	Stroop	12 (42)	28.9 (n/a)	115.8	0	0	n/a	12 (42)	30.3 (n/a)	118.3	ADHD>TD: R IFG, RACC
	Adult	Stroop	12 (42)	32.5 (n/a)	119.9	0	0	n/a	12 (42)	30.3 (n/a)	118.3	ADHD>TD: R IFG
L. Y. Fan, Shang, Tseng, Gau, and Chou (2018)	Pediat.	Stroop	27 (89)	12.1 (9-15)	105.2	0	0	n/a	27 (78)	11.8 (9-16)	110.4	ADHD < TD: R MFG, R post-CG, R SPL
Hwang et al. (2015)	Pediat.	Stroop	26 (65)	13.9 (n/a)	106.4	n/a	42	0	35 (51)	14.5 (n/a)	105.1	ADHD < TD: Medial FG, cerebellar tonsil
Iannaccone, Hauser, Staempfli, et al. (2015)	Pediat.	GNG/Flanker	18 (50)	14.5 (12-16)	108.5	72	72	n/a	18 (61)	14.8 (12-16)	114.4	ADHD<TD: R MFG/dIPFC/SFG; ADHD>TD: L/R MTG/STG, L insula/putamen/pre-CG, L ITG/FFG/PHG/MTG, L pons, L amygdala/ hippocampus, L FFG/ lingual gyrus, L post-CG/IPG, R brainstem/ pons/cerebellum
Janssen, Heslenfeld, Mourik, Logan, and Oosterlaan (2015)	Pediat.	Stop	21 (90)	10.6 (8-13)	98.6	90	90	n/a	17 (76)	10.3 (8-13)	108.7	ADHD<TD: L/R mPFC, R MFG, R IFG, R pre-CG, R ACC, L SFG; ADHD>TD: L/R ACC, L/R MFG, R mPFC, R PCC, R cuneus, L IFG, L post-CG, L pre-CG.
Konrad, Neufang, Hanisch, Fink, and Herpertz-Dahlmann (2006)	Pediat.	Flanker	16 (100)	10.2 (8-12)	103	0	0	0	16 (100)	10.3 (8-12)	105	ADHD<TD: L pre-CG, R putamen; ADHD>TD: L medial SPL
Kooistra et al. (2010)	Adult	GNG	11 (100)	21.5 (18-25)	110	100	0	n/a	11 (100)	10.1 (18-25)	125	ADHD>TD: L/R ACC, R supramarginal gyrus, L angular gyrus, L/R supramarginal gyrus, R occipital gyrus, L frontal pole, L/R MFG, L/R IFG, L SFG, paracingulate gyrus, medial thalamus, L caudate nucleus, L posterior putamen,

												L lateral occipital gyrus, L/R pre-CG, R post-CG, R mPFC, L PCC, L STG
J. Ma et al. (2012)	Pediat.	GNG	15 (53)	9.8 (8-12)	100.2	27	0	n/a	15 (53)	22.3 (8-12)	102.6	ADHD>TD: R ITF, R midbrain, R pre-CG, R calcarine, R IOG, R MOG, L/R cerebellum, L post-CG, L IFG, R hippocampus
I. Ma et al. (2016)	Pediat.	Stroop	25 (76)	15.4 (14-17)	98.3	n/a	60	n/a	33 (67)	15.3 (14-17)	108.9	--
Massat et al. (2018)	Pediat.	Stop	18 (44)	10.0 (8-12)	106.8	0	0	n/a	19 (47)	10.6 (8-12)	114.4	ADHD>TD: L caudate body, L ventral putamen R dorsal putamen, R insula, L caudate tail, R middle CC, R ACC
Passarotti et al. (2010)	Pediat.	Stop	11 (55)	13.1 (10-18)	107.6	48	0	n/a	15 (48)	9.9 (10-18)	101.2	ADHD<TD: R MFG, R mPFC, L/R I/SFG, L STG; ADHD>TD: R caudate tail, L caudate, L anterior cerebellar vermis
Peterson et al. (2009)	Pediat.	Stroop	16 (81)	14.1 (7-18)	101.2	100	0	n/a	20 (60)	14.1 (7-18)	118.5	ADHD<TD: L ACC, insula R, precuneus, thalamus, caudate; ADHD>TD: R SFG, hippocampus, L ACC
Rasmussen et al. (2016)	Adult	GNG	50 (82)	24.8 (n/a)	102.1	n/a	4	n/a	23 (70)	24.1 (n/a)	109.2	ADHD<TD: R supramarginal gyrus, superior parietal, angular gyrus, L/R MFG. R frontal pole, L/R SPL, L/R IFG, L/R pre-/post-CG, R caudate/ accumbens, L/R precuneus, R OFC, R posterior cingulate gyrus, R insula, R thalamus
Rubia, Smith, Brammer, Toone, and Taylor (2005)	Pediat.	Stop	16 (100)	13.2 (9-16)	100	0	0	n/a	21 (100)	13.4 (9-16)	95	ADHD<TD: R IFC, OFC, pre-CG, STL
Rubia, Halari, Cubillo, et al. (2011) ^d	Pediat.	Simon	12 (100)	13.0 (10-15)	90	0	0	0	13 (100)	14.0 (11-16)	102	ADHD<TD: R IFC/inferior parietal, L vmPFC/BG/thalamus, R SMA/ACC/PCC/SPL, L STL/MTL/occipital
Rubia, Halari, Mohammad, Taylor, and Brammer (2011) ^d	Pediat.	Stop	12 (100)	13.0 (10-15)	90	0	0	0	13 (100)	13.0 (11-16)	102	ADHD<TD: B IFC, insula, ACC, pre-SMA; thalamus; R MTL, occipital, IPL, precuneus, PCC, cerebellum
Schulz et al. (2004)	Pediat.	GNG	10 (100)	17.9 (15-19)	88.4	70	0	0	9 (100)	17.5 (16-19)	91.9	ADHD<TD: R pre-CG, R ITG, L hippocampus, R lingual gyrus, L/R cerebellum; ADHD>TD: L/R MFG, L/R IFG, L medial FG, L ACC, L/R IPL, R precuneus
Schulz et al. (2014)	Adult	GNG	14 (100)	23.3 (19-27)	n/a	71	0	0	14 (100)	22.8 (18-26)	n/a	--
Schulz et al. (2017)	Adult	Stroop	27 (89)	24.2 (21-28)	n/a	74	0	0	28 (86)	24.6 (21-28)	n/a	ADHD<TD: L/R cerebellum, L MTG, R lingual gyrus, R IFG, R MTG, R OFG, R SOG, R precuneus, R IPL. R MOG
Sebastian et al. (2012) ^e	Adult	GNG Stroop Stop	20 (55)	33.3 (n/a)	115.3	0	0	n/a	24 (46)	17.5 (n/a)	115.7	ADHD<TD: R caudate (GNG), R GP (Stop), R post-CG, L para-central lobule/mid-CC, L STG/MTG, R temporal pole, R GP (Stroop)
Shang, Sheng, Yang, Chou, and Gau (2018) ^f	Adult	Stroop	25 (48)	29.1 (17-44)	112.8	0	0	n/a	30 (50)	28.2 (20-42)	115.4	ADHD<TD: L IFG, L/R caudate nucleus
	Adult	Stroop	25 (56)	28.5 (17-50)	113.1	0	0	n/a	30 (50)	28.2 (20-42)	115.4	ADHD<TD: L IFG, L caudate nucleus, L SPL
Siniatchkin et al. (2012)	Pediat.	GNG	12 (83)	9.3 (7-13)	n/a	n/a	n/a	n/a	12 (75)	30.3 (7-13)	n/a	ADHD<TD: ACC, R dIPFC, R caudate nucleus

Smith, Taylor, Brammer, Toone, and Rubia (2006) ⁹	Pediat.	GNG	17(100)	12.8 (n/a)	n/a	0	0	0	18 (100)	9.3 (n/a)	n/a	ADHD<TD: L MFG
		Stroop	19 (100)	12.9 (n/a)	n/a	0	0	0	24 (100)	12.8 (n/a)	n/a	--
		Switch	14 (100)	13.3 (n/a)	n/a	0	0	0	27 (100)	12.9 (n/a)	n/a	ADHD<TD: L STG/IFG/pre-CG/insula, R MTG/IFG, IPL/STG
Spinelli et al. (2011)	Pediat.	GNG	13 (69)	10.6 (8-13)	109.2	n/a	15	n/a	17 (47)	13.3 (8-13)	108.8	ADHD>TD: R pre-CG, MFG
Szekely et al. (2018)	Adult	Stop	64 (56)	24.0 (n/a)	n/a	n/a	62.5	n/a	84 (57)	24.5 (n/a)	n/a	--
Tamm, Menon, Ringel, and Reiss (2004)	Pediat.	GNG	10 (100)	16.0 (14-18)	109.2	n/a	50	n/a	12 (100)	10.6 (14-16)	111.6	ADHD<TD: R ACC/SMA/ MFG; ADHD>TD: L M/STG
Thornton, Bray, Langevin, and Dewey (2018)	Pediat.	GNG	20 (90)	12.4 (8-17)	109.4	n/a	n/a	0	20 (40)	10.6 (8-17)	112.6	--
van Hulst, de Zeeuw, Rijks, Neggers, and Durston (2017)	Pediat.	GNG	24 (100)	11.2 (8-12)	105.6	n/a	n/a	n/a	26 (100)	10.5 (8-12)	117.3	--
van Rooij et al. (2015)	Pediat.	Stop	108 (64)	15.1 (8-17)	92.7	75	n/a	n/a	77 (49)	16.0 (9-17)	109.2	ADHD<TD: L STG, pre-CG; ADHD>TD: L SMA
	Adult	Stop	77 (78)	20.3 (18-25)	99.1	79	n/a	n/a	45 (33)	14.6 (18-23)	106.4	ADHD<TD: L SFG, R hippocampal gyrus
Zamorano et al. (2017)	Pediat.	Stroop	17 (100)	11.6 (10-12)	104.2	100	100	n/a	17 (100)	11.7 (10-12)	109.8	ADHD>TD: R medial/IFG

Abbreviations. N=sample size, y=year, pediat.=pediatric (child/adolescent) sample, life/cur stim exp=lifetime/current stimulant exposure, com ASD=comorbid autism spectrum disorder based on explicit reporting or study exclusion criteria, ADHD=attentional-deficit/hyperactivity disorder, TD=typically developing controls, L/R=Left/Right, GNG = Go/No-Go. Brain region (in alphabetical order): ACC=anterior cingulate cortex, BG=basal ganglia, dACC=dorsal ACC, dlPFC=dorsolateral prefrontal cortex, FFG=fusiform gyrus, I/M/medial/SFG=inferior/middle/medial/superior frontal gyrus, I/M/SOG=inferior/middle/superior occipital gyrus, I/M/STG=inferior/middle/superior temporal gyrus, I/SPL=inferior/superior parietal lobe, mPFC=medial prefrontal cortex, OFC=orbital frontal cortex, OFG=orbital frontal gyrus, PCC=posterior cingulate cortex, pre-/post-CG=pre-/post-central gyrus, PHG=parahippocampal gyrus, PMC=premotor cortex, SMA=supplementary motor area, STS=superior temporal sulcus

Table S2b. Sample Characteristics and Summary Findings of VBM and fMRI Cognitive Control Studies in ASD

Source	Age group	Task	Patients				Controls			Summary findings
			N (% male)	Age (Range), y	IQ	Com ADH D %	N (% male)	Age (Range), y	IQ	
(A) VBM studies in ASD										
Abell et al. (1999)	Adult	--	15 (80)	28.8 (n/a)	n/a	n/a	15 (80)	25.3 (n/a)	n/a	ASD<TD: R paracingulate, L IFG, L occipito-temporal junction; ASD>TD: L amygdala, L/R anterior cerebellum, L MTG, R ITG
Boddaert et al. (2004)	Pediat.	--	21 (76)	9.3 (7-15)	n/a	n/a	12 (58)	10.8 (7-15)	n/a	ASD<TD: L/R STS
Bonilha et al. (2008)	Pediat.	--	12 (100)	12.4 (8-15)	n/a	n/a	16 (100)	13.2 (n/a)	n/a	ASD>TD: L/R IFG, L/R cuneus, L/R ACC, R PCC, claustrum, L/R precuneus, L/R STG, L/R MTG, L ITG, thalamus (pulvinar), L/R SFG, L/R SPL, L insula, L putamen, R caudate, L/R FFG, L/R occipital gyrus, L lingual gyrus, L pre-CG, L post-CG, L/R thalamus, L/R PHG, L/R mPFC, R IPL
Brieber et al. (2007)	Pediat.	--	15 (100)	14.2 (10-16)	106.8	0	15 (100)	13.3 (10-16)	107.7	ASD<TD: L/R ITG/hippocampus/ amygdala, L MOG, L PMC, L hippocampus; ASD>TD: R supramarginal gyrus, L post-CG
Cai et al. (2018)	Pediat.	--	38 (84)	9.6 (5-16)	75.8	n/a	27 (96)	8.3 (5-14)	98.6	ASD<TD: R cerebellum anterior lobe, R precuneus, R angular gyrus; ASD>TD: L ITG, L/R MTG
Cheng, Chou, Fan, and Lin (2011)	Pediat.	--	25 (100)	13.7 (10-18)	101.6	n/a	25 (100)	13.5 (11-18)	109.0	ASD<TD: R IFG, R pre-CG, L post-CG, cuneus, thalamus, lingual gyrus, STG; ASD>TD: ACC, paracentral lobule, SPL, precuneus, MFG, FFG, subcallosal gyrus
Contarino, Bulgheroni, Annunziata, Erbetta, and Riva (2016)	Pediat.	--	25 (88)	6.1 (2-12)	56	n/a	25 (65)	6.1 (2-12)	103	--
M. Craig et al. (2007)	Adult	--	14 (0)	37.9 (n/a)	103.4	n/a	19 (0)	35.0 (n/a)	111.2	ASD<TD: Cuneus, L ITG/STG, R MTG, R ACC
D'Mello, Crocetti, Mostofsky, and Stoodley (2015)	Pediat.	--	35 (86)	10.4 (8-13)	n/a	n/a	35 (60)	10.4 (8-13)	n/a	ASD<TD: R lingual gyrus, R cerebellum, R angular gyrus; ASD>TD: L PCC/precuneus, R SFG, L MOG
Ecker et al. (2012)	Adult	--	89 (100)	27.0 (18-43)	110	n/a	89 (100)	28.0 (18-43)	113.0	ASD<TD: R occipital/ITG/MTG, R cerebellum, R posterior FFG, R lingual gyrus, L/R inferior and R SOG, R cuneus, precuneus, R PCC; ASD>TD: L /R ITG/MTG/STG/ L/R FFG/PHG/insula, L IFG, L putamen/caudate, L thalamus, L/R dlPFC/ MFG/pre-/post-CG, R IPL
Foster et al. (2015)	Pediat.	--	38 (100)	12.4 (6-17)	102.5	n/a	46 (100)	12.6 (7-17)	113.1	ASD<TD: R STG, L/R supramarginal gyrus, L cerebellum; ASD>TD: R central sulcus, L medial FG, L/R IFG, L/R pre-CG, L MFG, L pre-SMA, R SFG, L ACC, L OFC, L ITG/STG. L/R MTG, L Heschl's gyrus, R lingual gyrus, L FFG, L post-CG, L PCC, L precuneus, R supramarginal/angular gyrus, L IOG, L/R cuneus, L putamen, L caudate
Freitag et al. (2008)	Mixed	--	15 (87)	17.5 (n/a)	101.2	n/a	15 (87)	18.6 (n/a)	112.1	ASD<TD: R intraparietal sulcus

Greimel et al. (2013)	Mixed	--	47 (100)	18.3 (10-50)	107.5	13	51 (100)	21.4 (8-47)	112.5	ASD<TD: ACC, L/R posterior STS, R MTG
Groen, Buitelaar, van der Gaag, and Zwiers (2011)	Pediat.	--	17 (82)	14.4 (12-18)	98.0	n/a	25 (88)	15.5 (12-18)	105.0	--
Hyde, Samson, Evans, and Mottron (2010)	Mixed	--	15 (100)	22.7 (14-33)	100.4	n/a	15 (100)	19.2 (14-34)	106.6	ASD<TD: R post-CG, L/R pre-CG; ASD>TD: Brainstem/midbrain, reticular, medial FG/OFC, L/R MFG
Itahashi et al. (2015)	Adult	--	46 (100)	30.2 (19-50)	106.0	n/a	46 (100)	30.5 (19-47)	109.2	--
Katz et al. (2016)	Adult	--	23 (100)	26.6 (18-45)	n/a	n/a	32 (100)	29.8 (19-48)	n/a	ASD>TD: L/R OFC, ACC
Kaufmann et al. (2013)	Pediat.	--	10 (80)	14.7 (n/a)	102.3	n/a	10 (80)	13.8 (n/a)	109.5	ASD<TD: Lateral portion of R precuneus; ASD>TD: L medial FG, R precuneus
Ke et al. (2008)	Pediat.	--	17 (82)	8.9 (6-14)	108.8	n/a	15 (80)	9.7 (6-14)	109.8	ASD<TD: R PHG; ASD>TD: L/R supramarginal gyrus, post-CG, R MFG, R cerebellum
Kosaka et al. (2010)	Adult	--	32 (100)	23.8 (17-32)	101.6	n/a	40 (100)	22.5 (18-34)	109.7	ASD<TD: R insula, R IFG, R IPL
Kurth et al. (2011)	Pediat.	--	52 (73)	11.2 (5-20)	102.2	n/a	52 (73)	11.1 (6-19)	106.0	ASD<TD: Hypothalamus
Kwon, Ow, Pedatella, Lotspeich, and Reiss (2004)	Pediat.	--	20 (100)	13.5 (10-18)	n/a	n/a	13 (100)	13.6 (10-18)	n/a	ASD<TD: R ITG, R entorhinal cortex, R FFG
Langen et al. (2009)	Pediat.	--	99 (92)	12.9 (7-24)	107.6	n/a	89 (92)	12.4 (6-24)	110.0	--
Lim et al. (2015)	Pediat.	--	19 (100)	14.9 (11-17)	113.0	0	33 (100)	14.9 (11-17)	110.0	ASD>TD: L MTG/STG, L medial FG
Lin, Ni, Lai, Tseng, and Gau (2015)	Pediat. (Child)	--	28 (100)	10.7 (7-12)	106.9	9	43 (100)	10.6 (7-12)	115.2	ASD<TD: R post-CG, precuneus, L MOG; ASD>TD: L subcallosal gyrus, L/R sublobar areas
	Pediat. (Adolescent)	--	40 (100)	14.7 (13-17)	101.5		18 (100)	15.5 (13-17)	108.7	--
	Adult	--	18 (100)	22.2 (18-29)	99.6		29 (100)	23.4 (18-29)	116.8	ASD>TD: L/R SFG, L MFG
Lin, Tseng, Lai, Chang, and Gau (2017)	Pediat.	--	20 (100)	13.5 (8-19)	103.8	0	54 (100)	12.8 (8-19)	112.5	ASD>TD: L cerebellum crus I
McAlonan et al. (2002)	Adult	--	17 (90)	32.0 (18-49)	96.0	n/a	24 (92)	33.0 (18-49)	114.0	ASD<TD: R cerebellum, L/R lenticular nucleus, R CG, L/R MFG, R SFG, precuneus
McAlonan et al. (2008)	Pediat.	--	33 (82)	11.6 (7-16)	113.2	0	55 (86)	10.7 (7-16)	117.1	ASD<TD: L dlPFC, L/R BG, L/R inferior cerebellar vermis, L STS, posterior parietal cortices
Mengotti et al. (2011)	Pediat.	--	20 (90)	7.0 (4-14)	n/a	0	22 (91)	7.7 (4-11)	n/a	ASD<TD: L SMA, R IFG; ASD>TD: R IPL, L/R ITG, L SPL, R SOG, precuneus
Mueller et al. (2013)	Adult	--	12 (75)	35.5 (n/a)	111.3	n/a	12 (67)	33.3 (n/a)	110.8	ASD<TD: L/R TPJ, L/R temporal lobe, L/R IFG, L MFG, L frontal pole, L/R mPFC, paracingulate gyrus

Ni et al. (2018)	Pediat.	--	81 (100)	12.6 (7-17)	107.4	14.8	61 (100)	12.4 (7-17)	112.0	ASD<TD: L lateral occipital/superior parietal cortex, L frontal pole
Pereira et al. (2018)	Mixed		19 (82)	17.4 (14-25)	99.8	n/a	25 (66)	18.5 (14-25)	105.8	ASD<TD: L/R cerebellum, posterior lobe, R cerebellum anterior lobe, L FFG, L/R CG, L paracentral lobule, R MFG, L claustrum, R medial FG, L PHG L amygdala, L post-CG, R SFG; ASD>TD: L/R brainstem, R cerebellum, posterior lobe
Poulin-Lord et al. (2014)	Mixed	--	23 (87)	19.8 (14-30)	100.3	n/a	22 (86)	22.6 (15-35)	107.3	--
Poustka et al. (2012)	Pediat.	--	18 (89)	9.7 (6-12)	111.0	n/a	18 (89)	9.7 (6-12)	112.8	--
Radeloff et al. (2014)	Mixed	--	34 (91)	19.1 (14-33)	105.7	n/a	26 (85)	19.5 (14-27)	107.8	--
Retico et al. (2016)	Pediat.	--	76 (50)	4.6 (3-7)	71	n/a	76 (50)	4.6 (2-7)	73	ASD<TD: L posterior MTG, L cerebellum, L/R temporal pole; ASD>TD: L precuneus, L PCC, R posterior STG, R MTG/ITG, R mPFC, L SPL, mid-CC
Riedel et al. (2014)	Adult	--	30 (63.3)	35.4 (21-52)	124.5	n/a	30 (63)	35.5 (22-53)	123.6	--
Riva et al. (2013)	Pediat.	--	26 (89)	5.8 (2-10)	51.6	n/a	21 (62)	6.8 (3-10)	n/a	ASD<TD: L/R cerebellum crus II, vermis 8/9, L/R hippocampus, L IFG, L MOG/SOG, L ITG, L SFG. R IOG, R post-CG
Rojas et al. (2006)	Mixed	--	24 (100)	20.8 (7-44)	94.8	n/a	23 (100)	21.4 (7-44)	118.7	--
Sato et al. (2017)	Adult	--	36 (69.4)	27.0 (18-53)	110.4	n/a	36 (69)	24.9 (20-43)	n/a	ASD<TD: R IOG/ITG/MTG, L FFG, L/R amygdala, R PHG, R L/MFG, R SMA, L/R CG, ACC, L/R medial SFG, R OFG
Schmitz et al. (2006)	Adult	--	10 (100)	38.0 (18-52)	105.0	0	12 (100)	39.0 (18-52)	106	ASD>TD: L IFG, ACC, R SFG, L/R MFG
Toal et al. (2010)	Adult	--	65 (88)	31.0 (16-59)	98.0	n/a	33 (91)	32.0 (19-58)	105	ASD<TD: R cerebellum/PHG/ FFG, R ITG/STG L PHG/STG/TG/FFG/ cerebellum
Waiter et al. (2004)	Pediat.	--	16 (100)	15.4 (12-20)	100.4	n/a	16 (100)	15.5 (12-20)	99.7	ASD<TD: R thalamus; ASD>TD: L SFG, R FFG, R mPFC, L MTG, R PCC, L/R STG, L lingual gyrus, L IFG, L MFG, L IOG, L PHG
Wang et al. (2017)	Pediat.	--	31 (100)	4.8 (3-7)	62.5	n/a	31 (100)	4.8 (3-6)	97.1	ASD>TD: L STG, L post-CG
Wilson, Tregellas, Hagerman, Rogers, and Rojas (2009)	Adult	--	10 (80)	30.1 (22-47)	91.5	n/a	10 (70)	29.4 (21-43)	127.2	--
Q. Yang et al. (2018)	Pediat.	--	16 (63)	10.4 (6-15)	45.3	n/a	16 (63)	10.5 (6-15)	97.5	ASD<TD: L cerebellum, L caudate
(B) fMRI studies in ASD										
Ambrosino et al. (2014)	Pediat.	GNG	19 (100)	11.5 (9-12)	112	n/a	19 (100)	11.1 (9-14)	120	--
Chantiluke, Barrett, Giampietro, Santosh, et al. (2015)	Pediat.	Stop	19 (100)	14.7 (10-17)	112	0	25 (100)	13.4 (10-17)	109	ASD<TD: L IPL; ASD>TD: R IFC, L IFG/MFG

Daly et al. (2014)	Adult	GNG	14 (100)	31 (n/a)	115	n/a	14 (100)	31 (n/a)	123	ASD<TD: R IFG, L thalamus; ASD>TD: R caudate, R cerebellum
Denisova et al. (2017)	Mixed	Simon	20 (95)	22.7 (n/a)	n/a	n/a	20 (95)	23.2 (n/a)	n/a	ASD>TD: R lingual gyrus, R cerebellum, R ITG
Duerden et al. (2013)	Adult	GNG	16 (69)	27.2 (19-39)	112	0	17 (71)	30.7 (20-43)	114	ASD<TD: R MFG; ASD>TD: R IFG, R FFG
J. Fan et al. (2012)	Adult	Flanker	12 (75)	30 (n/a)	115	n/a	12 (83)	28 (n/a)	120	ASD<TD: ACC
Gooskens et al. (2018)	Pediat.	Stop	26 (35)	11.3 (8-12)	108.9	n/a	53 (45)	10.8 (8-12)	111.9	--
Kana, Keller, Minshew, and Just (2007)	Adult	GNG	12 (92)	26.8 (n/a)	110	n/a	12 (92)	22.5 (n/a)	117	ASD<TD: L ITG, R PHG, R calcarine sulcus, R PMC, R mid-CC, L/R post-CG, R insula/IFG, L lingual gyrus
Kennedy et al. (2006)	Adult	Stroop	15 (100)	25.5 (15-44)	96	n/a	14 (100)	26.1 (n/a)	n/a	ASD>TD: mPFC, precuneus
Prat, Stocco, Neuhaus, and Kleinhans (2016)	Adult	GNG	16 (63)	25.3 (18-35)	107.4	n/a	17 (65)	25.6 (19-44)	111.0	--
Schmitz et al. (2006)	Adult	GNG, Switch, Stroop	10 (100)	38 (18-52)	105	0	12 (100)	39 (18-52)	106	ASD>TD: L M/IFG, OFC (GNG), insula (Stroop), and IPL, mesial parietal lobe (Switch)
Shafritz, Dichter, Baranek, and Belger (2008)	Adult	Switch	18 (89)	22.3 (n/a)	103	n/a	15 (87)	24.3 (n/a)	111	ASD<TD: L dIPFC, ACC, L intraparietal sulcus, L BG, L insula
Shafritz, Bregman, Ikuta, and Szeszko (2015)	Mixed	GNG	15 (80)	18.1 (13-23)	102	0	15 (80)	18.4 (12-23)	115	ASD<TD: R IFG/insula
Solomon et al. (2014)	Pediat.	Switch	27 (19)	15.4 (12-18)	108	55	27 (19)	16.1 (12-18)	113	ASD<TD: L PCC, L lingual gyrus, L MOG
Vaidya et al. (2011)	Pediat.	Stroop	11 (100)	10.8 (7-12)	114	n/a	14 (100)	11.0 (n/a)	119	ASD<TD: ACC, L MFG, R caudate
van Hulst et al. (2017)	Pediat.	GNG	26 (100)	10.8 (8-12)	109.6	n/a	26 (100)	10.5 (8-12)	117.3	--
Velasquez et al. (2017)	Adult	GNG	19 (68)	25.8 (18-35)	111	0	22 (73)	29.0 (20-46)	112	ASD<TD: R ACC, post-CG, R lateral occipital cortex
Yerys et al. (2015)	Pediat.	Switch	20 (80)	11.3 (7-14)	115	n/a	19 (68)	11.4 (7-13)	120	ASD>TD: L MFG/ pre-CG, L SFG/ dACC, R IFG

Abbreviations. N=sample size, y=year, pediat.=pediatric (child/adolescent) sample, com ADHD=comorbid attention-deficit/hyperactivity disorder based on explicit reporting or study exclusion criteria, ASD=autism spectrum disorder, TD=typically developing controls, L/R=Left/Right, GNG = Go/No-Go. Brain region (in alphabetical order): ACC=anterior cingulate cortex, BG=basal ganglia, dACC=dorsal ACC, dIPFC=dorsolateral prefrontal cortex, FFG=fusiform gyrus, I/M/SFG=inferior/middle/medial/superior frontal gyrus, I/M/SOG=inferior/middle/superior occipital gyrus, I/M/STG=inferior/middle/superior temporal gyrus, I/SPL=inferior/superior parietal lobe, mPFC=medial prefrontal cortex, OFC=orbital frontal cortex, OFG=orbital frontal gyrus, PCC=posterior cingulate cortex, pre-/post-CG=pre-/post-central gyrus, PHG=parahippocampal gyrus, PMC=premotor cortex, SMA=supplementary motor area, STS=superior temporal sulcus

Supplement 3: Meta-analyses in Pediatric Sample with ADHD and ASD

Table S3a. Characteristics of Overall Pediatric Sample and Age-, Sex- and IQ-matched Subgroups

	ADHD	ASD	TD _{ADHD}	TD _{ASD}
(A) Overall Sample				
VBM				
N	615	848	568	869
% males	81	87	82	86
Mean age (range), y	11.6 (6-20)	10.8 (2-24 ^a)	12.0 (6-20)	10.8 (2-24 ^a)
Mean FSIQ (SD)	102 (9.7)	94 (18.4)	110 (5.0)	106 (11.1)
fMRI during Cognitive Control				
N	568	--	572	--
% males	79	--	73	--
Mean age (range), y	13.0 (7-20)	--	13.1 (7-20)	--
Mean FSIQ (SD)	101 (6.7)	--	109 (5.5)	--
fMRI during Motor Response Inhibition				
N	389	--	395	--
% males	77	--	72	--
Mean age (range), y	13.1 (7-20)	--	13.0 (7-20)	--
Mean FSIQ (SD)	99 (7.1)	--	109 (5.5)	--
(B) Matched Subgroups				
VBM				
N	596	741	550	765
% males	82	86	82	84
Mean age (range), y	11.7 (6-20)	11.0 (2-24 ^a)	12.0 (6-20)	11.0 (2-24 ^a)
Mean FSIQ (SD)	101 (2.4)	96 (4.7)	109 (1.1)	106 (4.1)

Abbreviations: N = overall number of subjects, TD_{ASD} = typically developing controls in the ASD studies, TD_{ADHD} = typically developing controls in the ADHD studies, VBM = voxel-based morphometry, fMRI = functional Magnetic Resonance Imaging, % males = proportion of males among the samples, y = year, FSIQ = full scale IQ, and SD = standard deviation. The demographic characteristics presented here were calculated from the independent datasets with non-overlapping participants. Due to insufficient number of independent datasets, no fMRI meta-analysis in pediatric groups were carried out and no demographic information is reported here. ^aincluded one study primarily in adolescents with upper age limit of 24 years.

Table S3b. Structural and Functional Abnormalities in Pediatric Sample with ADHD and ASD

Contrasts	MNI coord. x, y, z	SDM Z	p-value	Voxels No	BA
(A) Structural Abnormalities					
(i) ADHD vs. TD					
ADHD < TD					
R STG/putamen/posterior insula*	56,-14,8	2.47	.0003	1709	48/22/38
rACC/vmPFC/vmOFC*	6,50,8	2.61	.0001	1223	32/10/11
R caudate*	18,12,10	2.44	.0003	353	--
L cerebellum hemispheric lobule VI	-28,-48,-34	2.08	.002	155	--
R cerebellum hemispheric lobule IX	16,-52,-46	1.93	.004	32	--
ADHD > TD					
Nil	--	--	--	--	--
(ii) ASD vs. TD					
ASD < TD					
L cerebellum lobule VIII*	-8,-66,-46	1.58	.0002	890	--
L MOG/SPG	-26,-82,40	1.59	.0002	415	19/7
Dorsomedial thalamus*	4,-4,16	1.48	.0003	144	--
Middle cingulate gyrus	0,-6,28	1.18	.002	23	--
ASD > TD					
L anterior temporal pole/STG/MTG*	-46,-10,-6	2.40	.00005	1051	21/48/22/20/38
Precuneus/PCC*	-4,-55,22	2.40	.00005	651	23/30/26
R MFG/SFG/dIPFC*	26,46,28	2.42	.00003	236	46/9
ADHD (vs. TD) vs. ASD (vs. TD)					
(iii) Overall Sample					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
R STG/MTG/Al	58,-12,0	2.29	.0006	938	22/21/48/38
mPFC/rACC/vmOFC*	6,58,10	2.46	.0003	906	10/32/11
L post-CG	-54,-12,34	2.02	.002	59	3/4
ASD (vs. TD) reduced vs. ADHD (vs. TD)					
L MOG	-24,-82,40	1.22	.0001	820	19
ASD (vs. TD) increased vs. ADHD (vs. TD)					
L posterior insula/STG/MTG*	-46,-8,-2	2.29	.0006	446	48/38
R MFG/dIPFC*	24,46,24	2.37	.0004	110	46
(iv) Matched Subgroups					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
R putamen/Al/STG	58,-12,2	2.09	.0006	1369	22/48
rACC/vmPFC/vmOFC*	6,60,12	2.36	.0001	1291	10/11/32
R caudate	14,14,8	1.87	.002	67	--
(B) Functional Abnormalities					
ADHD vs. TD					
(i) Cognitive Control					
ADHD < TD					
SMA/dmPFC*	4,6,50	1.72	.0001	554	6/32/24
L anterior MTG*	-50,-18,-10	1.63	.0003	197	48/22/20/21
L middle cerebellar peduncle	-14,-62,-36	1.26	.002	154	--
L post-CG*	-48,-10,56	1.23	.002	22	6
ADHD>TD					
L medial SFG	-8, 28, 48	1.09	.002	45	8/9/32

(ii) Motor Response Inhibition

ADHD < TD


R vIPFC/OFC/AI*	38,28,-16	1.74	.0005	379	47/38
L anterior MTG*	-52,-18,-10	2.06	.0001	323	22/21/10/48
vmOFC	-4,56,-6	1.43	.002	326	10
L lingual gyrus	-10,-32,-8	1.35	.003	47	27
L post-CG*	-48,-10,56	1.48	.002	27	6

ADHD > TD

L medial SFG	-8,28,50	1.29	.002	39	8
--------------	----------	------	------	----	---

Abbreviations: MNI = Montreal Neurological Institute, SDM = seed-based d mapping, BA = Brodmann Area, TD = typically developing controls, L/R = left/right. Brain regions (in alphabetical order): AI = anterior insula, dIPFC = dorsolateral prefrontal cortex, dmPFC = dorsomedial prefrontal cortex, mPFC = medial prefrontal cortex, MOG = middle occipital gyrus, M/SFG = middle/superior frontal gyrus, M/STG = middle/superior temporal gyrus, OFC = orbitofrontal cortex, PCC = posterior cingulate cortex, PHG = parahippocampal gyrus, pre-/post-CG = pre-/post-central gyrus, rACC = rostral anterior cingulate cortex, SMA = supplementary motor area, SPG = superior parietal gyrus, vIPFC = ventrolateral prefrontal cortex, vmOFC = ventromedial orbitofrontal cortex, vmPFC = ventromedial prefrontal cortex. Asterisks indicate overlapping impairments with findings in the age non-stratified overall samples. Bold fonts indicate overlapping disorder-differentiating impairment between the overall pediatric samples and age-, sex- and IQ-matched pediatric subgroups. No pediatric comparative fMRI meta-analyses involving the ASD population were carried out due to insufficient number of ASD independent datasets.

Figure S3. Structural and Functional Abnormalities in Pediatric Sample with ADHD and ASD



Abnormalities in the (A) gray matter volume (GMV) and (B) brain activations during (i) cognitive control and (ii) motor response inhibition only in ADHD relative to typically developing (TD) controls. A statistical threshold of $p < .005$ and a cluster extent of 20 voxels were used.

Supplement 4: Meta-analyses in Adults with ADHD and ASD

Table S4a. Characteristics of Adult Overall Sample and Subgroups Matched on Age, Sex and IQ

	ADHD	ASD	TD _{ADHD}	TD _{ASD}
(A) Overall Sample				
VBM				
<i>N</i>	611	417	531	427
% males	57	87	58	87
Mean age (range), y	32.0 (18-65)	29.1 (16 ^a -59)	32.4 (18-65)	27.7 (18-58)
Mean FSIQ (SD)	108 (5.6)	104 (11.5)	112 (5.7)	113 (6.0)
fMRI during Cognitive Control				
<i>N</i>	433	152	432	135
% males	75	85	65	84
Mean age (range), y	25.4 (17-50)	28.7 (15 ^b -52)	25.8 (18-50)	28.4 (18-52)
Mean FSIQ (SD)	106 (6.6)	108 (5.8)	112 (4.6)	114 (5.3)
fMRI during Motor Response Inhibition				
<i>N</i>	346	--	339	--
% males	77	--	65	--
Mean age (range), y	25.4 (18-50)	--	26.0 (18-50)	--
Mean FSIQ (SD)	103 (5.3)	--	111 (5.2)	--
(B) Matched Subgroups				
VBM				
<i>N</i>	287	232	275	223
% males	71	77	70	75
Mean age (range), y	27.7 (18-65)	31.4 (16 ^a -59)	28.4 (18-65)	28.5 (18-58)
Mean FSIQ (SD)	103 (5.7)	102 (13.6)	108 (4.5)	114 (8.7)
fMRI during Overall Cognitive Control				
<i>N</i>	257	152	241	135
% males	81	85	80	84
Mean age (range), y	26.5 (17-50)	28.7 (15 ^b -52)	26.0 (18-50)	28.4 (18-52)
Mean FSIQ (SD)	109 (6.8)	108 (5.8)	114 (3.8)	114 (5.3)

Abbreviations: *N* = overall number of subjects, TD_{ASD} = controls in the ASD studies, TD_{ADHD} = controls in the ADHD studies, VBM = voxel-based morphometry, fMRI = functional Magnetic Resonance Imaging, % males = proportion of males among the samples, y = year, FSIQ = full scale IQ, and SD = standard deviation. The demographic characteristics presented here were calculated from the independent datasets with non-overlapping participants. There were insufficient number of adult ASD fMRI datasets during motor control thus no meta-analyses involving adult ASD participants were conducted during prepotent response inhibition and the participant characteristics are not reported in this table. ^a included one study primarily in adults with lower age limit of 16 years. ^b included a study conducted primarily in adult with 3/12 participants aged 15-16 years old.

Table S4b. Structural and Functional Abnormalities in Adults with ADHD and ASD


Contrasts	MNI coord. x, y, z	SDM Z	p-value	Voxels No	BA
(A) Structural Abnormalities					
(i) ADHD vs. TD					
ADHD < TD					
vmOFC/subcallosal gyrus*	0,56,-24	1.81	.00005	952	11/25
R supramarginal gyrus/posterior MTG/STG	54,-42,24	1.45	.0005	680	48/42/40/22/21/41
R IFG/MFG	46,16,40	1.11	.003	53	44/9
R SMA	8,26,64	1.03	.004	27	8
ADHD>TD					
Nil	--	--	--	--	--
(ii) ASD vs. TD					
ASD < TD					
R putamen/posterior insula	38, 6, -2	1.86	.001	920	48
R cerebellum crus I	26, -78, -24	1.80	.001	174	19/18
L/R cuneus	6, -70, 12	1.85	.001	67	17/18
R lingual gyrus	18,-96,-10	1.84	.001	45	18
L MOG	-36,-94,-10	1.73	.002	30	18
ASD > TD					
L anterior MTG/ITG*	-60,-22,-16	2.15	.00002	266	21/20/22
L pre-/post-CG	-38,-16, 50	1.75	.0002	212	6/4
L SFG/MFG/dIPFC	-20, 28, 44	1.84	.0001	117	9/8/32
L posterior STG	-40, -22, -2	1.61	.0007	111	48
R pre-/post-CG	46, -18, 50	1.50	.002	98	6/4
L MTG	-44, 4, -26	1.68	.0005	37	20/21
ADHD (vs. TD) vs. ASD (vs. TD)					
(iii) Overall Sample					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
vmOFC/subcallosal gyrus*	-4,56,-22	1.87	.00004	672	11
vmOFC/gyrus rectus*	-10,16,-12	1.29	.002	65	11
R supramarginal gyrus	58,-36,38	1.42	.0007	269	40/48
ASD (vs. TD) reduced vs. ADHD (vs. TD)					
R cerebellum, crus I	26,-78,-24	1.37	.002	261	--
R insula	44,2,2	1.34	.002	104	--
R occipital pole	6,-72,12	1.47	.001	60	18
R lingual gyrus	18,-94,-10	1.59	.0006	50	18
L occipital pole	-36,-94,-10	1.32	.002	27	18
ASD (vs. TD) increased vs. ADHD (vs. TD)					
L MTG*	-60,-22,-16	1.56	.0003	154	21/20
L pre-CG	-38,-16,50	1.33	.001	90	6
L dIPFC/MFG/SFG	-20,30,46	1.45	.0006	62	9/8
(iv) Matched Subgroups					
ADHD (vs. TD) reduced vs. ASD (vs. TD)					
vmOFC/subcallosal gyrus*	0,48,-28	1.39	.0006	604	11
vmOFC/gyrus rectus*	6,14,-18	1.15	.002	87	11
R IFG/MFG	46,18,40	1.25	.001	59	44
ASD (vs. TD) reduced vs. ADHD (vs. TD)					
R amygdala/PHG/FFG	30,-2,-22	1.95	.0003	375	36/34/28
L FFG	-28,-32,-26	1.49	.003	84	20/30/27

ASD (vs. TD) increased vs. ADHD (vs. TD)						
R posterior STG*	56,-42,22	1.24	.001	180	42/48/41	
(B) Functional Abnormalities during Cognitive Control						
(i) ADHD vs. TD						
ADHD < TD						
L MFG/dIPFC*	-26,34,40	1.71	.0002	361	9/46/8	
R caudate/dorsomedial thalamus*	6,-6,12	1.76	.0001	364	--	
R AI/putamen*	30,24,2	1.40	.002	274	47/48	
L temporal pole/STG/insula*	-48,8,-14	1.43	.001	270	38/21/48	
L AI/IFG*	-36,20,6	1.37	.002	86	48/45	
R temporal pole/MTG/STG*	52,4,-18	1.31	.003	74	21/38	
R MTG/STG*	52,-12,-12	1.26	.003	41	22/20	
R posterior STG	56,-40,18	1.30	.003	29	42	
ADHD>TD						
Nil	--	--	--	--	--	
(ii) ASD vs. TD						
ASD < TD						
rdACC/dmPFC*	-2,36,22	1.99	.00001	2885	32/24	
L IPL/SPL*	-34,-46,52	1.43	.001	321	40/2/7	
ASD > TD						
Precuneus*	6,-48,60	1.77	.00004	1181	5/4	
L vIPFC/OFC*	-32,44,-14	1.35	.001	103	11/47	
L MFG/dIPFC*	-26,48,14	1.35	.001	57	46/10	
ADHD (vs. TD) vs. ASD (vs. TD)						
(iii) Overall Sample						
ASD (vs. TD) reduced vs. ADHD (vs. TD)						
mPFC/rACC/vmPFC/vmOFC*	0,36,20	2.70	.0000008	2948	32/24//9/10	
L IPL/SPL/post-CG	-34,-42,54	1.29	.001	108	40/2/3	
ASD (vs. TD) increased vs. ADHD (vs. TD)						
Precuneus/PCC*	-4,-40,52	2.35	.00001	1462	4/5/7	
L MFG/dIPFC	-28,48,8	1.62	.001	69	46/10	
L vIPFC/OFC*	-30,42,-8	1.55	.001	60	11/47	
(iv) Matched Subgroups						
ASD (vs. TD) reduced vs. ADHD (vs. TD)						
dmPFC/ACC/vmPFC/vmOFC*	0,34,20	2.63	.0000005	3241	32/24/9/10/8/11	
L IPL/SPL/post-CG*	-34,-42,54	1.22	.002	71	40/2/3	
ASD (vs. TD) increased vs. ADHD (vs. TD)						
Precuneus/PCC*	-4,-40,54	2.33	.000008	1548	4/5/7	
L vIPFC/OFC*	-30,42,-8	1.47	.002	33	11/47	
L MFG/dIPFC	-28,48,8	1.48	.002	30	46/10	
(C) Functional Abnormalities during Motor Response Inhibition						
(i) ADHD vs. TD						
ADHD < TD						
L MFG/dIPFC*	-28,28,48	1.81	.00006	554	9/46/8	
R AI/putamen/IFG*	32,22,4	1.70	.0002	456	47/48/45	
R MTG/STG	46,-20,-8	1.33	.002	255	48/22/21/20	
R posterior STG/supramarginal gyrus	54,-38,16	1.42	.001	165	42/48/41	
R IPL	48,-36,50	1.33	.002	69	2/40	
L Precuneus	-12,-32,58	1.39	.001	41	7	
ADHD > TD						

Nil	--	--	--	--	--
-----	----	----	----	----	----

Abbreviations: MNI=Montreal Neurological Institute, SDM=seed-based d mapping, BA=Brodmann Area, TD=typically developing controls, L/R = left/right. Brain regions (in alphabetical order): AI=anterior insula, dlPFC = dorsolateral prefrontal cortex, dmPFC = dorsomedial prefrontal cortex, FFG = fusiform gyrus, I/M/SFG = inferior/middle/superior frontal gyrus, I/M/STG=inferior/middle/superior temporal gyrus, I/SPL= inferior/superior parietal lobe, MOG = middle occipital gyrus, OFC = orbitofrontal cortex, PCC = posterior cingulate cortex, PHG = parahippocampal gyrus, pre-/post-CG = pre/postcentral gyrus, rACC = rostral anterior cingulate cortex, rdACC = rostr dorsolateral anterior cingulate cortex, SMA = supplemental motor area, vlPFC = ventrolateral prefrontal cortex, vmOFC = ventromedial orbitofrontal cortex, vmPFC = ventromedial prefrontal cortex. Asterisks indicate overlapping impairments with findings in the age non-stratified overall samples. Bold fonts indicate overlapping disorder-differentiating impairment between the adult overall samples and age-, sex- and IQ-matched adult subgroups. No adult fMRI meta-analyses during prepotent response inhibition involving the ASD population were carried out due to insufficient number of adult ASD independent datasets.

Figure S4. Structural and Functional Abnormalities in Adults with ADHD and ASD



Abnormalities in the (A) gray matter volume and brain activations during (B) cognitive control and (C) motor response inhibition. Rows (i) and (ii) show abnormalities in ADHD and ASD relative to typically developing (TD) controls. Abnormalities in ADHD versus ASD, each relative to TD controls are shown in (iii) overall adult samples and (iv) age-, sex-, and IQ-matched subgroups. No adult fMRI meta-analyses during motor control involving the ASD population were carried out due to insufficient number of adult ASD independent datasets. A statistical threshold of $p < .005$ with a cluster extent of 20 voxels were used in all analyses.

Supplement 5: Associations between Brain Abnormalities and Age

Correlational Analyses between Brain Abnormalities and Age

Among the most consistent disorder-differentiating VBM abnormalities, the ADHD-differentiating GMV reduction relative to ASD in medial orbitofrontal/anterior cingulate cortices was positively correlated with age (Spearman's $\rho = .18$, $p < .0005$), while the ASD-differentiating enhanced GMV relative to ADHD in right posterior temporal ($\rho = -.30$, $p < .0005$), right dorsolateral prefrontal ($\rho = -.43$, $p < .0005$) and left anterior posterior cortices ($\rho = -.29$, $p < .0005$) was negatively correlated with age (Table S4-i).

Among the most consistent disorder-differentiating fMRI abnormalities during cognitive control, the ASD-differentiating underactivation in left middle frontal/dorsolateral prefrontal cortex ($\rho = .24$, $p < .0005$), and the overactivation in precuneus ($\rho = .36$, $p < .0005$) and in left ventrolateral prefrontal cortex ($\rho = .50$, $p < .0005$) were all positively correlated with age. However, the ASD-differentiating underactivation in anterior cingulate/dorsomedial prefrontal and the overactivation in right fusiform gyrus/inferior occipital and in inferior frontal cortices were not correlated with age (Table S4-ii).

Finally, among the most consistent disorder-differentiating fMRI abnormalities during motor response control, the ADHD-differentiating underactivation in right caudate ($\rho = .44$, $p < .0005$) and inferior frontal gyrus ($\rho = .20$, $p = .002$) both were positively correlated with age. The ASD-differentiating underactivation in precuneus ($\rho = -.21$, $p = .02$) was negatively correlated with age, while the overactivation in left ventrolateral prefrontal/orbitofrontal ($\rho = .61$, $p < .0005$) and middle frontal/dorsolateral prefrontal cortices ($\rho = .61$, $p < .0005$) both were positively correlated with age. The ASD-differentiating underactivation in left cerebellum and right dorsolateral prefrontal cortex, and the ASD-differentiating overactivation in right inferior occipital gyrus/fusiform gyrus and the shared underactivation in ASD and ADHD in right AI were not correlated with age (Table S4-iii).

Table S5. Spearman's Correlations between Disorder-Differentiating or Shared Abnormalities and Age

	Peak MNI coord. x, y, z	ρ	p-value (Bonferroni adjusted)
(i) VBM Abnormalities			
ADHD (vs. TD) reduced vs. ASD (vs. TD)			
vmOFC/rACC	2, 48, -18	.18	<.0005
ASD (vs. TD) increased vs. ADHD (vs. TD)			
R MTG/STG/angular gyrus	50, -40, 8	-.30	<.0005
R MFG/SFG/dIPFC	24, 44, 24	-.43	<.0005
L anterior MTG/STG	-50, -20, -8	-.29	<.0005
(ii) Cognitive-Control fMRI Abnormalities			
ASD (vs. TD) reduced vs. ADHD (vs. TD)			
rACC/dmPFC	2, 28, 20	-.09	.54
L MFG/dIPFC	-40, 34, 28	.24	<.0005
ASD (vs. TD) increased vs. ADHD (vs. TD)			
Precuneus/PCC	-2, -40, 50	.36	<.0005
R FFG/IOG/ITG	36, -70, -8	.07	>.99
R IFG	44, 24, 18	-.13	.11
L vIPFC/OFC	-30, 40, -8	.50	<.0005
(iii) Motor-Response-Inhibition fMRI Abnormalities			
ADHD (vs. TD) reduced vs. ASD (vs. TD)			
R caudate	8, 20, 2	.44	<.0005
R IFG	42, 26, 20	.20	.002
ASD (vs. TD) reduced vs. ADHD (vs. TD)			
L cerebellum lobule IV/FFG/lingual gyrus	-16, -48, -12	.02	.81
R MFG/dIPFC	42, 14, 48	-.06	>.99
R precuneus	16, -36, 44	-.21	.018
ASD (vs. TD) increased vs. ADHD (vs. TD)			
R IOG/FFG	36, -70, -10	.06	>.99
L vIPFC/OFC	-28, 36, -8	.61	<.0005
L MFG/SFG/dIPFC	-28, 54, 8	.61	<.0005
Reduced in ASD (vs. TD) and ADHD (vs. TD)			
R AI	40, 20, -6	-.08	.54

Abbreviations: MNI = Montreal Neurological Institute, L/R=left/right, SDM = Seed-based d mapping, BA = Brodmann Area, TD = typically developing controls, r = Pearson correlation coefficient, brain regions (in alphabetical order): AI = anterior insula, dIPFC = dorsolateral prefrontal cortex, dmPFC = dorsomedial prefrontal cortex, FFG = fusiform gyrus, IFG M/SFG = inferior/middle/superior frontal gyrus, IOG = inferior occipital gyrus, I/M/STG = inferior/middle/superior temporal gyrus, OFC=orbitofrontal cortex, PCC = posterior cingulate cortex, rACC = rostral anterior cingulate cortex, vIPFC = ventrolateral prefrontal cortex, vmOFC = ventromedial orbitofrontal cortex.

Additional Analyses Covarying for Age in the Sub-Meta-Analyses with Participants Matched on Age, Sex and IQ


All disorder-differentiating findings in the VBM analysis, including the ADHD-differentiating reduced GMV in vmOFC/rACC (BA11; MNI coordinates: 2, 48, -18; $Z = 2.04$, $p = .002$, 193 voxels) remained after covarying for age, as did the ASD-differentiating increased GMV in right MTG/STG/angular gyrus (BA42/21/22/42; MNI coordinates: 50, -40, 8; $Z = 2.46$, $p = .0003$, 384 voxels), right MFG/SFG/dIPFC (BA46; MNI coordinates: 24, 44, 24; $Z = 2.11$, $p = .001$, 68 voxels) and left anterior MTG/STG (BA48/22/21/20; MNI coordinates: -50, -20, -8; $Z = 1.98$, $p = .002$, 58 voxels).

During cognitive control, the ASD-differentiating underactivation in rACC/dmPFC (BA32/24/9/10; MNI coordinates: 0, 30, 20; $Z = 2.27$, $p = .0000002$, 2809 voxels) and left MFG/dIPFC (BA46/45; MNI coordinates: -40,34,28; $Z = 1.02$, $p = .001$, 196 voxels), and all ASD-differentiating overactivation in precuneus/PCC (BA4/5; MNI coordinates: -2,-40,50; $Z = 1.18$, $p = .0009$, 348 voxels), right FFG/IOG/ITG (BA19/37; MNI coordinates: 36,-70,-8; $Z = 1.50$, $p = .00007$, 283 voxels), right IFG (BA48/45; MNI coordinates: 44,24,18; $Z = 1.25$, $p = .0005$, 112 voxels), and left vIPFC/OFC (BA11; MNI coordinates: -30, 40, -8; $Z = 1.06$, $p = .002$, 56 voxels) remained after covarying for age.

Specifically during motor response inhibition, with the exception of the shared right AI underactivation in both disorders, all other findings, i.e., the ADHD-differentiating underactivation in right caudate (BA25; MNI coordinates: 8,20,0; $Z = 1.35$, $p = .0008$, 188 voxels) and right inferior frontal gyrus (BA48/45; MNI coordinates: 42, 26, 20; $Z = 1.32$, $p = .0009$, 107 voxels); and the ASD-differentiating underactivation in left cerebellum/FFG/lingual gyrus (BA19/37/30; MNI coordinates: -16,-48,-12; $Z = 1.37$, $p = .0003$, 1045 voxels), right MFG/dIPFC (BA9; MNI coordinates: 42,14,48; $Z = 1.18$, $p = .001$, 77 voxels) and right precuneus (BA7; MNI coordinates: 16,-36,44; $Z = -1.27$, $p = .0006$, 49 voxels), and the ASD-differentiating overactivation in right IOG/FFG (BA19/37; MNI coordinates: 36,-72,-10; $Z = 1.80$, $p = .00004$, 461 voxels), and left vIPFC/OFC/MFG/SFG/dIPFC (BA11/10; MNI coordinates: -26,36,-8; $Z = 1.41$, $p = .0005$, 593 voxels) remained after covarying for age.


Supplement 6: Associations between Brain Abnormalities and Psychostimulant Exposure

Figure S6a. Association between VBM Abnormalities and Current Psychostimulant Exposure in ADHD



Current (i) and lifetime (ii) psychostimulant exposure were significantly associated with increased GMV in vmOFC (BA11; associations between SDM Z score and current and lifetime exposure peaked at MNI coordinates: -2, 52, -24; $Z = 1.99$, $p = .000002$, 362 voxels and coordinates: 0, 54, -26; $Z = 1.59$, $p = .0002$, 22 voxels, respectively).

Figure S6b. Association between fMRI Abnormalities and Lifetime Psychostimulant Exposure in ADHD



During overall cognitive control, lifetime psychostimulant exposure was positively associated with activation in left IFG (BA48; MNI coordinates: -46, 16, 4; $Z = 1.42$, $p = .0002$, 56 voxels). Current stimulant exposure was not significantly associated with brain functional activations in ADHD.

Supplement 7: Influences of Task Type or Performance in Disorder-differentiating Findings

Table S7. Task Types and Behavioural Performance Impairments in ADHD and ASD vs. TD

fMRI Studies	Task	Behavioural performance	Clinical group impaired?	Subgroup MA
ADHD Studies				
Banich et al. (2009)	Stroop	ADHD > TD (RT incongruent vs. congruent)	No	Yes
Bhaijiwala et al. (2014)	Stop	ADHD < TD (SSRT)	Yes	Yes
Booth et al. (2005)	GNG	ADHD < TD (Error rates, MRT)	Yes	Yes
Carmona et al. (2012)	GNG	ADHD < TD (MRT)	Yes	Yes
Chantiluke et al. (2015)	Stop	ADHD < TD (Omissions)	Yes	Yes
Chen et al. (2015)	GNG	ADHD = TD (Correct response, MRT)	No	Yes
Chou et al. (2015)	Stroop	ADHD < TD (Correct response, MRT)	Yes	Yes
Congdon et al. (2014)	Stop	ADHD = TD (SSRT, MRT, SDRT)	No	--
Cubillo et al. (2010)* Cubillo et al. (2011)*	Stop, Switch, Stroop	ADHD = TD (Stop: SSRT; switch: error rates and MRT; Stroop: error rates, MRT incongruent vs. congruent)	No	Yes
Cubillo et al. (2014)	Stop	ADHD = TD (SSRT, MRT, SDRT)	No	Yes
Dibbets et al. (2009)** Dibbets et al. (2010)**	GNG, Switch	ADHD < TD (Error rates) ADHD < TD (Omissions)	Yes	Yes
Durston et al. (2006)	GNG	ADHD = TD (Accuracy, MRT)	No	Yes
Fan et al. (2017)	Stroop	ADHD < TD (RT incongruent vs. congruent)	Yes	Yes
Fan et al. (2018)	Stroop	ADHD < TD (RT incongruent vs. congruent)	Yes	Yes
Hwang et al. (2015)	Stroop	ADHD < TD (MRT)	Yes	Yes
Iannaccone et al. (2015)	GNG/ Flanker	ADHD < TD (Accuracy)	Yes	Yes
Janssen et al. (2015)	Stop	ADHD < TD (SSRT, omissions)	Yes	--
Konrad et al. (2006)	Flanker	ADHD < TD (RT incongruent vs. congruent)	Yes	Yes
Kooistra et al. (2010)	GNG	ADHD < TD (MRT, SDRT)	Yes	Yes
Ma et al. (2012)	GNG	ADHD = TD (Commissions, omissions, MRT)	No	--
Ma et al. (2016)	Stroop	ADHD = TD (RT, error rates incongruent vs. congruent)	No	--
Massatt et al. (2018)	Stop	ADHD = TD (SSRT, MRT)	No	Yes
Passarotti et al. (2010)	Stop	ADHD < TD (Accuracy)	Yes	Yes
Peterson et al. (2009)	Stroop	ADHD = TD (RT incongruent vs. congruent)	No	Yes
Rasmussen et al. (2016)	GNG	ADHD < TD (Commissions)	Yes	--
Rubia et al. (2005)	Stop	ADHD < TD (SDRT, omissions)	Yes	Yes
Rubia et al. (2011)*** Rubia et al. (2011)***	Simon, Stop	ADHD < TD (MRT) ADHD < TD (SDRT)	Yes	Yes
Schulz et al. (2004)	GNG	ADHD < TD (Commissions)	Yes	Yes
Schulz et al. (2014)	GNG	ADHD < TD (Commissions, accuracy)	Yes	Yes
Schulz et al. (2017)	Stroop	ADHD = TD (MRT, error rates, omissions)	No	Yes
Sebastian et al. (2012)	GNG Stroop Stop	ADHD < TD (Stroop: interference effect; GNG: omissions; Stop: SSRT, SDRT)	Yes	Yes
Shang et al. (2018)	Stroop	ADHD = TD (MRT, SDRT)	No	Yes
Siniatchkin et al. (2012)	GNG	ADHD < TD (MRT)	Yes	Yes

Smith et al. (2006)	GNG, Stroop, Switch	ADHD < TD (GNG: MRT, SDRT; Switch: MRT; Stroop: SDRT)	Yes	Yes
Spinelli et al. (2011)	GNG	ADHD < TD (Omissions)	Yes	Yes
Szekely et al. (2018)	Stop	ADHD = TD (SSRT, MRT, accuracy)	No	--
Tamm et al. (2004)	GNG	ADHD < TD (Omissions, commissions)	Yes	Yes
Thornton et al. (2018)	GNG	ADHD < TD (Commissions)	Yes	--
Van Hulst et al. (2017)	GNG	ADHD < TD (Accuracy, SDRT, MRT)	Yes	Yes
Van Rooij et al. (2015) – adult	Stop	ADHD < TD (SSRT, errors, SDRT)	Yes	--
Van Rooij et al. (2015) – pediatric	Stop	ADHD < TD (SSRT, errors, SDRT)	Yes	--
Zamorano et al. (2017)	Stroop	ADHD > TD (RT incongruent vs. congruent)	No	Yes
ASD studies				
Ambrosino et al. (2014)	GNG	ASD = TD (Accuracy, MRT)	No	Yes
Chantiluke et al. (2015)	Stop	ASD = TD (SSRT, omissions, MRT)	No	Yes
Daly et al. (2014)	GNG	ASD = TD (MRT, probability of inhibition)	No	Yes
Denisova et al. (2017)	Stroop	ASD = TD (RT incongruent vs. congruent)	No	Yes
Duerden et al. (2013)	GNG	ASD = TD (Errors, MRT)	No	Yes
Fan et al. (2012)	Flanker	ASD < TD (Conflict errors, MRT)	Yes	Yes
Gooskens et al. (2018)	Stop	ASD = TD (SSRT, errors, MRT)	No	Yes
Kana et al. (2007)	GNG	ASD = TD (Commissions, MRT)	No	Yes
Kennedy et al. (2006)	Stroop	ASD = TD (Accuracy, MRT)	No	Yes
Prat et al. (2016)	GNG	ASD = TD (Accuracy, MRT)	No	Yes
Schmitz et al. (2006)	GNG, Switch, Stroop	ASD = TD (GNG: accuracy, MRT; Switch & Stroop: errors, RT incongruent vs. congruent)	No	Yes
Shafritz et al. (2008)	Switch	ASD < TD (Accuracy, RT)	Yes	Yes
Shafritz et al. (2015)	GNG	ASD = TD (Errors, MRT)	No	Yes
Solomon et al. (2014)	Switch	ASD < TD (Errors)	Yes	Yes
Vaidya et al. (2011)	Stroop	ASD < TD (RT incongruent vs. congruent)	Yes	Yes
Van Hulst et al. (2017)	GNG	ASD < TD (Accuracy, MRT, SDRT)	Yes	Yes
Velasquez et al (2017)	GNG	ASD = TD (Accuracy, MRT)	No	Yes
Yerys et al. (2015)	Switch	ASD < TD (Accuracy)	Yes	Yes

From the ADHD studies, 26/41 independent datasets demonstrated poorer performance in the patient group relative to controls. From the ASD studies, 6/18 independent datasets demonstrated poorer performance in the patient group relative to controls. Abbreviations: TD = typically developing controls, GNG = Go/No-Go, omissions = omission errors, commissions = commission errors, RT = response time, MRT = mean RT, SDRT = standard deviation of RT, SSRT = stop-signal reaction time. *, **, *** Datasets with largely overlapping participants, combined to control shared variance.

Comparing Task Type and Performance in ADHD and ASD vs. TD

A trend for an interaction between task type (motor response inhibition, cognitive interference, switch, combination of tasks) and diagnostic groups was observed overall, $\chi^2(3, N = 60) = 7.7, p = .06$, and in subgroups matched on age, sex and IQ, $\chi^2(3, N = 51) = 6.7, p = .08$. Furthermore, a significant interaction was found between performance (clinical group is impaired or unimpaired) and diagnostic groups in the overall sample, $\chi^2(1, N = 60) = 5.7, p = .02$, and in the subgroups matched on age, sex and IQ, i.e., fewer studies showed that people with ASD were impaired than

ADHD on the cognitive control tasks $\chi^2(1, N = 51) = 6.3, p = .02$. The interaction was also observed when motor response inhibition tasks only were included in the analyses using all available data, $\chi^2(1, N = 42) = 14.1, p < .001$ and in age-, sex- and IQ-matched subgroups $\chi^2(1, N = 29) = 15.2, p < .001$.

Influences of Task Type and Performance in the Disorder-differentiating Findings during Cognitive Control

To assess the influence of task type, a meta-analysis comparing the functional impairment during cognitive control, covarying separately for task type and performance, in ASD and ADHD was conducted in the overall sample and matched subgroups. Covarying for task type, ASD-differentiating reduced activation in the ACC/midcingulate/dmPFC (overall = BA32/24/9/8/10; MNI coordinates: 0, 32, 22; $Z = 2.44, p < .00001$, 3065 voxels; matched subgroups = BA32/24/9/10; MNI coordinates: 2, 28, 20; $Z = 2.30, p < .00001$, 2994 voxels); and increased activations in precuneus (overall = BA5/4/23; MNI coordinates: 0, -42, 50; $Z = 1.88, p = .00002$, 1203 voxels; matched subgroups = BA5/4; MNI coordinates: 0, -42, 50; $Z = 1.82, p = .0001$, 706 voxels) and right inferior occipital lobe (overall = BA19; MNI coordinates: 36, -70, -8; $Z = 1.28, p = .0003$, 51 voxels; matched subgroups = BA19; MNI coordinates: 36, -68, -8; $Z = 1.43, p = .002$, 66 voxels) relative to ADHD remained significant.

Covarying for behavioural performance, the disorder-differentiating impairment in the ASD relative to the ADHD group remained in the above regions, i.e., precuneus (overall = BA5/4; MNI coordinates: 0, -42, 50; $Z = 1.32, p = .0001$, 960 voxels; matched subgroups = BA5/4; MNI coordinates: 0, -42, 50; $Z = 1.09, p = .0001$, 281 voxels), right inferior occipital lobe (overall = BA19; MNI coordinates: 36, -70, -8; $Z = 1.17, p = .0004$, 119 voxels; matched subgroups = BA5/23/4; MNI coordinates: 38, -68, -8; $Z = 1.36, p = .0002$, 244 voxels) and ACC/midcingulate/dMPFC (overall = BA32/24/9/8/ 10; MNI coordinates: 0, 32, 22; $Z = 2.69, p < .00001$, 3214 voxels; matched subgroups = BA32/24/9/10/8; MNI coordinates: 0, 32, 22; $Z = 2.72, p < .00001$, 3286 voxels).

Influences of Performance in the Disorder-differentiating Findings during Motor Response Inhibition

When covarying for performance during motor response inhibition data, we observed ASD-differentiating underactivation in left lingual gyrus/FFG/cerebellum lobule (overall = BA19/37/30/18; MNI coordinates: -10, -58, -14; $Z = 1.13$, $p = .001$, 388 voxels; matched subgroups = BA19/37/30/18; MNI coordinates: -16, -48, -12; $Z = 1.01$, $p = .0004$, 747 voxels), rACC/dmPFC (overall data only = BA5/4/23; MNI coordinates: 4, 34, 20; $Z = 1.23$, $p = .0008$, 587 voxels); and left MFG/dlPFC (overall data only = BA45/48; MNI coordinates: -46, 34, 16; $Z = 1.24$, $p = .0007$, 261 voxels) and right MFG/dlPFC (overall data only = BA9/6/44; MNI coordinates: 42, 10, 48; $Z = 1.09$, $p = .002$, 49 voxels). We also observed overactivation in right IOG/FFG (overall = BA19/37; MNI coordinates: 36, -68, -8; $Z = 1.57$, $p = .000002$, 434 voxels; matched subgroups = BA19/37/18; MNI coordinates: 34, -68, -6; $Z = 1.35$, $p = .0000002$, 524 voxels), left vlPFC/OFC (overall data only = BA11/10/47/46; MNI coordinates: -28, 38, -16; $Z = 1.30$, $p = .00002$, 868 voxels) and left posterior MTG (overall data only = BA37/21; MNI coordinates: -48, -54, 4; $Z = 1.41$, $p = .00002$, 289 voxels). The ADHD-differentiating underactivation in right IFG and caudate did not survive covarying for task performance.

Supplement 8: Jack-knife Analyses in Brain Structure and Function Abnormalities

Table S8a. Jack-knife Analyses of Reduced GMV Clusters among ADHD VBM Studies

Study excluded	L/R vmOFC/v mPFC/rAC C R caudate	R putamen/p osterior insula/ST G	L pre-CG	R rostromedial PFC	L vIPFC
Ahrendts et al. (2011)	Yes	Yes	Yes	Yes	Yes
Montes et al. (2010)	Yes	Yes	Yes	Yes	Yes
Amico et al. (2011)	Yes	Yes	Yes	Yes	Yes
Bonath et al. (2016)	Yes	Yes	Yes	Yes	Yes
Bralten et al. (2016)	Yes	Yes	No	No	Yes
Brieber et al. (2007)	Yes	Yes	Yes	Yes	Yes
Carmona et al. (2005)	Yes	Yes	Yes	Yes	Yes
Depue et al. (2010)	Yes	Yes	Yes	Yes	Yes
Gehricke et al. (2017)	Yes	Yes	Yes	Yes	Yes
He et al. (2015)	Yes	Yes	Yes	Yes	Yes
Iannaccone et al. (2015a)	Yes	Yes	Yes	No	Yes
Jagger-Rickels et al. (2018)	Yes	Yes	Yes	Yes	Yes
Johnston et al. (2014)	Yes	Yes	Yes	Yes	Yes
Kappel et al. (2015) – pediatric	Yes	Yes	Yes	Yes	Yes
Kappel et al. (2015) – adult	Yes	Yes	Yes	Yes	Yes
Kaya et al. (2018)	Yes	Yes	Yes	Yes	Yes
Kobel et al. (2010)	Yes	Yes	Yes	Yes	Yes
Kumar et al. (2017)	Yes	Yes	Yes	Yes	Yes
Li et al. (2015)	Yes	Yes	Yes	Yes	Yes
Lim et al. (2015)	Yes	Yes	Yes	Yes	Yes
Maier et al. (2015)	Yes	Yes	Yes	Yes	Yes
McAlonan et al. (2007)	Yes	Yes	Yes	Yes	Yes
Moreno-Alcázar et al. (2016)	Yes	Yes	Yes	Yes	Yes
Onnink et al. (2014)	Yes	Yes	Yes	Yes	Yes
Overmeyer et al. (2001)	Yes	Yes	Yes	Yes	Yes
Ramesh and Rai (2013)	Yes	Yes	Yes	Yes	Yes
Roman-Urrestarazu et al. (2016)	Yes	Yes	Yes	Yes	Yes
Saad et al. (2017)	Yes	Yes	Yes	Yes	Yes
Sasayama et al. (2010)	Yes	Yes	Yes	Yes	Yes
Seidman et al. (2011)	Yes	Yes	Yes	Yes	Yes
Sethi et al. (2017)	Yes	Yes	Yes	Yes	Yes
Shimada et al. (2015)	Yes	Yes	Yes	Yes	Yes
Stevens and Haney-Caron (2012)	Yes	Yes	Yes	Yes	Yes
van Wingen et al. (2013)	Yes	No	Yes	Yes	Yes
Vilgis et al. (2016)	Yes	Yes	Yes	Yes	Yes
Villemonteix et al. (2015)	Yes	Yes	Yes	Yes	Yes
Wang et al. (2007)	Yes	Yes	Yes	Yes	Yes
Yang et al. (2008)	Yes	Yes	Yes	Yes	Yes

L/R= left/right, vmOFC = ventromedial orbital prefrontal cortex, vmPFC = ventromedial prefrontal cortex, rACC = rostral anterior cingulate cortex, , STG = superior temporal gyrus, pre-CG = precentral gyrus, vIPFC = ventrolateral prefrontal cortex.

Table S8b. Jack-knife Analyses of GMV Abnormalities among ASD VBM Studies

Study excluded	Reduced				Enhanced		
	L/R dACC/dmPFC	L cerebellum	R hippocampus/PHG/FG	Dorsomedial thalamus	L ITG/MTG/STG/posterior insula	L/R PCC/precuneus	R MFG/SFG/dIPFC
Abell et al. (1999)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Boddaert et al. (2004)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Bonilha et al. (2008)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Brieber et al. (2007)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Cai et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Cheng et al. (2011)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Contarino et al. (2016)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Craig et al. (2007)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
D'Mello et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Ecker et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Foster et al. (2015)	Yes	No	Yes	Yes	Yes	Yes	Yes
Freitag et al. (2008)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Greimel et al. (2013)	No	Yes	Yes	Yes	Yes	Yes	Yes
Groen et al. (2011)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Hyde et al. (2010)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Itahashi et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Katz et al. (2016)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Kaufmann et al. (2013)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Ke et al. (2008)	Yes	Yes	No	Yes	Yes	Yes	Yes
Kosaka et al. (2010)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Kurth et al. (2011)	Yes	Yes	Yes	No	Yes	Yes	Yes
Kwon et al. (2004)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Langen et al. (2009)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Lim et al. (2015)	Yes	No	Yes	Yes	Yes	Yes	Yes
Lin et al. (2015) – pediatric (child)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Lin et al. (2015) – pediatric (adolescent)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Lin et al. (2015) – adult	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Lin et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
McAlonan et al. (2002)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
McAlonan et al. (2008)	Yes	No	Yes	No	Yes	Yes	Yes
Mengotti et al. (2011)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Mueller et al. (2013)	Yes	Yes	Yes	No	Yes	Yes	Yes
Ni et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Pereira et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Poulin-Lord et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Poustka et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Radeloff et al. (2014)	Yes	Yes	No	Yes	Yes	Yes	Yes
Retico et al. (2016)	Yes	Yes	Yes	Yes	Yes	No	No
Riedel et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Riva et al. (2013)	Yes	Yes	No	Yes	Yes	Yes	Yes
Rojas et al. (2006)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Sato et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Schmitz et al. (2006)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Toal et al. (2010)	Yes	Yes	No	Yes	Yes	Yes	Yes
Waiter et al. (2004)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Wang et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Wilson et al. (2009)	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Yang et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes

L/R = left/right, dACC = dorsal anterior cingulate cortex, dmPFC = dorsomedial prefrontal cortex, PHG = parahippocampal gyrus, FG = fusiform gyrus, ITG/MTG/STG = inferior/middle/superior temporal gyrus, PCC= posterior cingulate cortex, MFG/SFG = middle/superior frontal gyrus

Table S8c. Jack-knife Analyses of Abnormal Brain Function during Cognitive Control among ADHD fMRI Studies

	R thalamus/ caudate	L M/STG/s uperior temporal pole	L/R SMA/dor sal cingulate	L IFG/AI/te mporal pole	R AI/putam en	L post- CG	L MFG/ dIPFC	R MTG
Banich et al. (2009)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Bhajibwala et al. (2014)	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes
Booth et al. (2005)	Yes	Yes	No	Yes	Yes	Yes	Yes	No
Carmona et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No
Chantiluke et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Chen et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Chou et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
(Congdon et al., 2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Cubillo et al. (2010), Cubillo et al. (2011)*	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes
Cubillo et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Dibbets et al. (2010), Dibbets et al. (2009)*	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes
Durstun et al. (2006)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Fan et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Fan et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Hwang et al. (2015)	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
Iannaccone et al. (2015b)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Janssen et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Konrad et al. (2006)	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes
Kooistra et al. (2010)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Ma et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Ma et al. (2016)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Massat et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Passarotti et al. (2010)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes
Peterson et al. (2009)	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
Rasmussen et al. (2016)	Yes	Yes	Yes	Yes	No	No	No	Yes
Rubia et al. (2005)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No
Rubia et al. (2011a), Rubia et al. (2011b)*	Yes	Yes	No	No	No	Yes	Yes	Yes
Schulz et al. (2004)	Yes	Yes	Yes	Yes	Yes	No	Yes	No
Schulz et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Schulz et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No
Sebastian et al. (2012)	Yes	Yes	Yes	No	No	Yes	Yes	No
Shang et al. (2018)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes
Siniatchkin et al. (2012)	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes
Smith et al. (2006)	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes
Spinelli et al. (2011)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Szekely et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Tamm et al. (2004)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Thornton et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
van Hulst et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
van Rooij et al. (2015) – pediatric	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes
van Rooij et al. (2015) – adult	Yes	Yes	Yes	No	Yes	Yes	Yes	No
Zamorano et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes

L/R = left/right, M/STG = middle/superior temporal gyrus, SMA = supplementary motor area, IFG = inferior frontal gyrus, AI = anterior insula, post-CG = post-cingulate gyrus and MFG = middle frontal gyrus, dIPFC = dorsolateral prefrontal cortex. * Data sets were combined as they were collected from the same participants

Table S8d. Jack-knife Analyses of Abnormal Brain Function during Cognitive Control among ASD fMRI Studies

Study excluded	Underactivation					Overactivation				
	L/R ACC/midci ngulate/d mPFC	L MFG/ dIPFC	R MFG/ dIPFC	L IPL	L lingual gyrus/cere bellum IV/V	L precuneus/ midcingulate	R IOG	L vIPFC/OFC	L MFG/ rostrolateral PFC	R IFG
Ambrosino et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Chantiluke et al. (2015)	Yes	No	No	Yes	No	Yes	Yes	Yes	Yes	Yes
Daly et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Denisova et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Duerden et al. (2013)	Yes	Yes	No	Yes	Yes	Yes	No	Yes	Yes	No
Fan et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Gooskens et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Kana et al. (2007)	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	Yes
Kennedy et al. (2006)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Prat et al. (2016)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Schmitz et al. (2006)	Yes	Yes	Yes	Yes	Yes	No	Yes	No	No	Yes
Shafritz et al. (2008)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Shafritz et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Solomon et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Vaidya et al. (2011)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
van Hulst et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Velasquez et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Yerys et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No

L/R = left/right, ACC = anterior cingulate cortex, dmPFC = dorsomedial prefrontal cortex, MFG = middle frontal gyrus, dIPFC = dorsolateral prefrontal cortex, IPL =inferior parietal lobe, IOG = inferior occipital gyrus, vIPFC = ventrolateral prefrontal cortex, OFC = orbitofrontal cortex, IFG = inferior frontal gyrus.

Table S8e. Jack-knife Analyses of Abnormal Brain Function during Motor Response Inhibition among ADHD fMRI Studies

Study excluded	L MFG/ dIPFC	L anterior MTG/STG	L post-CG	R IFG	R vIPFC/ OFC/AI	R caudate
Bhaijiwala et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes
Booth et al. (2005)	Yes	Yes	Yes	Yes	Yes	No
Carmona et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes
Chantiluke et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes
Chen et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes
Congdon et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes
Cubillo et al. (2010)	Yes	Yes	Yes	Yes	Yes	Yes
Cubillo et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes
Dibbets et al. (2009)	Yes	Yes	Yes	Yes	Yes	Yes
Durston et al. (2006)	Yes	Yes	Yes	Yes	Yes	Yes
Iannaccone et al. (2015b)	Yes	Yes	Yes	Yes	Yes	Yes
Janssen et al. (2015)	Yes	Yes	Yes	Yes	Yes	Yes
Kooistra et al. (2010)	Yes	Yes	Yes	Yes	Yes	Yes
Ma et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes
Massat et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes
Passarotti et al. (2010)	Yes	Yes	Yes	Yes	Yes	Yes
Rasmussen et al. (2016)	Yes	Yes	No	No	Yes	No
Rubia et al. (2005)	Yes	Yes	Yes	Yes	Yes	Yes
Rubia et al. (2011b)	Yes	Yes	Yes	Yes	Yes	Yes
Schulz et al. (2004)	Yes	Yes	Yes	Yes	Yes	Yes
Schulz et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes
Sebastian et al. (2012)	Yes	Yes	Yes	Yes	Yes	Yes
Siniatchkin et al. (2012)	Yes	Yes	Yes	Yes	Yes	No
Smith et al. (2006)	Yes	Yes	Yes	Yes	Yes	Yes
Spinelli et al. (2011)	Yes	Yes	Yes	Yes	Yes	Yes
Szekely et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes
Tamm et al. (2004)	Yes	Yes	Yes	Yes	Yes	Yes
Thornton et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes
van Hulst et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes
van Rooij et al. (2015) – pediatric	Yes	Yes	No	Yes	Yes	Yes
van Rooij et al. (2015) – adult	No	Yes	Yes	Yes	Yes	Yes


L/R = left/right, MFG = middle frontal gyrus, dIPFC = dorsolateral prefrontal gyrus, M/STG = middle/superior temporal gyrus, post-CG = post-cingulate gyrus, IFG = inferior frontal gyrus, vIPFC = ventrolateral prefrontal cortex, OFC = orbitofrontal cortex, AI = anterior insula.

Table S8f. Jack-knife Analyses of Abnormal Brain Function during Motor Response Inhibition among ASD fMRI Studies

Study excluded	Underactivation				Overactivation	
	R AI/vIPFC	L cerebellu m/ lingual gyrus/ FFG	R MFG/ dIPFC	R PCC/ precuneus	L vIPFC/ OFC	R IOG/FFG/ ITG
Ambrosino et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes
Chantiluke et al. (2015)	Yes	No	No	Yes	Yes	No
Daly et al. (2014)	Yes	Yes	Yes	Yes	Yes	Yes
Duerden et al. (2013)	Yes	Yes	No	Yes	Yes	No
Gooskens et al. (2018)	Yes	Yes	Yes	Yes	Yes	Yes
Kana et al. (2007)	Yes	No	Yes	No	Yes	Yes
Prat et al. (2016)	Yes	Yes	Yes	Yes	Yes	Yes
Schmitz et al. (2006)	Yes	Yes	Yes	Yes	No	Yes
Shafritz et al. (2015)	No	Yes	Yes	Yes	Yes	Yes
van Hulst et al. (2017)	Yes	Yes	Yes	Yes	Yes	Yes
Velasquez et al. (2017)	Yes	Yes	Yes	No	Yes	Yes

L/R = left/right, AI = anterior insula, vIPFC = ventrolateral prefrontal cortex, FFG = fusiform gyrus, MFG= middle frontal gyrus, dIPFC = dorsolateral prefrontal cortex, PCC= posterior cingulate cortex, OFC = orbitofrontal cortex, IOG =inferior occipital gyrus, ITG = inferior temporal gyrus.

Figure S8. Pictorial Representation of the Jack-knife Analyses



Pictorial representation of jack-knife reliability analyses for structural and functional abnormalities in ASD and ADHD. Colormap indicate the level of replicability of the significant clusters according to the number of study combinations, ranging from 0 to maximum number of study combination for each comparison (see Tables 6a-f). Darker colours indicate that the cluster is replicable in fewer combinations of studies.

References

- Abell, F., Krams, M., Ashburner, J., Passingham, R., Friston, K., Frackowiak, R., . . . Frith, U. (1999). The neuroanatomy of autism: A voxel-based whole brain analysis of structural scans. *Neuroreport*, 10(8), 1647-1651. doi:10.1097/00001756-199906030-00005
- Ahrendts, J., Rüscher, N., Wilke, M., Philipsen, A., Eickhoff, S. B., Glauche, V., . . . Tebartz Van Elst, L. (2011). Visual cortex abnormalities in adults with ADHD: A structural MRI study. *The World Journal of Biological Psychiatry*, 12(4), 260-270. doi:10.3109/15622975.2010.518624
- Ambrosino, S., Bos, D. J., van Raalten, T. R., Kobussen, N. A., van Belle, J., Oranje, B., & Durston, S. (2014). Functional connectivity during cognitive control in children with autism spectrum disorder: An independent component analysis. *Journal of Neural Transmission*, 121(9), 1145-1155. doi:10.1007/s00702-014-1237-8
- Amico, F., Stauber, J., Koutsouleris, N., & Frodl, T. (2011). Anterior cingulate cortex gray matter abnormalities in adults with attention deficit hyperactivity disorder: A voxel-based morphometry study. *Psychiatry Research: Neuroimaging*, 191(1), 31-35. doi:10.1016/j.psychres.2010.08.011
- Banich, M. T., Burgess, G. C., Depue, B. E., Ruzic, L., Bidwell, L. C., Hitt-Laustsen, S., . . . Willcutt, E. G. (2009). The neural basis of sustained and transient attentional control in young adults with ADHD. *Neuropsychologia*, 47(14), 3095-3104. doi:10.1016/j.neuropsychologia.2009.07.005
- Bhaijiwala, M., Chevrier, A., & Schachar, R. (2014). Withholding and canceling a response in ADHD adolescents. *Brain and Behavior*, 4(5), 602-614. doi:10.1002/brb3.244
- Boddaert, N., Chabane, N., Gervais, H., Good, C. D., Bourgeois, M., Plumet, M. H., . . . Zilbovicius, M. (2004). Superior temporal sulcus anatomical abnormalities in childhood autism: A voxel-based morphometry MRI study. *NeuroImage*, 23(1), 364-369. doi:10.1016/j.neuroimage.2004.06.016
- Bonath, B., Tegelbeckers, J., Wilke, M., Flechtner, H. H., & Krauel, K. (2016). Regional gray matter volume differences between adolescents with ADHD and typically developing controls: Further evidence for anterior cingulate involvement. *Journal of Attention Disorders*. doi:10.1177/1087054715619682
- Bonilha, L., Cendes, F., Rorden, C., Eckert, M., Dalgalarrondo, P., Li, L. M., & Steiner, C. E. (2008). Gray and white matter imbalance - Typical structural abnormality underlying classic autism? *Brain and Development*, 30(6), 396-401. doi:10.1016/j.braindev.2007.11.006
- Booth, J. R., Burman, D. D., Meyer, J. R., Lei, Z., Trommer, B. L., Davenport, N. D., . . . Mesulam, M. M. (2005). Larger deficits in brain networks for response inhibition than for visual selective attention in attention deficit hyperactivity disorder (ADHD). *Journal of Child Psychology and Psychiatry*, 46(1), 94-111. doi:10.1111/j.1469-7610.2004.00337.x
- Bralten, J., Greven, C. U., Franke, B., Mennes, M., Zwiers, M. P., Rommelse, N. N. J., . . . Buitelaar, J. K. (2016). Voxel-based morphometry analysis reveals frontal brain differences in participants with ADHD and their unaffected siblings. *Journal of Psychiatry & Neuroscience*, 41(4), 272-279. doi:10.1503/jpn.140377
- Brieber, S., Neufang, S., Bruning, N., Kamp-Becker, I., Remschmidt, H., Herpertz-Dahlmann, B., . . . Konrad, K. (2007). Structural brain abnormalities in adolescents with autism spectrum disorder and patients with attention deficit/hyperactivity disorder. *Journal of Child Psychology and Psychiatry*, 48(12), 1251-1258. doi:10.1111/j.1469-7610.2007.01799.x
- Cai, J., Hu, X., Guo, K., Yang, P., Situ, M., & Huang, Y. (2018). Increased left inferior temporal gyrus was found in both low function autism and high function autism. *Frontiers in Psychiatry*, 9, 542. doi:10.3389/fpsy.2018.00542
- Carmona, S., Hoekzema, E., Antoni Ramos-Quiroga, J., Richarte, V., Canals, C., Bosch, R., . . . Vilarroya, O. (2012). Response inhibition and reward anticipation in medication-naïve adults with attention-deficit/hyperactivity disorder: A within-subject case-control neuroimaging study. *Human Brain Mapping*, 33(10), 2350-2361. doi:10.1002/hbm.21368

- Carmona, S., Vilarroya, O., Bielsa, A., Tremols, V., Soliva, J. C., Rovira, M., . . . Bulbena, A. (2005). Global and regional gray matter reductions in ADHD: A voxel-based morphometric study. *Neuroscience Letters*, 389(2), 88-93. doi:10.1016/j.neulet.2005.07.020
- Chantiluke, K., Barrett, N., Giampietro, V., Santosh, P., Brammer, M., Simmons, A., . . . Rubia, K. (2015). Inverse fluoxetine effects on inhibitory brain activation in non-comorbid boys with ADHD and with ASD. *Psychopharmacology*, 232(12), 2071-2082. doi:10.1007/s00213-014-3837-2
- Chen, C.-Y., Yen, J.-Y., Yen, C.-F., Chen, C.-S., Liu, G.-C., Liang, C.-Y., & Ko, C.-H. (2015). Aberrant brain activation of error processing among adults with attention deficit and hyperactivity disorder. *Kaohsiung Journal of Medical Sciences*, 31(4), 179-187. doi:10.1016/j.kjms.2015.01.001
- Cheng, Y., Chou, K. H., Fan, Y. T., & Lin, C. P. (2011). ANS: Aberrant neurodevelopment of the social cognition network in adolescents with autism spectrum disorders. *PLoS One*, 6(4). doi:10.1371/journal.pone.0018905
- Chou, T.-L., Chia, S., Shang, C.-Y., & Gau, S. S.-F. (2015). Differential therapeutic effects of 12-week treatment of atomoxetine and methylphenidate on drug-naïve children with attention deficit/hyperactivity disorder: A counting Stroop functional MRI study. *European Neuropsychopharmacology*, 25(12), 2300-2310. doi:10.1016/j.euroneuro.2015.08.024
- Congdon, E., Altshuler, L. L., Mumford, J. A., Karlsgodt, K. H., Sabb, F. W., Ventura, J., . . . Poldrack, R. A. (2014). Neural activation during response inhibition in adult attention-deficit/hyperactivity disorder: Preliminary findings on the effects of medication and symptom severity. *Psychiatry Research: Neuroimaging*, 222(1-2), 17-28. doi:10.1016/j.psychresns.2014.02.002
- Contarino, V. E., Bulgheroni, S., Annunziata, S., Erbetta, A., & Riva, D. (2016). Widespread focal cortical alterations in autism spectrum disorder with intellectual disability detected by threshold-free cluster enhancement. *American Journal of Neuroradiology*, 37(9), 1721-1726. doi:10.3174/ajnr.A4779
- Craig, M., Zaman, S. H., Daly, E. M., Cutter, W. J., Robertson, D. M. W., Hallahan, B., . . . Murphy, D. G. M. (2007). Women with autistic-spectrum disorder: Magnetic resonance imaging study of brain anatomy. *British Journal of Psychiatry*, 191, 224-228. doi:10.1192/bjp.bp.106.034603
- Cubillo, A., Halari, R., Ecker, C., Giampietro, V., Taylor, E., & Rubia, K. (2010). Reduced activation and inter-regional functional connectivity of fronto-striatal networks in adults with childhood attention-deficit hyperactivity disorder (ADHD) and persisting symptoms during tasks of motor inhibition and cognitive switching. *Journal of Psychiatric Research*, 44(10), 629-639. doi:10.1016/j.jpsychires.2009.11.016
- Cubillo, A., Halari, R., Giampietro, V., Taylor, E., & Rubia, K. (2011). Fronto-striatal underactivation during interference inhibition and attention allocation in grown up children with attention deficit/hyperactivity disorder and persistent symptoms. *Psychiatry Research*, 193(1), 17-27. doi:10.1016/j.psychresns.2010.12.014
- Cubillo, A., Smith, A. B., Barrett, N., Giampietro, V., Brammer, M. J., Simmons, A., & Rubia, K. (2014). Shared and drug-specific effects of atomoxetine and methylphenidate on inhibitory brain dysfunction in medication-naïve ADHD boys. *Cerebral Cortex*, 24(1), 174-185. doi:10.1093/cercor/bhs296
- D'Mello, A. M., Crocetti, D., Mostofsky, S. H., & Stoodley, C. J. (2015). Cerebellar gray matter and lobular volumes correlate with core autism symptoms. *NeuroImage: Clinical*, 7, 631-639. doi:10.1016/j.nicl.2015.02.007
- Daly, E. M., Ecker, C., Hallahan, B., Deeley, Q., Craig, M., Murphy, C., . . . Murphy, D. G. M. (2014). Response inhibition and serotonin in autism: A functional MRI study using acute tryptophan depletion. *Brain*, 137, 2600-2610. doi:10.1093/brain/awu178
- Denisova, K., Zhao, G., Wang, Z., Goh, S., Huo, Y., & Peterson, B. S. (2017). Cortical interactions during the resolution of information processing demands in autism spectrum disorders. *Brain and Behavior*, 7(2), e00596. doi:10.1002/brb3.596
- Depue, B. E., Burgess, G. C., Bidwell, L. C., Willcutt, E. G., & Banich, M. T. (2010). Behavioral performance predicts grey matter reductions in the right inferior frontal gyrus in young

- adults with combined type ADHD. *Psychiatry Research: Neuroimaging*, 182(3), 231-237. doi:10.1016/j.psychres.2010.01.012
- Dibbets, P., Evers, E. A. T., Hurks, P. P. M., Bakker, K., & Jolles, J. (2010). Differential brain activation patterns in adult attention-deficit hyperactivity disorder (ADHD) associated with task switching. *Neuropsychology*, 24(4), 413-423. doi:10.1037/a0018997
- Dibbets, P., Evers, L., Hurks, P., Marchetta, N., & Jolles, J. (2009). Differences in feedback- and inhibition-related neural activity in adult ADHD. *Brain and Cognition*, 70(1), 73-83. doi:10.1016/j.bandc.2009.01.001
- Duerden, E. G., Taylor, M. J., Soorya, L. V., Wang, T., Fan, J., & Anagnostou, E. (2013). Neural correlates of inhibition of socially relevant stimuli in adults with autism spectrum disorder. *Brain Research*, 1533, 80-90. doi:10.1016/j.brainres.2013.08.021
- Durston, S., Mulder, M., Casey, B. J., Ziermans, T., & van Engeland, H. (2006). Activation in ventral prefrontal cortex is sensitive to genetic vulnerability for attention-deficit hyperactivity disorder. *Biological Psychiatry*, 60(10), 1062-1070. doi:10.1016/j.biopsych.2005.12.020
- Ecker, C., Suckling, J., Deoni, S. C., Lombardo, M. V., Bullmore, E. T., Baron-Cohen, S., . . . Murphy, D. G. M. (2012). Brain anatomy and its relationship to behavior in adults with autism spectrum disorder. *Archives of General Psychiatry*, 69(2), 195-209. doi:10.1001/archgenpsychiatry.2011.1251
- Fan, J., Bernardi, S., Van Dam, N. T., Anagnostou, E., Gu, X., Martin, L., . . . Hof, P. R. (2012). Functional deficits of the attentional networks in autism. *Brain and Behavior*, 2(5), 647-660. doi:10.1002/brb3.90
- Fan, L. Y., Chou, T. L., & Gau, S. S. (2017). Neural correlates of atomoxetine improving inhibitory control and visual processing in drug-naïve adults with attention-deficit/hyperactivity disorder. *Human Brain Mapping*, 38(10), 4850-4864. doi:10.1002/hbm.23683
- Fan, L. Y., Shang, C. Y., Tseng, W. I., Gau, S. S., & Chou, T. L. (2018). Visual processing as a potential endophenotype in youths with attention-deficit/hyperactivity disorder: A sibling study design using the counting Stroop functional MRI. *Human Brain Mapping*, 39(10), 3827-3835. doi:10.1002/hbm.24214
- Foster, N. E. V., Doyle-Thomas, K. A. R., Tryfon, A., Ouimet, T., Anagnostou, E., Evans, A. C., . . . Hyde, K. L. (2015). Structural gray matter differences during childhood development in autism spectrum disorder: A multimetric approach. *Pediatric Neurology*, 53(4), 350-359. doi:10.1016/j.pediatrneurol.2015.06.013
- Freitag, C. M., Konrad, C., Häberlen, M., Kleser, C., von Gontard, A., Reith, W., . . . Krick, C. (2008). Perception of biological motion in autism spectrum disorders. *Neuropsychologia*, 46(5), 1480-1494. doi:10.1016/j.neuropsychologia.2007.12.025
- Gehricke, J. G., Kruggel, F., Thampipop, T., Alejo, S. D., Tatos, E., Fallon, J., & Muftuler, L. T. (2017). The brain anatomy of attention-deficit/hyperactivity disorder in young adults - a magnetic resonance imaging study. *PLoS One*, 12(4). doi:10.1371/journal.pone.0175433
- Gooskens, B., Bos, D. J., Mensen, V. T., Shook, D. A., Bruchhage, M. M. K., Naaijen, J., . . . consortium, T. (2018). No evidence of differences in cognitive control in children with autism spectrum disorder or obsessive-compulsive disorder: An fMRI study. *Developmental Cognitive Neuroscience*, 100602. doi:10.1016/j.dcn.2018.11.004
- Greimel, E., Nehrkorn, B., Schulte-Ruether, M., Fink, G. R., Nickl-Jockschat, T., Herpertz-Dahlmann, B., . . . Eickhoff, S. B. (2013). Changes in grey matter development in autism spectrum disorder. *Brain Structure and Function*, 218(4), 929-942. doi:10.1007/s00429-012-0439-9
- Groen, W. B., Buitelaar, J. K., van der Gaag, R. J., & Zwiers, M. P. (2011). Pervasive microstructural abnormalities in autism: A DTI study. *Journal of Psychiatry & Neuroscience*, 36(1), 32-40. doi:10.1503/jpn.090100
- He, N., Li, F., Li, Y., Guo, L., Chen, L., Huang, X., . . . Gong, Q. (2015). Neuroanatomical deficits correlate with executive dysfunction in boys with attention deficit hyperactivity disorder. *Neuroscience Letters*, 600, 45-49. doi:10.1016/j.neulet.2015.05.062
- Hwang, S., White, S. F., Nolan, Z. T., Craig Williams, W., Sinclair, S., & Blair, R. J. R. (2015). Executive attention control and emotional responding in attention-deficit/hyperactivity

- disorder - A functional MRI study. *NeuroImage: Clinical*, 9, 545-554.
doi:10.1016/j.nicl.2015.10.005
- Hyde, K. L., Samson, F., Evans, A. C., & Mottron, L. (2010). Neuroanatomical differences in brain areas implicated in perceptual and other core features of autism revealed by cortical thickness analysis and voxel-based morphometry. *Human Brain Mapping*, 31(4), 556-566. doi:10.1002/hbm.20887
- Iannaccone, R., Hauser, T. U., Ball, J., Brandeis, D., Walitza, S., & Brem, S. (2015). Classifying adolescent attention-deficit/hyperactivity disorder (ADHD) based on functional and structural imaging. *European Child & Adolescent Psychiatry*, 24(10), 1279-1289. doi:10.1007/s00787-015-0678-4
- Iannaccone, R., Hauser, T. U., Staempfli, P., Walitza, S., Brandeis, D., & Brem, S. (2015). Conflict monitoring and error processing: New insights from simultaneous EEG-fMRI. *NeuroImage*, 105, 395-407. doi:10.1016/j.neuroimage.2014.10.028
- Itahashi, T., Yamada, T., Nakamura, M., Watanabe, H., Yamagata, B., Jimbo, D., . . . Hashimoto, R. (2015). Linked alterations in gray and white matter morphology in adults with high-functioning autism spectrum disorder: A multimodal brain imaging study. *NeuroImage: Clinical*, 7, 155-169. doi:10.1016/j.nicl.2014.11.019
- Jagger-Rickels, A. C., Kibby, M. Y., & Constance, J. M. (2018). Global gray matter morphometry differences between children with reading disability, ADHD, and comorbid reading disability/ADHD. *Brain and Language*, 185, 54-66. doi:10.1016/j.bandl.2018.08.004
- Janssen, T. W. P., Heslenfeld, D. J., Mourik, R. V., Logan, G. D., & Oosterlaan, J. (2015). Neural correlates of response inhibition in children with attention-deficit/hyperactivity disorder: A controlled version of the stop-signal task. *Psychiatry Research: Neuroimaging*, 233(2), 278-284. doi:10.1016/j.psychresns.2015.07.007
- Johnston, B. A., Mwangi, B., Matthews, K., Coghill, D., Konrad, K., & Steele, J. D. (2014). Brainstem abnormalities in attention deficit hyperactivity disorder support high accuracy individual diagnostic classification. *Human Brain Mapping*, 35(10), 5179-5189. doi:10.1002/hbm.22542
- Kana, R. K., Keller, T. A., Minshew, N. J., & Just, M. A. (2007). Inhibitory control in high-functioning autism: Decreased activation and underconnectivity in inhibition networks. *Biological Psychiatry*, 62(3), 198-206. doi:10.1016/j.biopsych.2006.08.004
- Kappel, V., Lorenz, R. C., Streifling, M., Renneberg, B., Lehmkuhl, U., Stroehle, A., . . . Beck, A. (2015). Effect of brain structure and function on reward anticipation in children and adults with attention deficit hyperactivity disorder combined subtype. *Social Cognitive and Affective Neuroscience*, 10(7), 945-951. doi:10.1093/scan/nsu135
- Katz, J., d'Albis, M. A., Boisgontier, J., Poupon, C., Mangin, J. F., Guevara, P., . . . Houenou, J. (2016). Similar white matter but opposite grey matter changes in schizophrenia and high-functioning autism. *Acta Psychiatrica Scandinavica*, 134(1), 31-39. doi:10.1111/acps.12579
- Kaufmann, L., Zotter, S., Pixner, S., Starke, M., Haberlandt, E., Steinmayr-Gensluckner, M., . . . Marksteiner, J. (2013). Brief report: CANTAB performance and brain structure in pediatric patients with Asperger syndrome. *Journal of Autism and Developmental Disorders*, 43(6), 1483-1490. doi:10.1007/s10803-012-1686-6
- Kaya, B. S., Metin, B., Tas, Z. C., Buyukaslan, A., Soysal, A., Hatiloglu, D., & Tarhan, N. (2018). Gray matter increase in motor cortex in pediatric ADHD: A voxel-based morphometry study. *Journal of attention disorders*, 22(7), 611-618. doi:10.1177/1087054716659139
- Ke, X., Hong, S., Tang, T., Zou, B., Li, H., Hang, Y., . . . Liu, Y. (2008). Voxel-based morphometry study on brain structure in children with high-functioning autism. *Neuroreport*, 19(9), 921-925. doi:10.1097/wnr.0b013e328300edf3
- Kennedy, D. P., Redcay, E., & Courchesne, E. (2006). Failing to deactivate: Resting functional abnormalities in autism. *Proceedings of the National Academy of Sciences of the United States of America*, 103(21), 8275-8280. doi:10.1073/pnas.0600674103
- Kobel, M., Bechtel, N., Specht, K., Klarhoefer, M., Weber, P., Scheffler, K., . . . Penner, I.-K. (2010). Structural and functional imaging approaches in attention deficit/hyperactivity disorder: Does the temporal lobe play a key role? *Psychiatry Research: Neuroimaging*, 183(3), 230-236. doi:10.1016/j.psychresns.2010.03.010

- Konrad, K., Neufang, S., Hanisch, C., Fink, G. R., & Herpertz-Dahlmann, B. (2006). Dysfunctional attentional networks in children with attention deficit/hyperactivity disorder: Evidence from an event-related functional magnetic resonance imaging study. *Biological Psychiatry*, 59(7), 643-651. doi:10.1016/j.biopsych.2005.08.013
- Kooistra, L., van der Meere, J. J., Edwards, J. D., Kaplan, B. J., Crawford, S., & Goodyear, B. G. (2010). Preliminary fMRI findings on the effects of event rate in adults with ADHD. *Journal of Neural Transmission*, 117(5), 655-662. doi:10.1007/s00702-010-0374-y
- Kosaka, H., Omori, M., Munesue, T., Ishitobi, M., Matsumura, Y., Takahashi, T., . . . Wada, Y. (2010). Smaller insula and inferior frontal volumes in young adults with pervasive developmental disorders. *NeuroImage*, 50(4), 1357-1363. doi:10.1016/j.neuroimage.2010.01.085
- Kumar, U., Arya, A., & Agarwal, V. (2017). Neural alterations in ADHD children as indicated by voxel-based cortical thickness and morphometry analysis. *Brain and Development*, 39(5), 403-410. doi:10.1016/j.braindev.2016.12.002
- Kurth, F., Narr, K. L., Woods, R. P., O'Neill, J., Alger, J. R., Caplan, R., . . . Levitt, J. G. (2011). Diminished gray matter within the hypothalamus in autism disorder: A potential link to hormonal effects? *Biological Psychiatry*, 70(3), 278-282. doi:10.1016/j.biopsych.2011.03.026
- Kwon, H., Ow, A. W., Pedatella, K. E., Lotspeich, L. J., & Reiss, A. L. (2004). Voxel-based morphometry elucidates structural neuroanatomy of high-functioning autism and Asperger syndrome. *Developmental Medicine & Child Neurology*, 46(11), 760-764. doi:10.1017/s0012162204001306
- Langen, M., Schnack, H. G., Nederveen, H., Bos, D., Lahuis, B. E., de Jonge, M. V., . . . Durston, S. (2009). Changes in the developmental trajectories of striatum in autism. *Biological Psychiatry*, 66(4), 327-333. doi:10.1016/j.biopsych.2009.03.017
- Li, X., Cao, Q., Pu, F., Li, D., Fan, Y., An, L., . . . Wang, Y. (2015). Abnormalities of structural covariance networks in drug-naïve boys with attention deficit hyperactivity disorder. *Psychiatry Research*, 231(3), 273-278. doi:10.1016/j.psychres.2015.01.006
- Lim, L., Chantiluke, K., Cubillo, A. I., Smith, A. B., Simmons, A., Mehta, M. A., & Rubia, K. (2015). Disorder-specific grey matter deficits in attention deficit hyperactivity disorder relative to autism spectrum disorder. *Psychological Medicine*, 45(5), 965-976. doi:10.1017/s0033291714001974
- Lin, H. Y., Ni, H.-C., Lai, M.-C., Tseng, W.-Y. I., & Gau, S. S.-F. (2015). Regional brain volume differences between males with and without autism spectrum disorder are highly age-dependent. *Molecular Autism*, 6, 29. doi:10.1186/s13229-015-0022-3
- Lin, H. Y., Tseng, W. I., Lai, M. C., Chang, Y. T., & Gau, S. S. (2017). Shared atypical brain anatomy and intrinsic functional architecture in male youth with autism spectrum disorder and their unaffected brothers. *Psychological Medicine*, 47(4), 639-654. doi:10.1017/S0033291716002695
- Ma, I., van Holstein, M., Mies, G. W., Mennes, M., Buitelaar, J., Cools, R., . . . Scheres, A. (2016). Ventral striatal hyperconnectivity during rewarded interference control in adolescents with ADHD. *Cortex*, 82, 225-236. doi:10.1016/j.cortex.2016.05.021
- Ma, J., Lei, D., Jin, X., Du, X., Jiang, F., Li, F., . . . Shen, X. (2012). Compensatory brain activation in children with attention deficit/hyperactivity disorder during a simplified go/no-go task. *Journal of Neural Transmission*, 119(5), 613-619. doi:10.1007/s00702-011-0744-0
- Maier, S., Perlov, E., Graf, E., Dieter, E., Sobanski, E., Rump, M., . . . Elst, L. T. v. (2015). Discrete global but no focal gray matter volume reductions in unmedicated adult patients with attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 80(12), 905-915. doi:10.1016/j.biopsych.2015.05.012
- Massat, I., Slama, H., Villemonteix, T., Mary, A., Baijot, S., Albajara Saenz, A., . . . Peigneux, P. (2018). Hyperactivity in motor response inhibition networks in unmedicated children with attention deficit-hyperactivity disorder. *The World Journal of Biological Psychiatry*, 19(2), 101-111. doi:10.1080/15622975.2016.1237040
- McAlonan, G. M., Cheung, V., Cheung, C., Chua, S. E., Murphy, D. G. M., Suckling, J., . . . Ho, T. P. (2007). Mapping brain structure in attention deficit-hyperactivity disorder: A voxel-

- based MRI study of regional grey and white matter volume. *Psychiatry Research-Neuroimaging*, 154(2), 171-180. doi:10.1016/j.psychresns.2006.09.006
- McAlonan, G. M., Daly, E., Kumari, V., Critchley, H. D., van Amelsvoort, T., Suckling, J., . . . Murphy, D. G. M. (2002). Brain anatomy and sensorimotor gating in Asperger's syndrome. *Brain*, 125, 1594-1606. doi:10.1093/brain/awf150
- McAlonan, G. M., Suckling, J., Wong, N., Cheung, V., Lienenkaemper, N., Cheung, C., & Chua, S. E. (2008). Distinct patterns of grey matter abnormality in high-functioning autism and Asperger's syndrome. *Journal of Child Psychology and Psychiatry*, 49(12), 1287-1295. doi:10.1111/j.1469-7610.2008.01933.x
- Mengotti, P., D'Agostini, S., Terlevic, R., De Colle, C., Biasizzo, E., Londero, D., . . . Brambilla, P. (2011). Altered white matter integrity and development in children with autism: A combined voxel-based morphometry and diffusion imaging study. *Brain Research Bulletin*, 84(2), 189-195. doi:10.1016/j.brainresbull.2010.12.002
- Montes, L. G. A., Ricardo-Garcell, J., de la Torre, L. B., Alcántara, H. P., García, R. B. M., Fernández-Bouzas, A., & Acosta, D. A. (2010). Clinical correlations of grey matter reductions in the caudate nucleus of adults with attention deficit hyperactivity disorder. *Journal of Psychiatry and Neuroscience*, 35(4), 238-246. doi:10.1503/jpn.090099
- Moreno-Alcázar, A., Ramos-Quiroga, J. A., Radua, J., Salavert, J., Palomar, G., Bosch, R., . . . Pomarol-Clotet, E. (2016). Brain abnormalities in adults with attention deficit hyperactivity disorder revealed by voxel-based morphometry. *Psychiatry Research: Neuroimaging*, 254, 41-47. doi:10.1016/j.psychresns.2016.06.002
- Mueller, S., Keeser, D., Samson, A. C., Kirsch, V., Blautzik, J., Grothe, M., . . . Meindl, T. (2013). Convergent findings of altered functional and structural brain connectivity in individuals with high functioning autism: A multimodal MRI study. *PLoS One*, 8(6), e67329. doi:10.1371/journal.pone.0067329
- Ni, H. C., Lin, H. Y., Tseng, W. I., Chiu, Y. N., Wu, Y. Y., Tsai, W. C., & Gau, S. S. (2018). Neural correlates of impaired self-regulation in male youths with autism spectrum disorder: A voxel-based morphometry study. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 82, 233-241. doi:10.1016/j.pnpbp.2017.11.008
- Norman, L. J., Carlisi, C., Lukito, S., Hart, H., Mataix-Cols, D., Radua, J., & Rubia, K. (2016). Structural and functional brain abnormalities in attention-deficit/hyperactivity disorder and obsessive-compulsive disorder: A comparative meta-analysis. *JAMA Psychiatry*. doi:10.1001/jamapsychiatry.2016.0700
- Onnink, A. M. H., Zwiers, M. P., Hoogman, M., Mostert, J. C., Kan, C. C., Buitelaar, J., & Franke, B. (2014). Brain alterations in adult ADHD: Effects of gender, treatment and comorbid depression. *European Neuropsychopharmacology*, 24(3), 397-409. doi:10.1016/j.euroneuro.2013.11.011
- Overmeyer, S., Bullmore, E. T., Suckling, J., Simmons, A., Williams, S. C., Santosh, P. J., & Taylor, E. (2001). Distributed grey and white matter deficits in hyperkinetic disorder: MRI evidence for anatomical abnormality in an attentional network. *Psychological Medicine*, 31(8), 1425-1435. doi:10.1017/s0033291701004706
- Passarotti, A. M., Sweeney, J. A., & Pavuluri, M. N. (2010). Neural correlates of response inhibition in pediatric bipolar disorder and attention deficit hyperactivity disorder. *Psychiatry Research: Neuroimaging*, 181(1), 36-43. doi:10.1016/j.psychresns.2009.07.002
- Pereira, A. M., Campos, B. M., Coan, A. C., Pegoraro, L. F., de Rezende, T. J. R., Obeso, I., . . . Cendes, F. (2018). Differences in cortical structure and functional mri connectivity in high functioning autism. *Frontiers in Neurology*, 9, 539. doi:10.3389/fneur.2018.00539
- Peterson, B. S., Potenza, M. N., Wang, Z., Zhu, H., Martin, A., Marsh, R., . . . Yu, S. (2009). An fMRI study of the effects of psychostimulants on default-mode processing during Stroop task performance in youths with ADHD. *American Journal of Psychiatry*, 166(11), 1286-1294. doi:10.1176/appi.ajp.2009.08050724
- Poulin-Lord, M. P., Barbeau, E. B., Soulières, I., Monchi, O., Doyon, J., Benali, H., & Motttron, L. (2014). Increased topographical variability of task-related activation in perceptive and motor associative regions in adult autistics. *NeuroImage: Clinical*, 4, 444-453. doi:10.1016/j.nicl.2014.02.008

- Poustka, L., Jennen-Steinmetz, C., Henze, R., Vomstein, K., Haffner, J., & Sieltjes, B. (2012). Fronto-temporal disconnectivity and symptom severity in children with autism spectrum disorder. *The World Journal of Biological Psychiatry*, 13(4), 269-280. doi:10.3109/15622975.2011.591824
- Prat, C. S., Stocco, A., Neuhaus, E., & Kleinhaus, N. M. (2016). Basal ganglia impairments in autism spectrum disorder are related to abnormal signal gating to prefrontal cortex. *Neuropsychologia*, 91, 268-281. doi:10.1016/j.neuropsychologia.2016.08.007
- Radeloff, D., Ciaramidaro, A., Siniatchkin, M., Hainz, D., Schlitt, S., Weber, B., . . . Freitag, C. M. (2014). Structural alterations of the social brain: A comparison between schizophrenia and autism. *PLoS One*, 9(9), e106539. doi:10.1371/journal.pone.0106539
- Radua, J., Mataix-Cols, D., Phillips, M. L., El-Hage, W., Kronhaus, D. M., Cardoner, N., & Surguladze, S. (2012). A new meta-analytic method for neuroimaging studies that combines reported peak coordinates and statistical parametric maps. *European Psychiatry*, 27(8), 605-611. doi:10.1016/j.eurpsy.2011.04.001
- Ramesh, M. G., & Rai, K. S. (2013). Region-wise gray matter volume alterations in brain of adolescents with attention deficit hyperactive disorder: A voxel based morphometric analysis. *Indian Journal of Physiology and Pharmacology*, 57(3), 270-279. Retrieved from <https://www.researchgate.net/publication/277133055>
- Rasmussen, J., Casey, B. J., van Erp, T. G., Tamm, L., Epstein, J. N., Buss, C., . . . Group, M. T. A. N. (2016). ADHD and cannabis use in young adults examined using fMRI of a Go/NoGo task. *Brain Imaging and Behavior*, 10(3), 761-771. doi:10.1007/s11682-015-9438-9
- Retico, A., Giuliano, A., Tancredi, R., Cosenza, A., Apicella, F., Narzisi, A., . . . Calderoni, S. (2016). The effect of gender on the neuroanatomy of children with autism spectrum disorders: A support vector machine case-control study. *Molecular Autism*, 7, 5. doi:10.1186/s13229-015-0067-3
- Riedel, A., Maier, S., Ulbrich, M., Biscaldi, M., Ebert, D., Fangmeier, T., . . . Elst, L. T. v. (2014). No significant brain volume decreases or increases in adults with high-functioning autism spectrum disorder and above average intelligence: A voxel-based morphometric study. *Psychiatry Research: Neuroimaging*, 223(2), 67-74. doi:10.1016/j.psychres.2014.05.013
- Riva, D., Annunziata, S., Contarino, V., Erbetta, A., Aquino, D., & Bulgheroni, S. (2013). Gray matter reduction in the vermis and CRUS-II is associated with social and interaction deficits in low-functioning children with autistic spectrum disorders: A VBM-DARTEL study. *Cerebellum*, 12(5), 676-685. doi:10.1007/s12311-013-0469-8
- Rojas, D. C., Peterson, E., Winterrowd, E., Reite, M. L., Rogers, S. J., & Tregellas, J. R. (2006). Regional gray matter volumetric changes in autism associated with social and repetitive behavior symptoms. *BMC Psychiatry*, 6, 56. doi:10.1186/1471-244x-6-56
- Roman-Urrestarazu, A., Lindholm, P., Moilanen, I., Kiviniemi, V., Miettunen, J., Jaaskelainen, E., . . . Murray, G. K. (2016). Brain structural deficits and working memory fMRI dysfunction in young adults who were diagnosed with ADHD in adolescence. *European Child & Adolescent Psychiatry*, 25(5), 529-538. doi:10.1007/s00787-015-0755-8
- Rubia, K., Halari, R., Cubillo, A., Smith, A. B., Mohammad, A. M., Brammer, M., & Taylor, E. (2011). Methylphenidate normalizes fronto-striatal underactivation during interference inhibition in medication-naïve boys with attention-deficit hyperactivity disorder. *Neuropsychopharmacology*, 36(8), 1575-1586. doi:10.1038/npp.2011.30
- Rubia, K., Halari, R., Mohammad, A. M., Taylor, E., & Brammer, M. (2011). Methylphenidate normalizes frontocingulate underactivation during error processing in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 70(3), 255-262. doi:10.1016/j.biopsych.2011.04.018
- Rubia, K., Smith, A. B., Brammer, M. J., Toone, B., & Taylor, E. (2005). Abnormal brain activation during inhibition and error detection in medication-naïve adolescents with ADHD. *American Journal of Psychiatry*, 162(6), 1067-1075. doi:10.1176/appi.ajp.162.6.1067
- Saad, J. F., Griffiths, K. R., Kohn, M. R., Clarke, S., Williams, L. M., & Korgaonkar, M. S. (2017). Regional brain network organization distinguishes the combined and inattentive

- subtypes of attention deficit hyperactivity disorder. *NeuroImage: Clinical*, 15, 383-390. doi:10.1016/j.nicl.2017.05.016
- Sasayama, D., Hayashida, A., Yamasue, H., Harada, Y., Kaneko, T., Kasai, K., . . . Amano, N. (2010). Neuroanatomical correlates of attention-deficit-hyperactivity disorder accounting for comorbid oppositional defiant disorder and conduct disorder. *Psychiatry and Clinical Neurosciences*, 64(4), 394-402. doi:10.1111/j.1440-1819.2010.02102.x
- Sato, W., Uono, S., Kochiyama, T., Yoshimura, S., Sawada, R., Kubota, Y., . . . Toichi, M. (2017). Structural correlates of reading the mind in the eyes in autism spectrum disorder. *Frontiers in Human Neuroscience*, 11, 361. doi:10.3389/fnhum.2017.00361
- Schmitz, N., Rubia, K., Daly, E., Smith, A., Williams, S., & Murphy, D. G. (2006). Neural correlates of executive function in autistic spectrum disorders. *Biological Psychiatry*, 59(1), 7-16. doi:10.1016/j.biopsych.2005.06.007
- Schulz, K. P., Bédard, A. C. V., Fan, J., Clerkin, S. M., Dima, D., Newcorn, J. H., & Halperin, J. M. (2014). Emotional bias of cognitive control in adults with childhood attention-deficit/hyperactivity disorder. *NeuroImage: Clinical*, 5, 1-9. doi:10.1016/j.nicl.2014.05.016
- Schulz, K. P., Fan, J., Tang, C. Y., Newcorn, J. H., Buchsbaum, M. S., Cheung, A. M., & Halperin, J. M. (2004). Response inhibition in adolescents diagnosed with attention deficit hyperactivity disorder during childhood: An event-related fMRI study. *American Journal of Psychiatry*, 161(9), 1650-1657. doi:10.1176/appi.ajp.161.9.1650
- Schulz, K. P., Li, X., Clerkin, S. M., Fan, J., Berwid, O. G., Newcorn, J. H., & Halperin, J. M. (2017). Prefrontal and parietal correlates of cognitive control related to the adult outcome of attention-deficit/hyperactivity disorder diagnosed in childhood. *Cortex*, 90, 1-11. doi:10.1016/j.cortex.2017.01.019
- Sebastian, A., Gerdes, B., Feige, B., Kloeppel, S., Lange, T., Philipsen, A., . . . Tiescher, O. (2012). Neural correlates of interference inhibition, action withholding and action cancellation in adult ADHD. *Psychiatry Research: Neuroimaging*, 202(2), 132-141. doi:10.1016/j.psychres.2012.02.010
- Seidman, L. J., Biederman, J., Liang, L., Valera, E. M., Monuteaux, M. C., Brown, A., . . . Makris, N. (2011). Gray matter alterations in adults with attention-deficit/hyperactivity disorder identified by voxel based morphometry. *Biological Psychiatry*, 69(9), 857-866. doi:10.1016/j.biopsych.2010.09.053
- Sethi, A., Evelyn-Rahr, E., Dowell, N., Jain, S., Voon, V., Critchley, H. D., . . . Cercignani, M. (2017). Magnetization transfer imaging identifies basal ganglia abnormalities in adult ADHD that are invisible to conventional T1 weighted voxel-based morphometry. *NeuroImage: Clinical*, 15, 8-14. doi:10.1016/j.nicl.2017.03.012
- Shafritz, K. M., Bregman, J. D., Ikuta, T., & Szeszko, P. R. (2015). Neural systems mediating decision-making and response inhibition for social and nonsocial stimuli in autism. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 60, 112-120. doi:10.1016/j.pnpbp.2015.03.001
- Shafritz, K. M., Dichter, G. S., Baranek, G. T., & Belger, A. (2008). The neural circuitry mediating shifts in behavioral response and cognitive set in autism. *Biological Psychiatry*, 63(10), 974-980. doi:10.1016/j.biopsych.2007.06.028
- Shang, C. Y., Sheng, C., Yang, L. K., Chou, T. L., & Gau, S. S. (2018). Differential brain activations in adult attention-deficit/ hyperactivity disorder subtypes: A counting Stroop functional MRI study. *Brain Imaging and Behavior*, 12(3), 882-890. doi:10.1007/s11682-017-9749-0
- Shimada, K., Fujisawa, T. X., Takiguchi, S., Naruse, H., Kosaka, H., Okazawa, H., & Tomoda, A. (2015). Ethnic differences in COMT genetic effects on striatal grey matter alterations associated with childhood ADHD: A voxel-based morphometry study in a Japanese sample. *The World Journal of Biological Psychiatry*, 1-7. doi:10.3109/15622975.2015.1102325
- Siniatchkin, M., Glatthaar, N., von Mueller, G. G., Prehn-Kristensen, A., Wolff, S., Knoechel, S., . . . Gerber, W.-D. (2012). Behavioural treatment increases activity in the cognitive neuronal networks in children with attention deficit/hyperactivity disorder. *Brain Topography*, 25(3), 332-344. doi:10.1007/s10548-012-0221-6

- Smith, A. B., Taylor, E., Brammer, M., Toone, B., & Rubia, K. (2006). Task-specific hypoactivation in prefrontal and temporoparietal brain regions during motor inhibition and task switching in medication-naïve children and adolescents with attention deficit hyperactivity disorder. *American Journal of Psychiatry*, 163(6), 1044-1051. doi:10.1176/ajp.2006.163.6.1044
- Solomon, M., Yoon, J. H., Ragland, J. D., Niendam, T. A., Lesh, T. A., Fairbrother, W., & Carter, C. S. (2014). The development of the neural substrates of cognitive control in adolescents with autism spectrum disorders. *Biological Psychiatry*, 76(5), 412-421. doi:10.1016/j.biopsych.2013.08.036
- Spinelli, S., Joel, S., Nelson, T. E., Vasa, R. A., Pekar, J. J., & Mostofsky, S. H. (2011). Different neural patterns are associated with trials preceding inhibitory errors in children with and without attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 50(7), 705-715. doi:10.1016/j.jaac.2011.03.014
- Stevens, M. C., & Haney-Caron, E. (2012). Comparison of brain volume abnormalities between ADHD and conduct disorder in adolescence. *Journal of Psychiatry and Neuroscience*, 37(6), 389-398. doi:10.1503/jpn.110148
- Szekely, E., Schwantes-An, T. L., Justice, C. M., Sabourin, J. A., Jansen, P. R., Muetzel, R. L., . . . Shaw, P. (2018). Genetic associations with childhood brain growth, defined in two longitudinal cohorts. *Genetic Epidemiology*, 42(4), 405-414. doi:10.1002/gepi.22122
- Tamm, L., Menon, V., Ringel, J., & Reiss, A. L. (2004). Event-related fMRI evidence of frontotemporal involvement in aberrant response inhibition and task switching in attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43(11), 1430-1440. doi:10.1097/01.chi.0000140452.51205.8d
- Thornton, S., Bray, S., Langevin, L. M., & Dewey, D. (2018). Functional brain correlates of motor response inhibition in children with developmental coordination disorder and attention deficit/hyperactivity disorder. *Human Movement Science*, 59, 134-142. doi:10.1016/j.humov.2018.03.018
- Toal, F., Daly, E. M., Page, L., Deeley, Q., Hallahan, B., Bloemen, O., . . . Murphy, D. G. M. (2010). Clinical and anatomical heterogeneity in autistic spectrum disorder: A structural MRI study. *Psychological Medicine*, 40(7), 1171-1181. doi:10.1017/s0033291709991541
- Vaidya, C. J., Foss-Feig, J., Shook, D., Kaplan, L., Kenworthy, L., & Gaillard, W. D. (2011). Controlling attention to gaze and arrows in childhood: An fMRI study of typical development and autism spectrum disorders. *Developmental Science*, 14(4), 911-924. doi:10.1111/j.1467-7687.2011.01041.x
- van Hulst, B. M., de Zeeuw, P., Rijks, Y., Neggers, S. F. W., & Durston, S. (2017). What to expect and when to expect it: An fMRI study of expectancy in children with ADHD symptoms. *European Child & Adolescent Psychiatry*, 26(5), 583-590. doi:10.1007/s00787-016-0921-7
- van Rooij, D., Hoekstra, P. J., Mennes, M., von Rhein, D., Thissen, A. J. A. M., Hestfeldt, D., . . . Hartman, C. A. (2015). Distinguishing adolescents with ADHD from their unaffected siblings and healthy comparison subjects by neural activation patterns during response inhibition. *American Journal of Psychiatry*, 172(7), 674-683. doi:10.1176/appi.ajp.2014.13121635
- van Wingen, G. A., van den Brink, W., Veltman, D. J., Schmaal, L., Dom, G., Booij, J., & Crunelle, C. L. (2013). Reduced striatal brain volumes in non-medicated adult ADHD patients with comorbid cocaine dependence. *Drug Alcohol and Dependence*, 131(3), 198-203. doi:10.1016/j.drugalcdep.2013.05.007
- Velasquez, F., Qin, X. A., Reilly, M. A., Neuhaus, E., Estes, A., Aylward, E., & Kleinhaus, N. M. (2017). Neural correlates of emotional inhibitory control in autism spectrum disorders. *Research in Developmental Disabilities*, 64, 64-77. doi:10.1016/j.ridd.2017.03.008
- Vilgis, V., Sun, L., Chen, J., Silk, T. J., & Vance, A. (2016). Global and local grey matter reductions in boys with ADHD combined type and ADHD inattentive type. *Psychiatry Research: Neuroimaging*, 254, 119-126. doi:10.1016/j.psychresns.2016.06.008
- Villemonteix, T., De Brito, S. A., Kavec, M., Baleriaux, D., Metens, T., Slama, H., . . . Massat, I. (2015). Grey matter volumes in treatment naïve vs. chronically treated children with attention deficit/hyperactivity disorder: A combined approach. *European Neuropsychopharmacology*, 25(8), 1118-1127. doi:10.1016/j.euroneuro.2015.04.015

- Waiter, G. D., Williams, J. H. G., Murray, A. D., Gilchrist, A., Perrett, D. I., & Whiten, A. (2004). A voxel-based investigation of brain structure in male adolescents with autistic spectrum disorder. *NeuroImage*, 22(2), 619-625. doi:10.1016/j.neuroimage.2004.02.029
- Wang, J., Fu, K., Chen, L., Duan, X., Guo, X., Chen, H., . . . Chen, H. (2017). Increased gray matter volume and resting-state functional connectivity in somatosensory cortex and their relationship with autistic symptoms in young boys with autism spectrum disorder. *Frontiers in Physiology*, 8, 588. doi:10.3389/fphys.2017.00588
- Wang, J., Jiang, T., Cao, Q., & Wang, Y. (2007). Characterizing anatomic differences in boys with attention-deficit/hyperactivity disorder with the use of deformation-based morphometry. *American Journal of Neuroradiology*, 28(3), 543-547.
- Wilson, L. B., Tregellas, J. R., Hagerman, R. J., Rogers, S. J., & Rojas, D. C. (2009). A voxel-based morphometry comparison of regional gray matter between fragile X syndrome and autism. *Psychiatry Research: Neuroimaging*, 174(2), 138-145. doi:10.1016/j.psychresns.2009.04.013
- Yang, P., Wang, P.-N., Chuang, K.-H., Jong, Y.-J., Chao, T.-C., & Wu, M.-T. (2008). Absence of gender effect on children with attention-deficit/hyperactivity disorder as assessed by optimized voxel-based morphometry. *Psychiatry Research: Neuroimaging*, 164(3), 245-253. doi:10.1016/j.psychresns.2007.12.013
- Yang, Q., Huang, P., Li, C., Fang, P., Zhao, N., Nan, J., . . . Cui, L. B. (2018). Mapping alterations of gray matter volume and white matter integrity in children with autism spectrum disorder: Evidence from fMRI findings. *Neuroreport*, 29(14), 1188-1192. doi:10.1097/wnr.0000000000001094
- Yerys, B. E., Antezana, L., Weinblatt, R., Jankowski, K. F., Strang, J., Vaidya, C. J., . . . Kenworthy, L. (2015). Neural correlates of set-shifting in children with autism. *Autism Research*, 8(4), 386-397. doi:10.1002/aur.1454
- Zamorano, F., Billeke, P., Kausel, L., Larrain, J., Stecher, X., Hurtado, J. M., . . . Aboitiz, F. (2017). Lateral prefrontal activity as a compensatory strategy for deficits of cortical processing in attention deficit hyperactivity disorder. *Scientific Report*, 7(1), 7181. doi:10.1038/s41598-017-07681-z